

A mechanistic approach to antidiarrheal and analgesic potentials of active secondary metabolites from encapsulated probiotics.

Humaria Naz¹, Bashir Ahmad¹, Saima Naz¹, Khaliq Ur Rahman², Abdur Rauf^{3*}, Syed Uzair Ali Shah⁴, Sri Fatmawati⁵, Seema Patel⁶, Sami Bawazeer⁷

¹Center of Biotechnology and Microbiology, University of Peshawar, Peshawar-KPK-25120, KPK, Pakistan

²Department of Chemistry, University of Swabi, Anbar-23561, Khyber Pakhtunkhwa, Pakistan

³Department of Biotechnology Abdul Wali Khan University, Mardan, Mardan, 23200, Pakistan

⁴Department of Pharmacy, University of Swabi, Anbar-23561, Khyber Pakhtunkhwa, Pakistan

⁵Department of Chemistry, Faculty of Mathematics and Natural Sciences, Institut Teknologi Nopember, Surabaya, Indonesia

⁶Bioinformatics and Medical Informatics Research Center, San Diego State University, San Diego-92182, USA

⁷Department of EMS. Paramedic, College of Public Health and Health Informatics, Umm Al-Qura University, Makkah, Saudi Arabia

Abstract

The results obtained from the different fractions of metabolites were selected for *in vivo* acute toxicity studies. The toxicological study revealed that the test samples are safe for animal use up to 800 mg/kg. The test sample showed dose dependent anti-nociceptive potentials, the significant effect was observed with 500 and 800 mg/kg b.w dose. The analgesic effect of diclofenac sodium (standard drug) was greater than test sample. The significant ($P < 0.01$) anti-nociceptive potentials was noticed at 800 mg/kg b.w. of the animal for normal saline (35.50 ± 0.84), chloroform (29.67 ± 1.22), ethyl acetate (30.67 ± 1.22) and (29.00 ± 0.93) chloroform fraction of *L. plantarum* secondary metabolites. The test sample reduced the motility of GIT in mice. The reduction in GIT motility is dose dependent and maximum effect was produced by 800 mg/kg dose. The test sample caused different degree of movement of charcoal meal in the intestine i.e. 24.17, 20.17 and 17.83 cm at the tested doses of 200, 500 and 800 mg/kg, respectively.

Keywords: Encapsulated probiotics, Secondary metabolites, Analgesic, Antidiarrheal.

Accepted on September 27, 2017

Introduction

The term probiotic refers to dietary supplements containing beneficial bacteria. The bacteria such as *Lactobacillus* *Bifidobacterium* and yeasts such as *Saccharomyces* are well-known probiotics [1]. The oral consumption of probiotics exerts a protective effect on the gut flora and confers beneficial effects on microbial disorders of the gut, as indicated by multiple studies [2]. The best sources of probiotic bacteria are fermented dairy products. Probiotics sector is a rapidly-growing part of food industry, the annual global sales of probiotic products is expected to rise up to \$42 billion by 2018 [3,4].

The presence of probiotic bacteria in dairy products has long been established but the occurrence of *Lactobacillus* species in fruits and vegetables has rarely been focused [5]. Carbohydrates, minerals, nitrogenous compound and low pH

environment are necessary for the growth of probiotics which is present in fruit and vegetable juices and thus serves as excellent substrate for probiotics proliferation [6]. Of course, in the past few years, several vegetables have been used to isolate novel strains of probiotics. Studies show that the best source of probiotic bacteria are dairy fermented products, but they can also be isolated from non-dairy products such as vegetables, fruits and cereals. Usually probiotics isolated from non-dairy products are microbiologically safer than isolated from dairy product. Probiotics is a rapidly growing business domain and annual global sales of probiotic products is expected to rise up to \$42 billion by 2018 [7,8].

While going through the literature, *Vitis vinifera* (grapes) and *Spinacia oleracea* (spinach) are selected as a source of probiotic bacteria in the current study.

It may survive over winter in temperate region of Pakistan. *S. oleracea* used for chemoprevention of cervical cancer, apoptosis (cell death), liver damage and control of bleeding in jaundice. The *V. vinifera* belongs to family *Vitaceae* and in Pakistan over 77% are grown in Khyber Pakhtunkhwa. The microenvironment and nutritive profile of *V. vinifera* and *S. oleracea* are quite conducive for probiotic bacteria to multiply [9]. The *V. vinifera* traditionally used to treat constipation, indigestion, fatigue, kidney disorder and muscular degeneration. The presence of probiotic bacteria in dairy products has long been established but the occurrence of *Lactobacillus* species in fruits and vegetables has rarely been focused. Patel et al. have extensively reviewed the immunomodulatory and anticancer potential of probiotics [10].

Most of the species of lactic acid bacteria, which were isolated from raw or spontaneously fermented vegetables and fruits like cabbage juice, Tomatoes, marrows, carrots, cucumbers, eggplants, red-beets, capers, pineapple, plums, kiwi, papaya, fennels, cherries, cabbages, cereals, steamed rice [11]. In this study, *Vitis vinifera* (grapes) from the family *Vitaceae* and *Spinacia oleracea* (spinach) from the family *Amaranthaceae* are selected as a source of probiotic bacteria in the current study. The microenvironment and nutritive profile of *V. vinifera* and *S. oleracea* are quite conducive for probiotic bacteria to multiply [12]. So, it is highly relevant to investigate the *in vivo* pharmacological activities of such probiotics and its secondary metabolites. In the present study, we aimed to explore and evaluate bioactivities including acute toxicity, gastrointestinal motility test and analgesic activity of *Lactobacillus*'s secondary metabolites from fruits (*V. vinifera*) and vegetables (*S. oleracea*).

Experimental

Materials and methods

The selected probiotics source in this study was *V. vinifera* and *S. oleracea*, collected from Charsadda, Khyber Pakhtunkhwa, Pakistan. The samples were collected in clean plastic bags and carried to the laboratory for analysis within 24 hours of purchase.

Media preparation

The de Man, Rogosa and Sharpe (MRS) agar and broth were used for bacterial growth [12]. The pH of media was adjusted to 6-7 by using 0.1 N NaOH and 0.1N HCl.

Isolation of probiotics

The grapes and spinach samples, carried to the laboratory were washed with water followed by a final rinse with distilled water. The samples were treated independently in sterile Petri plates. The media was autoclaved and then poured into sterile plates and allowed to solidify. One mL of *V. vinifera* and *S. oleracea* extract obtained was inoculated into 9 mL of sterile MRS broth were kept for 48 hours at 37°C at 120 rpm in shaking incubator. Sample was then isolated, diluted and

separated on MRS agar plate. Colonies were purified by sub-culturing on MRS agar plates and further characterized [13].

Characterization of selected isolates

The cultures were identified on the basis of criteria suggested by Harrigan [13].

Extraction of secondary metabolites

MRS broth media were prepared in 1 L of distilled water and 100 mL of it was poured in each 250 ml Erlenmeyer flasks. The flasks were plugged and autoclaved. The culture flasks were incubated in shaking incubator at 28°C for 7 d at 180 rpm. After incubation, few drops of HCl were poured into flasks and shaken vigorously to settle the media and increase the polarity of the aqueous phase. Media was filtered through filter paper and the filtrate was taken. Equal volume of ethyl acetate (EtOAc) and chloroform (CHCl₃) was added to the filtrate, and then shaken vigorously for about 20 minutes in a separating funnel. The organic phase was separated and washed with 2 M brine solution followed by purification with anhydrous sodium sulfate (Na₂SO₄). The organic phase containing the metabolites was dried in a rotary evaporator at 45°C and 150 rpm. The extract was taken in glass vials from evaporator by adding few ml of methanol, followed by filtration through a filter of 0.2 µm of mesh size. The filtrate was dried in a water bath and extract was then used for pharmacological assays [14].

Test animals

For the experiment, Adult Swiss albino mice (BALB/c) from either sex, weighing between (12-30) g, were collected from animal sources department NIH (National Institute of Health, Islamabad). The animals were maintained under normal laboratory conditions and kept in standard polypropylene cages at room temperature of 35 ± 2°C and relative humidity. The Swiss Albino (BALB/c) mice of either sex weighing 25±10 g were used for determining the acute toxicity, GIT motility test, and analgesic test.

Acute toxicity assay

The acute toxicity assay was performed according to the reported procedure [15]. Animals were divided into six groups (n=6). The secondary metabolites from the microbial isolates of the *V. vinifera* and *S. oleracea* samples at 200, 500, 800 and 1500 mg/kg (body weight of mice) was used as test dose. After intraperitoneally (i.p.) administering the test doses, the animals were kept under observation for 24 h. The number of survived and dead animals was noted by calculating percent mortality. The groups received the following treatments. Antidiarrheal/gastrointestinal motility test (GIT motility).

The GIT motility test was performed according to Marona and Lucchesi [14], with slight modifications where necessary. All the mice were kept in food deprivation for 3 h but only accessible to water and they were divided in six groups, each group containing 6 mice (n=6). Then the mice were weighed

A mechanistic approach to antidiarrheal and analgesic potentials of active secondary metabolites from encapsulated probiotics

and marked. Control solution (5 ml/kg), sample solutions of 200 mg/kg, 500 mg/kg and 800 mg/kg of all the extracts were administered by oral gavage. After 90 min, 0.3 ml of charcoal suspension was administered to all the mice. After 60 min mice were provided with free access to food. Then the mice were observed at 5 min intervals until feces with charcoal were eliminated. Charcoal was observed on the feces using normal light when it was easily visible, or using a microscope to help the identification of the black spots.

Antinociceptive activity

The acetic acid induced writhing test was used to know about analgesic effect of secondary metabolites of probiotics [15,22]. The selected test samples were screened for the presence of analgesic effect. These animals were divided into five different groups (n=6). Group I and II were served as negative and positive control, respectively. Normal saline at a dose of 10 ml/kg (body weight) was administered to Group I, while Diclofenac sodium at a dose of 10 mg/kg (body weight) was administered to Group II. They were fed according to recommended guidelines, but food supply was stopped 2 h before the start of activity [15,16]. The crude extract was administered to the remaining groups (III, IV and V) at a dose of 200, 500 and 800 mg/kg (body weight) intraperitoneally (i.p.). After 30 min of the above-mentioned treatments 1% acetic acid was administered to all groups through intraperitoneal route. Then after 5 min of acetic acid injection abdominal writhes (constrictions) were started to count for next 10 min. The percent analgesic effect was calculated as per given formula.

% Protection

$$= \frac{\text{No. of writhing in control} - \text{No. of writhing in test compound}}{\text{No. of writhing in control}} \times 10$$

Statistical analyses

Data was expressed as Mean \pm SEM (Standard error of Mean). The results were analyzed statistically by ANOVA followed by Dunnett's test. Results below $p < 0.05$ and $p < 0.01$ were considered statistically significant.

Results

Acute toxicity

The animals were observed for 24 h for mortality, but no dose was found to be toxic and all doses were found to be safe. Therefore, the test samples were considered to be relatively safe up to a dose of 200-1500 mg/kg body weight of mice. The results are shown in Table 1.

Antinociceptive activity

This study showed that EtOAc and CHCl_3 fractions of *L. plantarum* and *L. acidophilus* possess analgesic activity. The results showed that the pain relief was achieved in a dose dependent manner as shown in table 2. The dominant and

significant ($P < 0.01$) analgesic action was noticed at dose of 800 mg/kg, which was (29.67 ± 1.22) for CHCl_3 fraction of *L. plantarum*, (30.67 ± 1.22) for EtOAc fraction of *L. acidophilus* and (29.00 ± 0.93) CHCl_3 fraction of *L. acidophilus*. The percent inhibition of writhing is shown in Figure 1. The inhibitory effect of diclofenac sodium was greater than that of the highest dose of test samples of all fractions.

Table 1. Acute toxicity of the EtOAc, CHCl_3 fractions of *L. plantarum* and *L. acidophilus*.

Test samples	Dose (mg/kg)	Gross effect after 4 h	Mortality after 24 h
Test sample	200	-	-
	500	-	-
	800	-	-
	1500	-	-




Figure 1. Percent Analgesic activity of (a) CHCl_3 fraction of *L. plantarum* (b) CHCl_3 fraction of *L. acidophilus* (c) EtOAc fraction of *L. acidophilus* secondary metabolites. Bars present as mean \pm S.E.M. (n=6). ANOVA followed by Dunnett's test was applied on the data to find the level of significance in comparison with negative control.

S. No	Treatment	Dose mg/kg	ml or No. of writhing (10 min) (Mean+SEM)
1	Normal saline	10	52.00 \pm 1.59
2	<i>L. plantarum</i> CHCl_3	200	47.83 \pm 1.53
		500	41.33 \pm 1.29
		800	29.67 \pm 1.22**
3	<i>L. acidophilus</i> CHCl_3	200	44.67 \pm 0.91
		500	31.50 \pm 0.80
		800	29.00 \pm 0.93
4	EtOAc	200	50.17 \pm 1.20
		500	40.00 \pm 1.37
		800	30.67 \pm 1.22**
5	STD (Diclofenac sod.)	10	18.00 \pm 1.39

Table 2. Analgesic activity CHCl_3 and EtOAc fractions of *L. plantarum* and *L. acidophilus* secondary metabolites.

GIT motility test

The entire test sample caused a significant movement of charcoal meal in the intestine of mice results. The percent movement of charcoal meal through the small intestine after treatment with CHCl₃ fraction of *L. plantarum* secondary metabolites was 26.50, 24.00 and 20.33 cm, at the doses of 200, 500 and 800 mg/kg, respectively. In saline-treated groups, the charcoal meal travelled 41.50. The percent GIT motility of CHCl₃ fraction of *L. plantarum* secondary metabolites 24.17, 20.17 and 17.83, respectively at (200, 500 and 800 mg b.w) and atropine sulfate assessed by charcoal propulsion. GIT Motility of CHCl₃ fraction of *L. acidophilus* secondary metabolites 26.83, 22.17 and 13.83 respectively at (200, 500 and 800 mg b.w) (Table3).

Table 5. The effect of CHCl₃ and EtOAc fractions of *L. plantarum* and *L. acidophilus* on GIT Motility.

S. No	Treatment	Dose ml or mg/kg	Mean total length of intestine (cm)	Mean charcoal movement	
1	Normal saline	10	49.00 ± 0.80	35.50 ± 0.84	
2	L. plantarum	CHCl ₃	200	53.00 ± 0.37	26.50 ± 0.56
			500	53.00 ± 24.00 ± 0.68 0 0.35	
			800	50.00 ± 20.33 ± 0.80 0 0.33	
3	L. acidophilus	CHCl ₃	200	45.00 ± 0.57	26.83 ± 1.19
			500	47.00 ± 22.17 ± 0.68 0 0.37	
			800	48.00 ± 13.83 ± 1.01 0 0.57	
4	EtOAc	200	47.00 ± 24.17 ± 1.24 0 0.57		
			500		47.00 ± 20.17 ± 0.47 0 0.37
			800		45.00 ± 17.83 ± 0.97 0 0.57

Discussion

The present study mainly focused on the isolation of *Lactobacillus*'s secondary metabolites from fruits (*V. vinifera*) and vegetables (*S. oleracea*) and to confirm their use for the management of pain and diarrhea on scientific basis. The various strains of *Lactobacillus* have already been isolated and characterized from dairy sources but literature is limited regarding fruits and vegetables [17]. Therefore, we tried to explore the potential of *Lactobacillus* from *V. vinifera* and *S. oleracea*. The MRS broth and agar media is used for initial culturing, culture enrichment of *Lactobacillus* sp. and then they

were characterized as suggested by Harington [12]. We isolated the *L. plantarum* and *L. acidophilus* from broth culture. Furthermore, we fractionated these cultures with chloroform (CHCl₃) and ethylacetate (EtOAc) for its secondary metabolites, which were finally assessed for analgesic and antidiarrheal effect.

The acetic acid induced pain model is a common, rapid, sensitive and easy method for the determination of peripheral analgesic effect of plant extracts and other drugs as well [18]. The chemicals produced in peritoneal fluids are responsible for pain which may appear in the form of abdominal constrictions. Writhing inhibition by various substances may be involved in the decreased production or inhibition of prostanooids, which is considered for pain inhibition through peripheral mechanism [19,20]. It is suggested that the test sample its peripheral analgesic effect through inhibition of abdominal receptors. The active chemical(s) present in test sample produced analgesic effect in the form of reduction in abdominal constrictions (writhing). Therefore, we evaluated the analgesic potential of our test fractions by using acetic acid induced writhing test. Our data showed analgesic potential of probiotics and their fractions of secondary metabolites to much or less extent in all animal models used. The analgesic effect of fractions was found to be dose dependent i.e., maximum at dose of 800 mg/kg, however, all the fraction exhibited less analgesic potential than produced by standard drug i.e. diclofenac. The predominant analgesic effect among all fractions was exhibited by CHCl₃ fraction of *L. plantarum*. Previous studies have reported analgesic, antidiarrheal and anti-inflammatory potentials of probiotics [21] other several flavone derivatives [22].

Multiple comparison tests revealed that *Lactobacillus* fractions at concentration of 800 mg/kg significantly reduced the GIT motility. The test sample may cause relaxation of intestine by interaction with M3 receptor found in small intestine, where acetylcholine can act and activation of M3 receptors will lead to increase contraction [20,21]. The effect of antispasmodic agents is to alleviate the symptoms of abdominal cramps. The test samples reduce GIT motility and can be used for the management of diarrhea and in abdominal spasm.

Conclusion

In the present study, we concluded that beneficial *Lactobacillus* strains can be isolated from fruits (*V. vinifera*) and vegetables (*S. oleracea*) like *L. plantarum* and *L. acidophilus*. These *lactobacillus* strains produced secondary metabolites which are effective to produce analgesia and alter GIT motility.

References

1. Weissinger WR, Chantarapanont W, Beuchat LR. Survival and growth of Salmonella baidon in shredded lettuce and diced tomatoes, and effectiveness of chlorinated water as a sanitizer. Int J Food Microbiol 2000; 62. 123-13.

A mechanistic approach to antidiarrheal and analgesic potentials of active secondary metabolites from encapsulated probiotics

2. Trias R, Bañeras L, Montesinos E, Badosa E. Lactic acid bacteria from fresh fruit and vegetables as biocontrol agents of phytopathogenic bacteria and fungi. *Int Microbiol* 2008; 11: 231-236.
3. Alvarez-Olmos MI, Oberhelman RA. Probiotic agents in infectious diseases: A modern perspective on a traditional therapy. *Clin Infec Dis* 2001; 32: 1567-1576.
4. Salminen S. Probiotics: scientific support for use. *Food Technol* 1999; 53: 66.
5. Chambel L, Chelo IM, Zé-Zé L, Pedro LG, Santos MA, Tenreiro R. *Leuconostoc pseudoficulneum* sp. nov., isolated from a ripe fig. *Int J Syst Evol Microbiol* 2006; 56: 1375-1381.
6. Gill HS, Guarner F. Probiotics and human health: a clinical perspective. *Postgrad Med J* 2004; 80: 516-526.
7. Naidu AS, Bidlack WR, Clemens RA. Probiotic spectra of lactic acid bacteria (LAB). *Crit Rev Food Sci Nutr* 1999; 39: 13-126.
8. Bae S, Fleet GH, Heard GM. Lactic acid bacteria associated with wine grapes from several Australian vineyards. *J Appl Microbiol* 2006; 100: 712-727.
9. Quewand AC, Salminen SJ. The health effects of cultured milk products with viable and non-viable bacteria. *Int Dairy J* 1998; 8:749-758.
10. Patel S, Shukla R, Goyal A. Probiotics in valorization of innate immunity across various animal models. *J functional foods* 2015; 14, 549-561
11. Patel S, Goyal A. Evolving roles of probiotics in cancer prophylaxis and therapy. *Probiotics Antimicrob Proteins* 2013; 5: 59-67.
12. Chambel L, Chelo IM, Zé-Zé L, Pedro LG, Santos MA. *Leuconostoc pseudoficulneum* sp. nov., isolated from a ripe fig. *Int J Syst Evol Microbiol* 2006; 56: 1375-1381.
13. Harrigan WF. *Laboratory method in food microbiology*. Gulf professional publishing, 1998.
14. Marona HR, Lucchesi MB. Protocol to refine intestinal motility test in mice. *Lab Anim* 2004; 38: 257-260.
15. Bashir S, Alam M, Ahmad B, Aman A, Ali J. Screening of *Ferula narthex* Boiss Crude Methanolic Extract for Analgesic, Gastrointestinal Motility and Insecticidal Activity. *Middle-East J Sci Res* 2013; 14: 471-475.
16. Rastogi S, Pandey MM, Rawat AK. An ethno-medicinal, phytochemical and pharmacological profile of *Desmodium gangeticum* (L.) DC. and *Desmodium adscendens* (Sw.) DC. *J ethnopharmacol* 2011; 136: 283-296
17. Wedajo B. Lactic Acid Bacteria: Benefits, Selection Criteria and Probiotic Potential in Fermented Food. *J. Prob Health* 2015; 3:129
18. Bentley GA, Newton SH, Starr J. Studies on nociceptive action of a-agonist drugs and their interactions with opioids mechanisms. *Br J Pharmacol* 1983; 79:125
19. Duarte ID, Nakamura M, Ferreira SH. Participation of the sympathetic system in acetic acid-induced writhing in mice. *Braz J Med Biol Res* 1988; 21: 341-343.
20. Khan H, Saeed M, Gilani AU, Khan MA, Dar A. The antinociceptive activity of *Polygonatum verticillatum* rhizomes in pain models. *J Ethnopharmacol* 2010; 127: 521-527.
21. Quigley EM. Probiotics in irritable bowel syndrome: an immunomodulatory strategy? *J Am Coll Nutr* 2007; 26: 684S-90S.
22. Sannigrahi S, Mazumder UK, Pal D, Mishra ML, Maity S. flavonoids of *enhydra fluctuans* exhibits analgesic and anti-inflammatory activity in different animal models, *Pak. J Pharm Sci* 2011; 24: 369-375.

***Correspondence to**

Abdur Rauf

Head Department of Chemistry

University of Swabi

Anbar

Khyber Pakhtunkhwa

Pakistan

E-mail: mashaljcs@yahoo.com

Tel Number: +923469488944