

## Intermittent low-magnitude high-frequency vibration promotes osteogenic protein expression and inhibits osteoclastogenic cytokine secretion in osteoblasts.

Bin Zhao<sup>1#</sup>, Ruixin Li<sup>2#</sup>, Shihuan Cheng<sup>3</sup>, Xin Dong<sup>4</sup>, Lei Tan<sup>1</sup>, Yuying Chen<sup>5</sup>, Baohui Hao<sup>1</sup>, Dong Zhu<sup>1\*</sup>

<sup>1</sup>Department of Orthopedic Traumatology, First Hospital of Jilin University, Changchun, PR China

<sup>2</sup>Institute of Medical Equipment, the Academy of Military Medical Sciences, Tianjin, PR China

<sup>3</sup>Department of Rehabilitation Medicine, First Hospital of Jilin University, Changchun, PR China

<sup>4</sup>Medical Record Department, Chinese People's Liberation Army 461 Hospital, Changchun, PR China

<sup>5</sup>The Second Affiliated Hospital of Dalian Medical University, Dalian, PR China

#These authors contributed equally to this work

### Abstract

**Background:** Accumulating evidence has indicated that Low-Magnitude High-Frequency Vibration (LMHFV) can promote fracture healing, and thus, holds promise as a treatment for osteoporosis. However, few studies have reported the effects of LMHFV, especially intermittent LMHFV (iLMHFV) on osteoblasts *in vitro*. In this study, we examined the effects of sostenuto and intermittent LMHFV (sLMHFV and iLMHFV, respectively) on MC3T3-E1 preosteoblast proliferation and production of osteogenic and osteoclastogenic proteins.

**Methods:** The cell proliferation was determined using MTT assay for up to 7 d. The expression levels of the osteogenic proteins Alkaline Phosphatase (ALP), Osteocalcin (OCN), and Collagen type I (COL I), and the osteoclastogenic protein nuclear factor- $\kappa$ B ligand (RANKL) and its inhibitor Osteoprotegerin (OPG) were determined by Real time PCR and/or Western blot analysis. The ALP activity was determined using a fluorometric assay. The secretion of osteoclastogenic cytokines RANKL and M-CSF was assessed by ELISA.

**Results:** Our results showed that neither sLMHFV nor iLMHFV affected MC3T3-E1 cell proliferation. However, both sLMHFV and iLMHFV increased the expression of the osteogenic proteins ALP, OCN, and COL I. SLMHFV and iLMHFV also inhibited the secretion of the osteoclastogenic cytokines RANKL and M-CSF and decreased the OPG/RANKL ratio. These osteogenic and anti-osteoclastogenic effects were most profound with short iLMHFV.

**Conclusion:** These finding could eventually contribute to the development of potential treatment regimen for osteoporosis using LMHFV.

**Keywords:** Sostenuto low-magnitude high-frequency vibration, Intermittent low-magnitude high-frequency vibration, Osteoblast, Osteogenic proteins, Osteoclastogenic cytokines.

Accepted on December 1, 2017

### Introduction

Bone fracture is the most common injury in traffic accidents and falls, especially in elderly patients with osteoporosis [1,2]. 5%-10% of all patients with bone fracture often suffer from delayed unions or non-unions, which may result in long-term disability [3]. There are about 8.9 million of osteoporosis-related fractures each year globally, which brings a great burden to public health [2].

Bone homeostasis is an intricate balance between bone formation by osteoblasts and bone resorption by osteoclasts. During the process of fracture healing, osteoblasts must proliferate, differentiate, and induce mineralization within bone extracellular matrix. Osteoblast differentiation plays a vital role during bone fracture healing. This process is characterized by increased expression of osteogenic proteins such as Alkaline Phosphatase (ALP), Osteocalcin (OCN), and Collagen type I (COL I). Mechanical forces can affect bone formation and

repair through regulation of osteoblast differentiation and the intricate interplay between osteoblasts and osteoclasts [4-6]. Osteoblasts regulate osteoclast differentiation by producing osteoclastogenic cytokines including Macrophage Colony-Stimulating Factor (M-CSF) and nuclear factor- $\kappa$ B ligand (RANKL). Mechanical forces can also inhibit the production of these osteoblast-derived osteoclastogenic cytokines, thereby slowing down the process of bone resorption [5,7-9].

Sostenuto Low-Magnitude High-Frequency Vibration (sLMHFV) in the range of 0.2-0.3 g and 32-37 Hz increases bone density and prevent bone from developing osteoporosis [10-13]. It also promotes closed femoral fracture healing [14]. In agreement with these *in vivo* effects, sLMHFV (0.3 g, 40 Hz, 30 min/12 h) increases osteoblast differentiation, matrix synthesis, and mineralization *in vitro* [15,16]. Interestingly, studies have also shown that intermittent LMHFV (iLMHFV) is more effective than sLMHFV in improving the biomechanical properties and microstructure of bone tissue in ovariectomized rats [17]. However, there are few reports on the effects of iLMHFV on osteoblasts *in vitro*.

In this study, we examined the effects of sLMHFV and iLMHFV on osteoblast proliferation, osteogenic protein expression, and osteoclastogenic cytokine production *in vitro*. We found that iLMHFV was more effective than sLMHFV in promoting osteogenic protein expression and inhibiting osteoclastogenic cytokine secretion. Moreover, short iLMHFV can provide optimal osteogenic and anti-osteoclastogenic effects on osteoblasts.

## Materials and Methods

### MC3T3-E1 cell culture

The mouse preosteoblast MC3T3-E1 cell line was obtained from the Cell Resource Center IBMS CAMS/PUMS (Beijing, China). The cells were cultured in modified Eagles minimum essential medium ( $\alpha$ -MEM; Invitrogen, USA) supplemented with 10% Fetal Bovine Serum (FBS) and 1% penicillin-streptomycin. The cells were maintained at 37°C, 5% CO<sub>2</sub> in a humidified incubator. After having reached around 90% confluence, the cells were harvested by treatment with 10% trypsin-EDTA (Sigma), seeded in six-well plates, and cultured for another 24 h prior to vibration treatment.

### LMHFV treatment

The MC3T3-E1 cells were subjected to sLMHFV or iLMHFV (0.25 g, 35 Hz) treatment on a six-well vibration table (DH40020) manufactured at Hebei University of Technology. For studies on osteoblast phenotypic gene expression, we set up six groups: Control, sostenuto vibration 0.5 h (SV0.5 h), vibration 15 minX2-interval 0.5 h (IV0.5 h), vibration 15 minX2-interval 1 h (IV1 h), vibration 15 minX2-interval 2 h (IV2 h), vibration 15 minX2-interval 3 h (IV3 h), vibration 15 minX2-interval 6 h (IV6 h). Untreated cells were used as negative control. For studies on osteoblast proliferation, the cells were treated each day with sostenuto vibration or

intermittent vibration as above for a total of 1, 3, 5, or 7 d. Untreated cells were used as negative control.

### The MTT assay

The effects of iLMHFV and sLMHFV treatment on MC3T3-E1 cell proliferation were determined using the Methyl Thiazolyltetrazolium (MTT) assay. Briefly, MC3T3-E1 cells were cultured for another 24 h after vibration treatment. MTT (Promega) solution (5 mg/ml, 1 ml/well) was added, and the cells were incubated for 4 h. After that, the supernatant was removed, and 2 ml Dimethyl Sulfoxide (DMSO) was added to dissolve the formazan crystals formed. Absorbance at 490 nm was recorded by a scanning multiwell spectrophotometer (iMark, Bio-Rad, US). Each experiment was performed in triplicate.

### Alkaline phosphatase (ALP) activity analysis

After vibration treatment, MC3T3-E1 cells were harvested and lysed in lysis buffer (10 mM HEPES, 250 mM sucrose, 5 mM Tris-HCl, 0.1% TritonX-100, pH 7.5) with short bursts of sonication. The cell lysates were centrifuged at 4°C overnight. The ALP activity in the supernatant was determined using a fluorometric detection kit (Nanjing KeyGENBioTECH Co. Ltd, Nanjing, China). Relative ALP units were normalized to sample protein concentration. Each experiment was performed in triplicate.

### RNA extraction and real time PCR

MC3T3-E1 cells were harvested after vibration treatment. Total RNA was extracted using Trizol reagent (Beijing Biosntech Co. Ltd, Beijing China). RNA (10  $\mu$ l) was reverse transcribed using random hexamer primers and transcriptase from the Superscript III Reverse Transcriptase Kit (TIANGEN Biotech (Beijing) Co., Ltd Beijing China). Real time PCR was carried out on an IQ5 Bio-rad Real-time PCR System following manufacturer's instructions. Primer sequences are shown in Table 1. Relative mRNA expression was calculated using the  $\Delta\Delta C(t)$  method [18] and normalized to  $\beta$ -actin. Each experiment was performed in triplicate.

### Western blot analysis

MC3T3-E1 cells were harvested after vibration treatment. Total cell lysates were prepared in RIPA lysis buffer containing protease and phosphatase inhibitors. Nuclear lysates were prepared as previously described [19]. Protein concentration was determined using the BCA method. Samples (20  $\mu$ g proteins) were separated by SDS-PAGE and transferred to PVDF membranes (Millipore, USA). After blocking in 5% skim milk, the membranes were incubated overnight with anti-OCN and anti-Col I antibodies (Santa Cruz, USA), respectively, at 4°C. After washing, the membranes were incubated with horseradish peroxidase-conjugated secondary antibody. The protein bands were visualized using enhanced chemiluminescence detection kit (TIANGEN BIOTECH CO., LTD, Beijing, China). Relative protein expression was

## Intermittent low-magnitude high-frequency vibration promotes osteogenic protein expression and inhibits osteoclastogenic cytokine secretion in osteoblasts

normalized to GAPDH and presented as fold of control. Each experiment was performed in triplicate.

### ELISA

Secretion of osteoclastogenic cytokines RANKL and M-CSF were assessed by enzyme-linked immunosorbent assay (ELISA). After vibration treatment, RANKL and M-CSF levels in the MC3T3-E1 cell culture medium were determined using RANKL and M-CSF ELISA kits (Cloud-Clone Corp., USA), respectively, following manufacturer's instructions. Data were normalized to total protein content. Each experiment was performed in triplicate.


### Statistical analysis

Statistical analysis was performed using SPSS13.0 software. All results are presented as mean  $\pm$  S.D. (Standard Deviation). Data were interpreted for normal distribution using Shapiro-Wild test. Multiple comparisons were analysed using one-way ANOVA followed by Duncan's multiple range test. Differences with a P-value less than 0.05 were considered statistically significant.

## Results

### Neither LMHFV nor iLMHFV affects MC3T3-E1 cell proliferation


First, we examined the effects of sLMHFV and iLMHFV on MC3T3-E1 cell proliferation for up to 7 d. Each day, the cells were treated with sostenuto vibration for 0.5 h (SV0.5 h) or vibration 15 minX2-interval 0.5 h (IV0.5 h), vibration 15 minX2-interval 1 h (IV1 h), vibration 15 minX2-interval 2 h (IV2 h), vibration 15 minX2-interval 3 h (IV3 h), vibration 15 minX2-interval 6 h (IV6 h). Our MTT assay results indicated that sLMHFV or iLMHFV-treated cells proliferated at the same rate as the untreated cells throughout the experiment (Figure 1). Thus, sLMHFV and iLMHFV does not have obvious effects on osteoblast proliferation.



**Figure 1.** Neither sLMHFV nor iLMHFV affects MC3T3-E1 cell proliferation. MC3T3-E1 cells were treated each day with sLMHFV for 0.5 h or iLMHFV for 0.5 h, 1 h, 3 h, or 6 h, for a total of 1 d, 3 d, 5 d, and 7 d, respectively. Cell proliferation was determined by the MTT assay.  $n=3$ .

### sLMHFV and iLMHFV increase ALP mRNA expression level and enzymatic activity in MC3T3-E1 cells

Alkaline Phosphatase (ALP), a key enzyme that provides high concentrations of phosphate at the site of mineral deposition, is a key factor of osteoblast early differentiation [20]. We examined the effects of sLMHFV and iLMHFV on ALP mRNA expression level and enzymatic activity in MC3T3-E1 cells. The cells were treated with sostenuto vibration for 0.5 h or intermittent vibration for 0.5 h, 1 h, 3 h, or 6 h. The ALP mRNA levels and enzymatic activity were determined by real time PCR and a fluorometric ALP assay, respectively. As shown in Figure 2A, both sLMHFV and iLMHFV-treated cells displayed significantly higher ALP mRNA levels compared to untreated cells. Interestingly, 0.5 h or 1 h intermittent vibration was significantly more effective in increasing ALP expression than 0.5 h constant vibration with 0.5 h group 110% more and 1h group 20% more, respectively. Moreover, 0.5 h intermittent vibration appeared to be optimal, with longer vibration being less effective. Consistent with the mRNA expression results, the ALP enzymatic activity was significantly enhanced by both sLMHFV and iLMHFV treatment, with iLMHFV demonstrating more robust effects (Figure 2B).




**Figure 2.** Both sLMHFV and iLMHFV significantly increase ALP mRNA expression and activity. MC3T3-E1 cells were treated with sLMHFV for 0.5 h or iLMHFV for 0.5 h, 1 h, 3 h, or 6 h. The ALP mRNA expression (A) and enzymatic activity (B) were determined by real time PCR and a fluorometric ALP assay, respectively.  $n=3$ , \* $P<0.05$  versus Control; # $P<0.05$  versus SV0.5 h.


### sLMHFV and iLMHFV induce OCN and COL I expression in MC3T3-E1 cells

We subsequently examined the effects of sLMHFV and iLMHFV on the mRNA and protein expression level of the osteogenic proteins OCN and COL I, the key markers of osteoblast differentiation at later stage. OCN, a protein secreted solely by osteoblasts, is present in bone mineralization and calcium ion homeostasis. COL I is the main component of bone ECM (Extra Cellular Matrix). MC3T3-E1 cells were treated with sostenuto vibration for 0.5 h or intermittent vibration for 0.5 h, 1 h, 3 h, or 6 h. The OCN and COL I mRNA and protein levels were determined by real time PCR and Western blot analysis, respectively. As shown in Figures 3A and 4A, the mRNA levels of OCN and COL I in MC3T3-E1 cells significantly increased after sLMHFV or iLMHFV treatment. Moreover, intermittent vibration (OCN: 0.5 h, 1 h, or 3 h, COL I: 0.5 h) was superior to constant vibration (0.5 h) in inducing

these osteogenic proteins. Consistent with these mRNA results, sLMHFV and iLMHFV treatment significantly elevated OCN and COL I protein levels in MC3T3-E1 cells, with iLMHFV demonstrating more robust effects (Figures 3B and 3C and 4B and 4C). Similar to what was observed with ALP, the optimal effect on OCN and COL I was obtained with 0.5 h intermittent vibration.



**Figure 3.** Both sLMHFV and iLMHFV significantly increase OCN mRNA and protein expression. MC3T3-E1 cells were treated with sLMHFV for 0.5 h or iLMHFV for 0.5 h, 1 h, 3 h, or 6 h. The OCN mRNA (A) and protein expression (B and C) were determined by real time PCR and Western blot analysis, respectively. n=3, \*P<0.05 versus control, #P<0.05 versus SV0.5 h.




**Figure 4.** Both sLMHFV and iLMHFV significantly increase COL I mRNA and protein expression. MC3T3-E1 cells were treated with sLMHFV for 0.5 h or iLMHFV for 0.5 h, 1 h, 3 h, or 6 h. The COL I mRNA (A) and protein expression (B and C) were determined by real time PCR and Western blot analysis, respectively. n=3, \*P<0.05 versus control, #P<0.05 versus SV0.5 h.

**SLMHFV and iLMHFV increase the OPG/RANKL ratio in MC3T3-E1 cells**

Osteoblasts regulate osteoclast differentiation through production of M-CSF, RANKL, and OPG. OPG is a decoy receptor for RANKL, which binds to RANKL blocks the RANKL-RANK interaction between osteoblast and osteoclast precursors, thereby inhibiting osteoclastogenesis. The ratio of OPG/RANKL determines osteoclast differentiation and consequently the bone resorption rate [14]. We examined the effects of sLMHFV and iLMHFV on the OPG/RANKL ratio in MC3T3-E1 cells. The cells were treated with sostenuto vibration for 0.5 h or intermittent vibration for 0.5 h, 1 h, 3 h, or 6 h. The OPG and RANKL mRNA levels were determined by real time PCR and the OPG/RANKL ratios were subsequently calculated. As shown in Figure 5, sLMHFV and iLMHFV treatment significantly increased the OPG/RANKL

ratio in MC3T3-E1 cells, suggesting that these treatments inhibit the osteoclastogenic potential of osteoblasts. Similar to what was observed with osteogenic protein expression, iLMHFV (0.5 h, 1 h, or 3 h) treatment outperformed sLMHFV (0.5 h) in increasing the OPG/RANKL ratio, with 0.5 h, 1 h, and 3 h has 124%, 105%, and 96% higher than sLMHFV (0.5 h), and the optimal effects were obtained with 0.5 h intermittent vibration.



**Figure 5.** SLMHFV and iLMHFV significantly increase the OPG/RANKL ratio. MC3T3-E1 cells were treated with sostenuto vibration for 0.5 h or intermittent vibration for 0.5 h, 1 h, 3 h, or 6 h. The OPG and RANKL mRNA levels were determined by real time PCR, and the OPG/RANKL ratios were subsequently calculated. n=3, \*P<0.05 versus Control, #P<0.05 versus SV0.5 h.

**sLMHFV and iLMHFV inhibit M-CSF and RANKL secretion by MC3T3-E1 cells**

It has been reported that mechanical loading can suppress osteoclast activation through inhibition of the production of osteoblast-derived RANKL and M-CSF [5,7,8,14]. In this study, we examined the effects of sLMHFV and iLMHFV on M-CSF and RANKL secretion by MC3T3-E1 cells. The cells were treated with sostenuto vibration for 0.5 h or intermittent vibration for 0.5 h, 1 h, 3 h, or 6 h. The secreted RANKL and M-CSF protein in the culture medium were determined by ELISA. Our results revealed that both sLMHFV and iLMHFV treatment led to significantly decreased M-CSF and RANKL concentrations in the culture medium (Table 2), suggesting that these treatments suppress osteoblast-mediated osteoclast activation. In addition, iLMHFV (0.5 h or 1 h) outperformed sLMHFV (0.5 h) in inhibiting M-CSF and RANKL secretion, and the optimal inhibitory effects on both cytokines were obtained with 0.5 h intermittent vibration.

MC3T3-E1 cells were treated with sostenuto vibration for 0.5 h or intermittent vibration for 0.5 h, 1 h, 3 h, or 6 h. The RANKL and M-CSF protein concentrations in the culture medium were determined by ELISA. n=3, \*P<0.05 versus control, #P<0.05 versus SV0.5 h.

**Table 1.** Primer sequences for real time PCR.

Gene (mouse)	Forward	Reverse
ALP	GAAGGAGGCAGGATTGAC	ATCAGCAGTAACCACAGTC

*Intermittent low-magnitude high-frequency vibration promotes osteogenic protein expression and inhibits osteoclastogenic cytokine secretion in osteoblasts*

OCN	CCATCTTTCTGCTCACTCT	ACTACCTTATTGCCCTCCT	RANKL	ATCAGAAGACAGCACTCAC	GGCTTGTTTCATCCTCCT
COL1	CAGTGGCGGTTATGACTT	CTGCGGATGTTCTCAATCT	$\beta$ -actin	GATTACTGCTCTGGCTCCT	ACTCCTGCTTGCTGATCC
OPG	GCCCTGACCACTCTTATAC	AACGCCCTTCCTCACACT			

**Table 2.** sLMHFV and iLMHFV inhibit M-CSF and RANKL secretion by MC3T3-E1 cells.

Cytokines	Control	SV0.5 h	IV0.5 h	IV1 h	IV3 h	IV6 h
RANKL (pg/ml)	273 ± 15	263 ± 15*	200 ± 18* <sup>#</sup>	231 ± 13* <sup>#</sup>	274 ± 13	277 ± 17
M-CSF (pg/ml)	70.7 ± 2.8	54.1 ± 3.6*	41.1 ± 3.4* <sup>#</sup>	49.8 ± 2.6* <sup>#</sup>	57.6 ± 2.4*	60.6 ± 2.2*

**Discussion**

Mechanical loading has been well recognized as one of the key regulators for the mechanical and biological structure and function of bone. Bone-related cells respond to a diverse range of mechanical stimulations such as tensile stress from stretching [19,21-26], shear stress from fluid flow [27], four-point bending [28], and hydrostatic pressure [29]. In recent years, accumulating evidence has demonstrated that LMHFV promotes bone formation *in vivo* [9,17]. However, the effects of LMHFV on bone cells *in vitro* have not been comprehensively investigated.

In this study, we examined the effects of sLMHFV and iLMHFV on osteoblast proliferation, osteogenic protein expression, and osteoclastogenic cytokine production, using the MC3T3-E1 preosteoblasts as an *in vitro* model system. Our MTT assay results indicated that sLMHFV or iLMHFV treatment for up to 7 d did not cause significant changes in MC3T3-E1 cell proliferation. Past studies on the effects of LMHFV on preosteoblast proliferation have generated controversial results. LMHFV has been reported to inhibit the proliferation of human periodontal ligament stem cells and Bone Marrow-Derived Mesenchymal Stromal Cells (BMSC) [16,30]. However, LMHFV has also been reported to promote BMSC proliferation [31]. In this study, we found that MC3T3-E1 cell proliferation was not affected by sLMHFV or iLMHFV under the experimental conditions tested. Interestingly, studies have shown that the effects of pressure loading on osteoblast proliferation are frequency and duration-dependent [32]. Taking all previous studies and our data into consideration, we can believe that the effects of LMHFV on preosteoblast proliferation depend highly on the cell type and vibration parameters.

Many studies have shown that mechanical stimulation promotes osteogenic differentiation and expression of ALP, COL I and OCN, proteins required for bone matrix synthesis, both *in vitro* and *in vivo* [33-35]. Specifically, LMHFV has been shown to strengthen bones and promote fracture healing *in vivo* [10,12,14,15]. In addition, sLMHFV has been reported to promote osteogenic differentiation *in vitro* [16]. In this study, we found that both sLMHFV and iLMHFV significantly increased the expression of ALP, COL I, and OCN in MC3T3-E1 preosteoblasts, indicating that both sLMHFV and iLMHFV promote osteogenic differentiation *in vitro*. Intriguingly, we found that iLMHFV was more effective than sLMHFV in

inducing osteoblast differentiation, suggesting that iLMHFV may have greater osteogenic effects. We speculate that osteoblasts could adapt to the stimulation, thus fail to respond as rigorous as it does at the beginning, and could not translate the mechanical stimulation to biological responses in the continuous manner. When osteoblasts are subject to an intermittent rest between stimulation, they can regain the ability and sensitivity to vibration stimulation, thus creating a biomechanical response far better than continuous stimulation.

Previous *in vitro* studies have shown that sLMHFV can inhibit osteoclastic differentiation and osteoclastic resorption by suppressing the production of osteocyte-derived osteoclastogenic cytokines [14,36]. In this study, we found that both sLMHFV and iLMHFV significantly decreased the level of M-CSF and RANKL secreted by MC3T3-E1 cells. sLMHFV and iLMHFV also significantly increased the OPG/RANKL ratio, a critical regulator of osteoclast formation, activation, and survival. Thus, our results suggested that both sLMHFV and iLMHFV have anti-resorption potential in bone remodeling and homeostasis. Intriguingly, iLMHFV has more profound effect than sLMHFV in inhibiting the production of osteoclastogenic cytokines, suggesting that iLMHFV may contribute more in protecting bone tissue from osteoclastic absorption.

Taken together, our results suggested that, compared with sLMHFV, iLMHFV has more profound effect in enhancing bone formation and healing. These *in vitro* results are on par with previous *in vivo* studies, which showed that iLMHFV loading was superior to sLMHFV loading in upregulation osteogenic proteins and promoting healing in osteoporotic fractures [17,37]. Sustained mechanical loading may cause adaptive tolerance to the stimuli, resulting in gradually diminishing effects [38]. We believe that periodical rest between LMHFV may help maintain or even restore the sensitivity to the stimulation of cells, and thus generating greater effect towards desired function. Moreover, we found that 0.5 h of iLMHFV can provide optimal results, with longer treatment being less effective. These results were in agreement with previous findings that early exposure to LMHFV can positively modulate osteoporotic bone and osteoblasts; however, the beneficial effects seem to not give a consistent performance over time [39]. We believe that this is because prolonged intermittent vibration could result in similar effect as

sostenuto vibration, which may cause fatigue of cells to mechanical stimuli, leading to desensitization over time.

Our findings could provide useful insights in the development of clinically applicable LMHFV treatment regimen to strengthen bones and promote fracture healing. Clinical trials are required to investigate the therapeutic potential of LMHFV to treat osteoporosis in human in order to harness our findings to benefit more patients.

## Conclusion

In conclusion, we have shown that both sLMHFV and iLMHFV can increase the expression level of the osteogenic proteins ALP, OCN, and COL I in MC3T3-E1 preosteoblasts. sLMHFV and iLMHFV also can inhibit the secretion of osteoclastogenic cytokines M-CSF and RANKL, as well as can increase the OPN/RANKL ratio. These osteogenic and anti-osteoclastogenic effects were most profound with short iLMHFV.

## Acknowledgments

This study was supported by the Key Program and General Programs of the National Natural Science Foundation of China (No. 11432016, 11272134 and 11602093).

## Conflict of Interest Statement

The authors declare no conflict of interest.

## References

- Ganveer GB, Tiwari RR. Injury pattern among non-fatal road traffic accident cases: a cross-sectional study in Central India. *Indian J Med Sci* 2005; 59: 9-12.
- Blotman F, Cortet B, Hilliquin P, Avouac B, Allaert FA, Pouchain D, Gaudin AF, Cotté FE, El Hasnaoui A. Characterisation of patients with postmenopausal osteoporosis in French primary healthcare. *Drugs Aging* 2007; 24: 603-614.
- Nelson FR, Brighton CT, Ryaby J, Simon BJ, Nielson JH, Lorich DG, Bolander M, Seelig J. Use of physical forces in bone healing. *J Am Acad Orthop Surg* 2003; 11: 344-354.
- Ehrlich PJ, Lanyon LE. Mechanical strain and bone cell function: a review. *Osteoporos Int* 2002; 13: 688-700.
- Rubin J, Murphy TC, Rahnert J, Song H, Nanes MS, Greenfield EM, Jo H, Fan X. Mechanical inhibition of RANKL expression is regulated by H-Ras-GTPase. *J Biol Chem* 2006; 281: 1412-1418.
- Chen JH, Liu C, You L, Simmons CA. Boning up on Wolff's Law: mechanical regulation of the cells that make and maintain bone. *J Biomech* 2010; 43: 108-118.
- Fan X, Rahnert JA, Murphy TC, Nanes MS, Greenfield EM, Rubin J. Response to mechanical strain in an immortalized pre-osteoblast cell is dependent on ERK1/2. *J Cell Physiol* 2006; 207: 454-460.
- Ogawa T, Zhang X, Naert I, Vermaelen P, Deroose CM, Sasaki K, Duyck J. The effect of whole-body vibration on peri-implant bone healing in rats. *Clin Oral Implants Res* 2011; 22: 302-307.
- Zhang R, Gong H, Zhu D, Gao J, Fang J, Fan Y. Seven day insertion rest in whole body vibration improves multi-level bone quality in tail suspension rats. *PLoS One* 2014; 9: 92312.
- Rubin C, Turner AS, Bain S, Mallinckrodt C, McLeod K. Anabolism. Low mechanical signals strengthen long bones. *Nature* 2001; 412: 603-604.
- Prisby RD, Lafage-Proust MH, Malaval L, Belli A, Vico L. Effects of whole body vibration on the skeleton and other organ systems in man and animal models: what we know and what we need to know. *Ageing Res Rev* 2008; 7: 319-329.
- Ozcivici E, Luu YK, Rubin CT, Judex S. Low-level vibrations retain bone marrow's osteogenic potential and augment recovery of trabecular bone during reambulation. *PLoS One* 2010; 5: 11178.
- Wysocki A, Butler M, Shamliyan T, Kane RL. Whole-body vibration therapy for osteoporosis: state of the science. *Ann Intern Med* 2011; 155: 680-686, W206-213.
- Zhou Y, Guan X, Liu T, Wang X, Yu M, Yang G, Wang H. Whole body vibration improves osseointegration by up-regulating osteoblastic activity but down-regulating osteoblast-mediated osteoclastogenesis via ERK1/2 pathway. *Bone* 2015; 71: 17-24.
- Leung KS, Shi HF, Cheung WH, Qin L, Ng WK, Tam KF, Tang N. Low-magnitude high-frequency vibration accelerates callus formation, mineralization, and fracture healing in rats. *J Orthop Res* 2009; 27: 458-465.
- Zhou Y, Guan X, Zhu Z, Gao S, Zhang C, Li C, Zhou K, Hou W, Yu H. Osteogenic differentiation of bone marrow-derived mesenchymal stromal cells on bone-derived scaffolds: effect of microvibration and role of ERK1/2 activation. *Eur Cell Mater* 2011; 22: 12-25.
- Ma R, Zhu D, Gong H, Gu G, Huang X, Gao J, Zhang X. High-frequency and low-magnitude whole body vibration with rest days is more effective in improving skeletal micro-morphology and biomechanical properties in ovariectomised rodents. *Hip Int* 2012; 22: 218-226.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001; 25: 402-408.
- Ziros PG, Gil AP, Georgakopoulos T, Habeos I, Kletsas D, Basdra EK, Papavassiliou AG. The bone-specific transcriptional regulator Cbfa1 is a target of mechanical signals in osteoblastic cells. *J Biol Chem* 2002; 277: 23934-23941.
- Ducy P, Zhang R, Geoffroy V, Ridall AL, Karsenty G. *Osf2/Cbfa1*: a transcriptional activator of osteoblast differentiation. *Cell* 1997; 89: 747-754.
- Brighton CT, Strafford B, Gross SB, Leatherwood DF, Williams JL, Pollack SR. The proliferative and synthetic response of isolated calvarial bone cells of rats to cyclic

*Intermittent low-magnitude high-frequency vibration promotes osteogenic protein expression and inhibits osteoclastogenic cytokine secretion in osteoblasts*

- biaxial mechanical strain. *J Bone Joint Surg Am* 1991; 73: 320-331.
22. Zhuang H, Wang W, Tahernia AD, Levitz CL, Luchetti WT, Brighton CT. Mechanical strain-induced proliferation of osteoblastic cells parallels increased TGF-beta 1 mRNA. *Biochem Biophys Res Commun* 1996; 229: 449-453.
23. Matsuda N, Morita N, Matsuda K, Watanabe M. Proliferation and differentiation of human osteoblastic cells associated with differential activation of MAP kinases in response to epidermal growth factor, hypoxia, and mechanical stress in vitro. *Biochem Biophys Res Commun* 1998; 249: 350-354.
24. Cheng M, Zaman G, Rawlinson SC, Mohan S, Baylink DJ, Lanyon LE. Mechanical strain stimulates ROS cell proliferation through IGF-II and estrogen through IGF-I. *J Bone Miner Res* 1999; 14: 1742-1750.
25. Nagatomi J, Arulanandam BP, Metzger DW, Meunier A, Bizios R. Frequency- and duration-dependent effects of cyclic pressure on select bone cell functions. *Tissue Eng* 2001; 7: 717-728.
26. Weyts FA, Bosmans B, Niesing R, van Leeuwen JP, Weinans H. Mechanical control of human osteoblast apoptosis and proliferation in relation to differentiation. *Calcif Tissue Int* 2003; 72: 505-512.
27. Mikuni-Takagaki Y, Suzuki Y, Kawase T, Saito S. Distinct responses of different populations of bone cells to mechanical stress. *Endocrinology* 1996; 137: 2028-2035.
28. You J, Yellowley CE, Donahue HJ, Zhang Y, Chen Q, Jacobs CR. Substrate deformation levels associated with routine physical activity are less stimulatory to bone cells relative to loading-induced oscillatory fluid flow. *J Biomech Eng* 2000; 122: 387-393.
29. Kaspar D, Seidl W, Neidlinger-Wilke C, Ignatius A, Claes L. Dynamic cell stretching increases human osteoblast proliferation and CICP synthesis but decreases osteocalcin synthesis and alkaline phosphatase activity. *J Biomech* 2000; 33: 45-51.
30. Zhang C, Li J, Zhang L, Zhou Y, Hou W, Quan H, Li X, Chen Y, Yu H. Effects of mechanical vibration on proliferation and osteogenic differentiation of human periodontal ligament stem cells. *Arch Oral Biol* 2012; 57: 1395-1407.
31. Luu YK, Capilla E, Rosen CJ, Gilsanz V, Pessin JE, Judex S, Rubin CT. Mechanical stimulation of mesenchymal stem cell proliferation and differentiation promotes osteogenesis while preventing dietary-induced obesity. *J Bone Miner Res* 2009; 24: 50-61.
32. Garcia-Giralt N, Nogues X, Enjuanes A, Puig J, Mellibovsky L, Bay-Jensen A, Carreras R, Balcells S, Díez-Pérez A, Grinberg D. Two new single-nucleotide polymorphisms in the COL1A1 upstream regulatory region and their relationship to bone mineral density. *J Bone Miner Res* 2002; 17: 384-393.
33. Tanno M, Furukawa KI, Ueyama K, Harata S, Motomura S. Uniaxial cyclic stretch induces osteogenic differentiation and synthesis of bone morphogenetic proteins of spinal ligament cells derived from patients with ossification of the posterior longitudinal ligaments. *Bone* 2003; 33: 475-484.
34. Guo Y, Zhang CQ, Zeng QC, Li RX, Liu L, Hao QX, Shi CH, Zhang XZ, Yan YX. Mechanical strain promotes osteoblast ECM formation and improves its osteoinductive potential. *Biomed Eng* 2012; 11: 80.
35. Lu HF, Mai ZH, Xu Y, Wang W, Ai H. Mechanical loading induced expression of bone morphogenetic protein-2, alkaline phosphatase activity, and collagen synthesis in osteoblastic MC3T3-E1 cells. *Chin Med J (Engl)* 2012; 125: 4093-4097.
36. Lau E, Al-Dujaili S, Guenther A, Liu D, Wang L, You L. Effect of low-magnitude, high-frequency vibration on osteocytes in the regulation of osteoclasts. *Bone* 2010; 46: 1508-1515.
37. Li M, Wu W, Tan L, Mu D, Zhu D, Wang J, Zhao B. Low-magnitude mechanical vibration regulates expression of osteogenic proteins in ovariectomized rats. *Biochem Biophys Res Commun* 2015; 465: 344-348.
38. Davies PF. Flow-mediated endothelial mechanotransduction. *Physiol Rev* 1995; 75: 519-560.
39. Qing F, Xie P, Liem YS, Chen Y, Chen X, Zhu X, Fan Y, Yang X, Zhang X. Administration duration influences the effects of low-magnitude, high-frequency vibration on ovariectomized rat bone. *J Orthop Res* 2016; 34: 1147-1157.

**\*Correspondence to**

Dong Zhu  
Department of Orthopedic Traumatology Trauma  
First Hospital of Jilin University  
PR China