

Pathogenicity determinants and epidemiology of uropathogenic *E. coli* (UPEC) strains isolated from children with urinary tract infection (UTI) to define distinct pathotypes.

Shahin Najar Peerayeh¹, Masoumeh Navidinia^{2*}, Fatemeh Fallah³, Bita Bakhshi¹, Jamshid Jamali⁴

¹Bacteriology Department, Tarbiat Modares University, Tehran, IR Iran

²Medical Laboratory Sciences Department, School of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

³Pediatric Infectious Research Center, School of Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

⁴Department of Biostatistics and Epidemiology, School of Health, Mashhad University of Medical Sciences, Mashhad, IR Iran

Abstract

Background/purpose: One of the most common childhood diseases is Urinary Tract Infection (UTI). Without diagnosis and treatment, it can cause irreparable effects. *Escherichia coli* cause UTI in 75% of cases. Unlike diarrheagenic *E. coli* has certain pathotypes, *E. coli* causing UTI, are not well known. For further information, we considered pathogenicity determinants and epidemiology of uropathogenic *E. coli* (UPEC) strains isolated from children with Urinary Tract Infection (UTI) to define distinct pathotypes.

Methods: One hundred *E. coli* strains (50 UPEC and 50 commensal) isolated from children with UTI were examined. Some virulence factors and specific genes were examined by PCR method. Genetic diversity was evaluated by phylogenetic typing groups.

Results: Some pathogenicity determinants were more prevalent in UPEC strains rather than commensal *E. coli* strains, significantly. There were PAI IICFT073, PAI II J96, PAI I536, PAI ICFT073, PAIII536, PAI IV, *gafD*, *focG*, *vat*, *usp*, *hlyD*, *sat*, *cnf1*, *picU*, *fliC(H7)*, *kpsMTII*, *kpsMTIII*. UPEC were mainly found in phylogenetic typing groups B2 and D, while in commensal isolates, phylogenetic groups A and D were the most common.

Conclusion: We need a simple pathotypes screening test which can be used either as or could be beneficial to facilitate along with other experiments in establishing an UTI assessment. Unfortunately, due to the high variation in pathogenicity determinants of UPEC strains, pathotypes could not be determined using genotype and virulence factors. Knowledge of the molecular details of UPEC is mainstay of successful strategies development for treatment of UTI and prevention of its subsequent complications.

Keywords: Pathogenicity determinants, Uropathogenic *E. coli* (UPEC), Pathotypes, Children, Urinary tract infection (UTI).

Accepted on March 14, 2018

Introduction

Urinary Tract Infection (UTI) is the most prevalent infectious diseases, and very problematic worldwide [1]. Uropathogenic *E. coli* (UPEC) which can colonize successfully in urinary tract is the primary etiologic agents associated with UTI. UPEC isolates express a wide range of virulence factors and specific genes. In fact, UPEC can exceed some types of host cells that comprise the stratified layers of the bladder urothelium, such as differentiated superficial facet cells, less mature intermediate,

and basal epithelial cells. Invasion of host cell is facilitated both the establishment and permanence of UPEC within the urinary tract. Pathogenic extraintestinal *E. coli* isolates or ExPEC (such as UPEC) and commensal *E. coli* typically differ in phylogenetic group and virulent characteristics. According to previous studies, pathogenic ExPEC isolates belong to phylogenetic groups B2, and D. While, commensal *E. coli* strains belong to groups A and B1. In addition, pathogenic ExPEC isolates carry specialized pathogenic factors, i.e., traits that confer pathogenic potential, which are uncommon between

commensal isolates, UPEC strains and other *E. coli* strains are involved in various extraintestinal infections. These strains are distinguished from commensal strains by particular pathogenic factors such as adherence characters, toxins genes, and iron uptake systems. Bacterial adherence depending on the assembly in fimbrial projecting or afimbrial aggregates. The toxins related to ExPEC strains mostly display cytotoxic necrotizing factor and hemolysin, contribute to destruction of eukaryotic cells.

Siderophores give to the strains the advantage of obtaining iron from the ambience to exist and reproduce. Co-expression of virulence factors contributes to the host defense system defeat and onset of infection. Pathogenicity-Associated Islands (PAI) are codifying genes localized in distinctive area on the bacterial chromosome. In addition, ExPEC strains carry some virulence factors that are rarely between commensal *E. coli* isolates. Some of these virulence factors which is encoded by PAIs, preparing a mechanism for coordinated horizontal factors, such as properties that confer the ability to transfer pathogenic virulence genes. Hacker et al. first defined PAI character as mobile genetic elements in the late 1980s. These contain short direct repeats of fragments of DNA more than >10 kb nearby to the tRNA genes and comprise insertion sequences, integrases, and have a high percent G+C content that varies from the host bacterial. Particular sets of virulence factors are associated with UTI caused by *E. coli*. A variety of virulence genes are contributed by bacterial strains operating them by an individual pathogenesis process, are named a "pathotypes." Consideration of co-occurrence of potential UTI virulence factors between different *E. coli* isolates from commensal and urine collections provides documents for defining multiple pathotypes of UPEC, but recent understanding of critical genetic discrepancy to define the pathotypes, is limited. Finding of *E. coli* extra genes involved in uropathogenesis and consideration of their distribution will further describe UPEC pathotypes and permit to a more analysis of details of how these pathotypes might vary in how they cause infection [2-6]. For further information, we considered pathogenicity determinants and epidemiology of Uropathogenic *E. coli* (UPEC) strains isolated from children with Urinary tract Infection (UTI) to define distinct pathotypes.

Methods

UTI definition

Totally, 100 *E. coli* strains isolated from children presenting symptomatic UTI of both sex and different ages (2-12 y old) were hospitalized in the nephrology ward or were visited outpatient in Mofid Children Hospital, Tehran, Iran. Isolated *E. coli* colonies were recognized by standard bacteriological procedures. Commensal *E. coli* strains, including the well-characterized strains MG1655 and HS were used as controls. All distinct colonies were recognized morphologically and were stored for molecular examination, as described previously by Plos and Foxman. Classification of phylogenetic grouping of the *E. coli* strains was performed by PCR-based method

using a combination of three DNA markers (*chuA*, *yjaA*, *TspE4.C2*) [7,8].

Detection of *E. coli* virulence determinants

All isolates were tested by PCR method for the existence of 31 bacterial genes related to UTI that as following as: phylogenetic typing groups genes; *chuA*, *yjaA*, *TspE4.C2*, adhesions groups; *afa*, *bmaE*, *fimH*, *gafD*, *focG*; protectin related genes; *kpsMT (K1)*, *kpsMTII*, *kpsMTIII*, *rfa* (*O4 LPS*), common toxins related to UPEC; *vat*, *usp*, *cvaC*, *hlyD*, *cdtB*, *sat*, *cnf1*, *picU* and, different PAI; PAI *IICFT073*, PAI *II536*, PAI *III536*, PAI *I536*, PAI *IV536*, PAI *ICFT073*, PAI *I J96*, PAI *III J96*, Miscellaneous genes; *fliC (H7)* *ibeA*, *ompT*. These pathogenic factors were part of a large set of virulence genes described previously by Johnson et al. Boiled whole-cell lysates 400 ml were used as DNA template and amplification was performed in a 25 µl reaction mixture containing 4 mM MgCl₂, 2.5 µl reaction buffer 10X, 2.5 U iTaq TM DNA polymerase (5 U/µl), 0.5 mM each primer and 5 µl DNA template. Reactions were performed in a Gene PCR System (Eppendorff). A reaction mixture without a DNA template was used as a negative control. Isolates producing a clear zone of hemolysis around colonies on sheep-blood agar, considered to be positive for hemolysin production [2,9].

Statistics

Continuous variables were expressed as mean ± Standard Deviation (SD). Discrete variables were reported as frequency and percentage. Chi square test and Fisher's exact test were used to access the relation between quantities' variables. For all statistical analyses, a p-value of <0.05 was considered to be significant. Statistical analysis was conducted using the SPSS version 21.

Results

Characteristics of patients

Total of 100 *E. coli* isolates were analysed. Of these, 50 were collected from midstream clean catch urine and 50 isolated from stool of the same patients who were in the nephrology ward in Mofid Children Hospital in Tehran.

Phylogenetic typing group distribution

The distribution of commensal *E. coli* isolates and UPEC isolates among the four phylogenetic groups is as following: Of the 50 commensal *E. coli* isolates, 44% fell into group A, 16% into B2, 32% into D, and UPEC isolates fell into group D, 54% into B2, 8% into A.

Pathogenicity island genes distribution

Distribution PAI genes, such as PAI *ICFT073* (74 vs. 26%), PAI *IICFT073* (38 vs. 14%), PAI *I536* (36 vs. 6%), PAI *IV536* (86 vs. 42%), PAI *II J96* (30 vs. 10%) were more frequent virulence markers in UPEC isolates than commensal *E. coli* and PAI

Pathogenicity determinants and epidemiology of uropathogenic E. coli (UPEC) strains isolated from children with urinary tract infection (UTI) to define distinct pathotypes

II536 (22 vs. 4%), *PAI III536* (6 vs. 0%), *PAI I J96* (4 vs. 0%) markers was almost similar in UPEC isolates and commensal *E. coli* (Table 1).

Adhesion genes distribution

Distribution of adhesion genes, such as *bmaE* (16 vs. 6%), *gafD* (20 vs. 2%), *focG* (22 vs. 6%) were more frequent virulence markers in UPEC than commensal *E. coli* and *afa* (6 vs. 10%), *fimH* (92 vs. 98%) markers were almost similar in both isolates (Table 2).

Toxin related genes distribution

Distribution of toxin genes, such as *vat* (96 vs. 4%), *usp* (54 vs. 6%), *hlyD* (26 vs. 2%), *cdtB* (18 vs. 10%), *sat* (44 vs. 8%), *cnfI* (26 vs. 0%), *picU* (42 vs. 2%) were more frequent virulence markers in UPEC isolates than commensal *E. coli*. *cvaC* (20 vs. 66%) was most frequent marker in commensal *E. coli* (Table 3).

Miscellaneous genes distribution

Distribution of miscellaneous genes, such as *fliC* (H7(26 vs. 10%)), *ompT* (62 vs. 58%) were more frequent virulence factors in UPEC than commensal *E. coli*, and *ibeA* (14 vs. 26%) was almost similar in both of them (Table 4).

Protectins genes distribution

Distribution of protectins genes, such as *kpsMTII* (70 vs. 46%) was more frequent virulence gene in UPEC rather than commensal *E. coli*. The prevalence of *kpsMTI* (K1) (46 vs. 54%), *kpsMTIII* (14 vs. 2%), *rfc* (O4 LPS) (6 vs. 2%), were almost similar in both of them (Table 5).

Table 1. PAI distribution between commensal *E. coli* and UPEC in children with UTI in Iran.

PAI genes	Commensal <i>E. coli</i>	UPEC	p-value
<i>PAI III 536</i>	0 (0%)	3 (6%)	0.324
<i>PAI IV 536</i>	21 (42%)	43 (86%)	<0.001
<i>PAI II CFTO73</i>	7 (14%)	19 (38%)	<0.001
<i>PAI I 536</i>	3 (6%)	18 (36%)	<0.001
<i>PAI II 536</i>	2 (4%)	11 (22%)	<0.001
<i>PAI I J96</i>	0 (0%)	2 (4%)	0.329
<i>PAI II J96</i>	5 (10%)	15 (30%)	0.001
<i>PAI I CFTO73</i>	13 (26%)	37 (74%)	<0.001

Table 2. Adhesion genes distribution between commensal *E. coli* and UPEC in children with UTI in Iran.

Adhesion genes	Commensal <i>E. coli</i>	UPEC	p-value
<i>afa</i>	5 (10%)	3 (6%)	0.298

<i>bma E</i>	3 (6%)	8 (16%)	0.007
<i>fim H</i>	49 (98%)	46 (92%)	0.534
<i>gaf D</i>	1 (2%)	10 (20%)	<0.001
<i>foc G</i>	3 (6%)	11 (22.0)	0.006

Table 3. Toxin related genes distribution between commensal *E. coli* and UPEC in children with UTI in Iran.

Toxin genes	Commensal <i>E. coli</i>	UPEC	p-value
<i>cdtB</i>	5 (10%)	9 (18%)	0.194
<i>hlyD</i>	1 (2%)	13 (26%)	0.001
<i>cnf1</i>	0 (0%)	13 (26%)	<0.001
<i>cva C</i>	33 (66%)	10 (20%)	<0.001
<i>usp</i>	3 (6%)	27 (54%)	<0.001
<i>vat</i>	2 (4%)	48 (96%)	<0.001
<i>sat</i>	4 (8%)	22 (44%)	<0.001
<i>picU</i>	1 (2%)	21 (42%)	<0.001

Table 4. Miscellaneous genes distribution between UPEC and commensal *E. coli* in children with UTI in Iran.

Miscellaneous genes	Commensal <i>E. coli</i>	UPEC	p-value
<i>fliC H</i>	5 (10%)	13 (26%)	0.020
<i>ibeA</i>	13 (26%)	7 (14%)	0.325
<i>ompT</i>	29 (58%)	31 (62%)	0.895

Table 5. Protectins genes distribution between commensal *E. coli* and UPEC in children with UTI in Iran.

Protectins genes	Commensal <i>E. coli</i>	UPEC	p-value
<i>kps MTI</i>	27 (54%)	23 (46%)	0.707
<i>kps MT II</i>	23 (46%)	35 (70%)	<0.001
<i>kps MT III</i>	1 (2%)	7 (14%)	0.083
<i>rfc</i>	1 (2%)	3 (6%)	0.700

Discussion

It is thought to the pathogenic *E. coli* strains are related to the presence of virulence factors. According to products called virulence factors, *E. coli* bacteria adhere selectively to the uroepithelial mucosa, promote colonization and persist in the urinary tract, induce a local inflammatory response, and sometimes promote tissue destruction. The result of a complex combination of special attributes of the *E. coli* causes to movement of a bacterium from the intestinal tract to the kidney and bladder. Our goal in this study was to pathogenicity determinants and epidemiology of Uropathogenic *E. coli* (UPEC) strains isolated from children with UTI to define distinct pathotypes. The phylogenetic groups' distribution

varies considerably among UPEC and commensal *E. coli* isolates [10].

Group A were the most prevalent phylogenetic group (41%) between the commensal *E. coli*, and that group B2 isolates were the least whereas in UPEC isolates, group B2 being the most prevalent (54%) and group B1 being least prevalent (4%). Our results were similar with those reported by Duriez et al. who detected that group B2 was seldom *E. coli* isolates (11%), whereas groups A and B1 were the most common (40% and 34%, respectively) [11].

Also, according to studies of Duriez et al. and Picard et al. group B2 has been reported with different prevalence in commensal *E. coli* isolates between 2-19% [11-14].

Some studies detected groups A and D in enteroaggregative *E. coli*. It is proven that other pathotypes such as enteropathogenic *E. coli* (EPEC) and enterotoxigenic *E. coli* (ETEC) belong to phylogenetic groups A and B2 [15-19].

As the same like as Johnson study, we also identified that the phylogenetic group B2 were dominant in urine isolates. One of the main reasons due to being more virulence factors in UPEC compared to other groups [20].

A significant difference on phylogenetic groups B2 and D distribution between UPEC and commensal *E. coli* due to some reasons. Being additional alternative route for causing disease and infection and being specificity of bacterial pathogenesis are the most reason on these results. ExPEC especially UPEC mainly belong to the phylogenetic group B2 and D groups while commensal *E. coli* mostly belong to group A and B1 [21-27].

As we mentioned above, the most dominant phylogenetic groups in UPEC was B2 or D. Also, we identified phylogenetic group D rather than A in commensal *E. coli*. Previous studies have demonstrated that human health are potentially affected by UPEC strains, especially group B2 with virulence genes profiles, which in the face with a high proportion of commensal isolates existence in intestinal, these genes could be transferred them [28,29].

In contrast in Duriez study, it was recognized that group B was the most prevalent between both commensal and UPEC isolates, and recommended that the detected differences with respect to previous results could be impressed by the population and age selected for study. In the present research, children were 2-12 y old; therefore, the discrepancy between the two studies cannot be illustrated by the age selected. Duriez et al. studied females aged between 18-39 y old, which is an age range with a high occurrence of the proportion of UPEC asymptomatic carriers. It was interesting that the mean number of PAIs in isolates related to groups B2 and A was alike in both UPEC and commensal *E. coli*. In comparison, isolates related to group D significantly had PAIs rather than intestinal flora, but was non-significant between isolates related to group B1. In based on Duriez et al.'s study, they compared virulence factors and phylogenetic groups between commensal strains and recognized that commensal isolates related to groups A,

B1 and D. They exposed fewer virulence factors rather than ExPEC strains [11].

Actually, that commensal *E. coli* related to group B2 are less virulent than strains isolated from clinical specimens, as observed in this study and previous studies, reinforce the hypothesis that it is mainly the most virulent isolates related to these groups in intestinal tract, can produce UTI and other infections. Some of virulence factor genes locate on genetic elements called on Pathogenicity Islands (PAI) in the vicinity of tRNA genes. In UPEC these islands are detected and have proven their role [30,31].

Two multiplex PCR for detection of 8 pathogenesis-related PAI in this study were used. The prevalence of PAI in UPEC was% 89 compared to commensal *E. coli* 38%. There are evidences that the intestinal environment have *E. coli* strains, mainly belong to B2 group which contain large numbers of PAI. *PAI IV536* island pathogenesis has been seen the most common PAI in both groups. This PAI is detected a lot in the Enterobacteriaceae family [18,32-36].

We were observed *PAI IV536* (also called HPI) 19% in commensal *E. coli* isolates compared to 43% in UPEC isolates. Being high prevalence of *PAI IV536* in isolates is led to the hypothesis that HPI is a structural island to be assumed as a pathogenesis island [37,38].

However, *in-vivo* experiments showed that HPI had important role in ExPEC strains. Similar with Middendorf et al. study, the high frequency of *PAI IV536* could be explained by PAI stability in *E. coli* [39].

Based on Bingen-Bidois study in 2002 on urosepsis producing *E. coli*, the frequency of PAIs was reported as follows: *PAI IV536* (92%), *PAI IJ96* (24%), *PAI ICFT073* (19%), *PAI I536* (1%), but neither *PAI Ij96* nor *PAI II536* was observed, and *PAI IICFT073* studied *PAI III536* were not studied. Results were showed the high frequency of *PAI IV536* but low frequency of *PAI I536* (19% vs. 73%), and higher frequency of *PAI IJ96* (24%) compare to *PAI ICFT073* (19%) [40]. They suggested that the role of *PAI II536* and *PAI IJ96* in urosepsis pathogenesis may not be important, and no difference was identified in distribution rate of *PAI II536* among *E. coli* isolates from patients with urosepsis or pyelonephritis. Dobrindt et al. reported that 64.5% of UPEC isolates and 39.3% of non-pathogenic *E. coli* had *PAI III536* and concluded that *PAI II536* was more common than *PAI I536*, *PAI III536* [41]. Middendorf et al. reported that *PAI II536* and *PAI III536* were very unstable and hence were easily lost and could be explained the difference between our findings and the findings of Dobrindt [39,41].

Similar with the other studies, PAIs numbers in UPEC and commensal isolates showed that the average number of PAI in isolates belonging B2 and A groups were the same. PAIs numbers in D group were significantly higher in commensal *E. coli*, but it in the case of B1 group was not significant. Similar PAI combination was seen in many isolates that had the same number of PAIs regardless of phylogenetic groups and their origin. Although, PAI acquisition is not a random phenomenon

Pathogenicity determinants and epidemiology of uropathogenic E. coli (UPEC) strains isolated from children with urinary tract infection (UTI) to define distinct pathotypes

and is carried out in a planned mechanism. It is proven that PAI IV536 initially gains on chromosome and establishes to stable condition.

PAI IV536 is often seen alone in strains except group B2 but sometimes there are exceptions, and it also is detected along with PAI ICFT073. PAI IICFT073 may be the third island, which gained followed by PAI IJ96 and PAI I536. The PAI II536 and III536 in the final stage earn. This fact explains why they are so unusual and can be seen in highly virulent strains. PAI IJ96 and PAI IICFT073 compete on goal to replace on tRNA^{phe}. The results indicate that PAI IICFT073 have more affiliation compare to PAI IJ96 for the target. However, when commensal populations of *E. coli* and ExPEC consider based on their phylogenetic groups reveals more differences. Because *E. coli* isolated from feces mainly belong to B1 and A groups while the population of ExPEC often belong to B2 group and has a lot number of virulence factors. Therefore, it is unclear whether all *E. coli* isolated from the intestinal tract of healthy people in a specific time, regardless of phylogenetic groups become commensal or just isolates belonging to groups A and B1 are commensal.

Similarly, we don't know in the case of *E. coli* belongs to A and B1 groups isolated from patients with UTI should be considered as natural pathogens or commensal *E. coli* in a host with healthy immune system. These findings suggest that intestinal flora may act as a reservoir and can stimulate UTI by means of commensal *E. coli* with the B2 phylogenetic group.

The prevalence of *afa* gene in UPEC and commensal *E. coli* was 6%, 10% respectively. Gene prevalence in different studies was 2-11% due to the low prevalence of this gene in the *E. coli* strains [42-46]. *fimH* is common in *E. coli* strains. In fact, most clinical isolates both virulent and non-virulent can produce type 1 fimbriae. In epidemiological studies, there were no evidence of the relationship between type 1 fimbriae and severity of the infection. Several studies in experimental models indicated that type 1 fimbriae could have an effective role in the stability of *E. coli* in the urinary tract. Renovation of *fimH*-1 adhesion greatly reduces UPEC ability to colonize the urinary tract in human volunteers and mice. In this study, *fimH* prevalence has been reported 92% in UPEC, and 98% in commensal *E. coli* [47-49].

It is noteworthy that *fimH* adhesion in UPEC plays important role to connect and invade host cells and produce intracellular bacterial apartments (biofilm formation) [48]. According to genotyping study, *fimH* almost more than 90% of UPEC strains and pathogenic *E. coli* in birds, was reported [50].

In Johnson et al.'s study, *gafD* detected more than 20% of *E. coli* isolates with other virulence factors such as *sfaS*, *focG*, *afa/dra*, *bmaE*, *gafD*, *cnf1*, *cdtB*, *cvaC*, *ibeA* and most was frequently in relationship with phylogenetic group B2 which was agreement with our study. In this study, *focG* prevalence among UPEC, commensal *E. coli* was 22%, 6%, respectively, which was agreement with Johnsons findings [29].

In this study, 20% of strains belonged to phylogenetic group B2 gene had *hlyD* gene, while only in 4% of group A and 2%

of B1 and 10% of group D, *hlyD* gene was detected. Forty two percent of UPEC isolates had hemolytic activity and 26% of gene carried *hlyD*. The reason due to be the low percentage of *hlyD* gene carriers strains (10%) belonged to group B2 [24].

We identified *cvaC* 20% and 66% among UPEC and commensal *E. coli*, respectively. Low prevalence of *cvaC* in UPEC strains suggests that this gene may be placed on non-ColV plasmid or PAI such as pTJ100. In Johnson study, based on bird pathogenic *E. coli* examination, they reported *hlyA* 41%, *cnf1* 16%, *cdtB* 8%, but about 28% of UPEC had *cnf1*. We reported *hlyD* (26%, 2%), *cnf1* (26%, 0%), *cdtB* (18%, 10%) in UPEC and commensal *E. coli*, respectively. This disagreement is caused by differences in study population [25,41,51].

By means of PCR method, *fimH*, *kpsM*, *hlyD*, *usp*, *cnf1*, *afa* were detected in UPEC strains, 92%, 46%, 26%, 54%, 26%, 6%, respectively. In this study, the prevalence of *fimH* in UPEC strains was high. These results demonstrate that type 1 fimbriae are important virulence factor. The type 1 fimbriae have been shown to enhance inflammation and played an important role in the pathogenesis of ascending UTIs.

Connell et al. reported that infected mice by strains 01: K1: H7 with present type 1 fimbriae had more stimulation of neutrophil cells rather than infected mice by type 1 negative strains. Our results showed that *fimH* was associated to P and S fimbriae in UPEC strains [52-56].

In a study conducted in Japan, *usp* gene was detected in 80% of 195 strains isolated from cystitis, and 93% of the 76 strains isolated from pyelonephritis [57]. According to different studies, *usp* has often been observed associated with pyelonephritis rather than cystitis. In a study conducted by Kanamaru, *usp* was identified in 22.2% of isolates. The difference between results due to differences in studied population (women) or between clones of bacterial isolated from women in Brazil and Japan [58,59].

usp gene is homologous with zonula occludens in *Vibrio cholera*. In a study conducted in Japan, *usp* was diagnosed in 54% of *E. coli* strains isolated from healthy subjects stool samples, 80% isolated from cystitis and 93% of strains pertaining to pyelonephritis [57].

Most of UPEC strains have capsule group 2 (K1, K5) are coded by the operon *kps*. Capsules in UPEC associated with pyelonephritis is very common [60]. In this study, UPEC strains contained approximately 70% *kpsMTII*. All isolates was observed in relation to other virulence factors in UPEC including *kpsMTII* and *hlyA* [29,58].

In first time *cdt* producing *E. coli* were observed in relation to children with enteritis [61]. In other studies, *cdt* gene was observed in strains isolated from urosepsis *E. coli* and fecal *E. coli* and in patients with various symptoms such as diarrhea, encephalopathy [29,62].

cdt gene cause irreversible inhibition of cell cycle at the G₂/M and produce single nuclear giant cells. The results suggest that cystitis can cause *cdtB* negative strains but this *E. coli* is less

virulent compare to strains cause pyelonephritis, urosepsis and diarrhea.

Specifically, we reported 96% vat in UPEC isolates compared to 4% in commensal *E. coli* strains. In Tiba et al.'s study, 39% of the isolates in bird's pathogen (APEC) revealed vat that these mainly related in the phylogenetic group D. In commensal strains and UPEC, *pic* was more common in phylogenetic groups B2. *sat* (44%) was more common in related to UPEC isolates and also, *picU* (42%) was observed in UPEC more than commensal *E. coli*, too [57]. Agreement with Timothy and Germon studies, *ibeA* observed in *E. coli* pathogens related to UTI and along with other numerous virulence factors more naturally associated to this infection [63,64].

In another study, *ibeA* was reported in about 26% of pathogenic *E. coli* strains isolated from chickens, and was not observed in non-pathogenic strains. Also, *ibeA* in APEC, *E. coli* isolated from vagina, and infant meningitis was observed (26%), (32%), (33-40%), respectively. So, we concluded that this factor significantly associated with pathogenic strains [17,64,65].

Compared with the study of Rodriguez-Siek, we reported 26% *fliC* (*H7*) in UPEC but they reported 4.8% in APEC. The difference between results due to differences in studied population. Similar to Zhao et al. *ompT* was reported 68% in our study [50].

Based on Rodriguez-Siek study in 2005, most of UPEC causing UTI in human presented capsules *kpsMT* (*K1*), *kpsMTIII* genes. These genes identified rarely in APEC. So it has been concluded that these genes closely related to ExPEC strains such as UPEC. Specifically, capsular antigen K1 often has been observed in UPEC [65-67].

However, significant relationship between these factors and producing UTI cause this hypothesis that extraintestinal movement of bacteria after acquisition of some virulence factor and ability to ascending urinary tract is very important to cause disease. It is proven that these virulence factors present in strains cause meningitis rather than other types of *E. coli* [63].

Similar to Bert, *kpsMTIII* have been observed significantly more compare with the other capsular genes. Even, *kpsMT III* and *rfa* (*O4LPS*) have been reported less than 10% of the cases. In Zhao et al.'s reports, *kpsMT* (*K1*) was involved in the synthesis of capsules in 45% APEC and UPEC [50,68].

The present study provides molecular and epidemiological information about virulence factor genes found in two groups of *E. coli*. It is necessary to have a simple pathotypes screening test which can be beneficial to facilitate along with other experiments in establishing an UTI assessment. Unfortunately, due to the high variation in pathogenicity determinants of UPEC strains, pathotypes could not be determined using pathogenicity determinants. Knowledge of the molecular details of UPEC is mainstay of successful strategies development for treatment of UTI and prevention of its subsequent complications.

Funding Information

The design of the study, collection, analysis, interpretation of data, and in writing the manuscript was supported by a grant from Tarbiat Modares University, Faculty of Medical Sciences, and Tehran, Iran. There is no conflict of interests.

Acknowledgements

We thank Prof. A. Karimi and Dr. S. Maham, Dr. M. Mohkam and N. Heidari for their support.

References

1. Navidinia M, Teymouri AR, Goudarzi M. Assessment of correlation between urinary secretory IgA (sIgA) levels and different types of urinary tract infection (UTI) in various age groups. *J Paramed Sci (JPS)* 2018; 9: 51-55.
2. Navidinia M, Peerayeh SN, Fallah F, Bakhshi B, Sajadinia RS. Phylogenetic grouping and pathotypic comparison of urine and commensal *Escherichia coli* isolates from children with urinary tract infection. *Braz J Microbiol* 2014; 45: 509-514.
3. Navidinia M, Najari Peerayeh S, Fallah F, Bakhshi B, Adabian S, Alimehr S. Distribution of the Pathogenicity Islands markers (PAIs) in uropathogenic *E. coli* isolated from children in Mofid children hospital. *Arch Ped Infect Dis* 2013; 1: 75-79.
4. Marrs CF, Zhang L, Foxman B. *Escherichia coli* mediated urinary tract infections: are there distinct uropathogenic *E. coli* (UPEC) pathotypes? *FEMS Microbiol Lett* 2005; 252: 183-190.
5. Karimi A, Fallah F, Navidinia M, Malekan MA. Detection of integron elements and gene groups encoding ESBLs and their prevalence in *Escherichia coli* and *Klebsiella* isolated from urine samples by PCR method. *Afr J Microbiol Res* 2012; 6: 1806-1809.
6. Fallah F, Karimi A, Goudarzi M, Shiva F, Navidinia M, Jahromi MH. Determination of integron frequency by a polymerase chain reaction restriction fragment length polymorphism method in multidrug-resistant *Escherichia coli*, which causes urinary tract infections. *Microb Drug Resist* 2012; 18: 546-549.
7. Plos K, Connell H, Jodal U, Marklund BI, Mårild S, Wettergren B. Intestinal carriage of P fimbriated *Escherichia coli* and the susceptibility to urinary tract infection in young children. *J Infect Dis* 1995; 171: 625-631.
8. Foxman B, Manning SD, Tallman P, Bauer R, Zhang L, Koopman JS. Uropathogenic *Escherichia coli* are more likely than commensal *E. coli* to be shared between heterosexual sex partners. *Am J Epidemiol* 2002; 156: 1133-1140.
9. Ghenghesh KS, Elkateb E, Barbash N, Abdel Nada R, Ahmed SF, Rahouma A. Uropathogens from diabetic patients in Libya: Virulence factors and phylogenetic groups of *Escherichia coli* isolates. *J Med Microbiol* 2009; 58: 1006-1014.

Pathogenicity determinants and epidemiology of uropathogenic E. coli (UPEC) strains isolated from children with urinary tract infection (UTI) to define distinct pathotypes

10. Giray B, UÇAR FB, Aydemir SS. Characterization of uropathogenic *E. coli* in relation to virulence factors. *Turk J Med Sci* 2012; 42: 1328-1337.
11. Duriez P, Clermont O, Bonacorsi S, Bingen E, Chaventre A, Elion J. Commensal *Escherichia coli* isolates are phylogenetically distributed among geographically distinct human populations. *Microbiol* 2001; 147: 1671-1676.
12. Picard B, Garcia JS, Gouriou S, Duriez P, Brahim N, Bingen E. The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect Immun* 1999; 67: 546-553.
13. Picard B, Picard-Pasquier N, Krishnamoorthy R, Goulet P. Characterization of highly virulent *Escherichia coli* strains by ribosomal DNA restriction fragment length polymorphism. *FEMS Microbiol Lett* 1991; 82: 183-188.
14. Picard B, Journet-Mancy C, Picard-Pasquier N, Goulet P. Genetic structures of the B2 and B1 *Escherichia coli* strains responsible for extra-intestinal infections. *J Gen Microbiol* 1993; 139: 3079-3088.
15. Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. *Am J Med* 2002; 113: 5-13.
16. Moreno E, Prats G, Sabate M, Perez T, Johnson JR, Andreu A. Quinolone, fluoroquinolone and trimethoprim sulfamethoxazole resistance in relation to virulence determinants and phylogenetic background among uropathogenic *Escherichia coli*. *J Antimicrob Chemother* 2006; 57: 204-211.
17. Johnson JR, Delavari P, Kuskowski M, Stell AL. Phylogenetic distribution of extraintestinal virulence-associated traits in *Escherichia coli*. *J Infect Dis* 2001; 183: 78-88.
18. Zhang L, Foxman B, Manning SD, Tallman P. Molecular epidemiologic approaches to urinary tract infection gene discovery in uropathogenic *Escherichia coli*. *Infect Immun* 2008; 38: 1150-1158.
19. Zhang L, Foxman B, Marrs C. Both urinary and rectal *Escherichia coli* isolates are dominated by strains of phylogenetic group B2. *J Clin Microbiol* 2002; 40: 3951-3955.
20. Johnson JR, Obryan TT, Sandberg T. Phylogenetic and pathotypic comparison of concurrent urine and rectal *Escherichia coli* isolates from men with febrile urinary tract infection. *J Clin Microbiol* 2005; 43: 3895-3900.
21. Clermont O, Bonacorsi S, Bingen E. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* 2000; 66: 4555-4558.
22. Abraham S, Chapman TA, Zhang R, Chin J, Mabbett AN, Totsika M. Molecular characterization of *Escherichia coli* strains that cause symptomatic and asymptomatic urinary tract infections. *J Clin Microbiol* 2012; 50: 1027-1030.
23. Patti G, Mannini A, Balistreri M, Schito AM. Virulence factors in urinary *Escherichia coli* strains: phylogenetic background and quinolone and fluoroquinolone resistance. *J Clin Microbiol* 2008; 46: 480-487.
24. Johnson JR, Russo TA. Uropathogenic *Escherichia coli* as agents of diverse non-urinary tract extraintestinal infections. *J Infect Dis* 2002; 186: 859-864.
25. Rodriguez-Siek KE, Giddings CW, Doetkott C, Johnson TJ, Fakhr MK, Nolan LK. Comparison of *Escherichia coli* isolates implicated in human urinary tract infection and avian colibacillosis. *Microbiol* 2005; 151: 2097-2110.
26. Moreno E, Prats G, Sabate M, Perez T, Johnson JR, Andreu A. Quinolone, fluoroquinolone and trimethoprim sulfamethoxazole resistance in relation to virulence determinants and phylogenetic background among uropathogenic *Escherichia coli*. *J Antimicrob Chemother* 2006; 57: 204-211.
27. Tenaillon O, Skurnik D, Picard B, Denamur E. The population genetics of commensal *Escherichia coli*. *Nat Rev Microbiol* 2010; 8: 207-217.
28. Burman WJ, Breese PE, Murray BE, Singh KV, Batal HA, MacKenzie TD, Ogle JW, Wilson ML, Reves RR, Mehler PS. Conventional and molecular epidemiology of trimethoprim-sulfamethoxazole resistance among urinary *Escherichia coli* isolates. *Am J Med* 2003; 115: 358-364.
29. Johnson JR, Stell AL, Delavari P, Murray AC, Kuskowski M, Gaastra W. Phylogenetic and pathotypic similarities between *Escherichia coli* isolates from urinary tract infections in dogs and extraintestinal infections in humans. *J Infect Dis* 2001; 183: 897-906.
30. Blum G, Ott M, Lischewski A, Ritter A, Imrich H, Tschape H, Hacker J. Excision of large DNA regions termed pathogenicity islands from tRNA-specific loci in the chromosome of an *Escherichia coli* wild-type pathogen. *Infect Immun* 1994; 62: 606-614.
31. Silva GJD, Mendonca N. Association between antimicrobial resistance and virulence in *Escherichia coli*. *Virulence* 2012; 3: 18-28.
32. Sandip P, Linardopoulou EV, Billig M, Tchesnokova V, Price LB, Johnson JR, Chattopadhyay S, Sokurenko EV. Role of homologous recombination in adaptive diversification of extra-intestinal *Escherichia coli*. *J bacteriol* 2012; 8: 1-48.
33. Kanamaru S, Kurazano H, Nakano M, Terai A, Ogawa O, Yamamoto S. Subtyping of uropathogenic *Escherichia coli* according to the pathogenicity island encoding uropathogenic-specific protein: comparison with phylogenetic groups. *Int J Urol* 2006; 13: 754-760.
34. Schubert S, Rakin A, Karch H, Carniel E, Heesemann J. Prevalence of the high-pathogenicity island of *Yersinia* species among *Escherichia coli* strains that are pathogenic to humans. *Infect Immun* 1998; 66: 480-485.
35. Schubert S, Rakin A, Heesemann J. The *Yersinia* high pathogenicity island (HPI): evolutionary and functional aspects. *Int J Med Microbiol* 2004; 294: 83-94.
36. Schubert S, Picard B, Gouriou S, Heesemann J, Denamur E. *Yersinia* high-pathogenicity island contributes to virulence in *Escherichia coli* causing extraintestinal infections. *Infect Immun* 2002; 70: 5335-5337.

37. Oelschlaeger TA, Dobrindt U, Hacker J. Virulence factors of uropathogens. *Curr Opin Urol* 2003; 21: 12-33.
38. Karch H, Schubert S, Zhang D, Zhang W, Schmidt H, Olschlager T, Hacker J. A genomic island, termed high-pathogenicity island, is present in certain non-O157 Shiga toxin-producing *Escherichia coli* clonal lineages. *Infect Immun* 1999; 67: 5994-6001.
39. Middendorf B, Hochhut B, Leipold K, Dobrindt U, Blum-Oehler G, Hacker J. Instability of pathogenicity islands in uropathogenic *Escherichia coli* 536. *J Bacteriol* 2004; 186: 3086-3096.
40. Bingen-Bidois M, Clermont O, Bonacorsi S, Terki M, Brahimi N, Loukil C, Barraud D, Bingen E. Phylogenetic analysis and prevalence of urosepsis strains of *Escherichia coli* bearing pathogenicity island-like domains. *Infect Immun* 2002; 70: 3216-3226.
41. Dobrindt U, Blum-Oehler G, Nagy G, Schneider G, Johann A, Gottschalk G, Hacker J. Genetic structure and distribution of four pathogenicity islands (PAI I (536) to PAI IV (536) of uropathogenic *Escherichia coli* strain 536. *Infect Immun* 2002; 70: 6365-6372.
42. Blanco M, Blanco J, Blanco JE, Alonso MP, Abalia I, Rodriguez E, Bilbao JR, Umaran A. Virulence factors and o serogroups of *Escherichia coli* as a cause of community acquired urinary infections. *Enferm Infect Microbiol Clin* 1995; 13: 236-241.
43. Santo E, Macedo C, Marin JM. Virulence factors of uropathogenic *Escherichia coli* from a university hospital in Ribeirao Preto, Sao Paulo Brazil. *Rev Inst Med Trop Sao Paulo* 2006; 48: 185-188.
44. Abe CM, Salvador FA, Falsetti IN, Vieira MA, Blanco J, Blanco JE, Blanco M, Machado AM, Elias WP, Hernandez RT, Gomes TA. Uropathogenic *Escherichia coli* (UPEC) strains may carry virulence properties of diarrhoeagenic *E. coli*. *FEMS Immunol Med Microbiol* 2008; 52: 397-406.
45. Blanco M, Blanco JE, Alonso MP, Mora A, Balsalobre C, Munoa F, Juarez A, Blanco J. Detection of pap, sfa and afa adhesin-encoding operons in uropathogenic *Escherichia coli* strains: relationship with expression of adhesins and production of toxins. *Res Microbiol* 1997; 148: 745-755.
46. Takahashi A, Kanamaru S, Kurazono H, Kunishima Y, Tsukamoto T, Ogawa O, Yamamoto S. *Escherichia coli* isolates associated with uncomplicated and complicated cystitis and asymptomatic bacteriuria possess similar phylogenies, virulence genes, and O-serogroup profiles. *J Clin Microbiol* 2006; 44: 4589-4592.
47. Mougeot FKB, Buckles ELC, Lockatell V, Hebel JR, Johnson DE, Tang CM, Donnenberg MS. Type 1 fimbriae and extracellular polysaccharides are preeminent uropathogenic *Escherichia coli* virulence determinants in the murine urinary tract. *Mol Microbiol* 2002; 45: 1079-1093.
48. Wiles TJ, Kulesus RR, Mulvey MA. Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Exp Mol Path* 2008; 85: 11-19.
49. Brumbaugh AR, Mobley HLT. Preventing urinary tract infection: progress toward an effective *Escherichia coli* vaccine. *Expert Rev Vaccines* 2012; 11: 663-676.
50. Zhao L, Gao S, Huan H, Xu X, Zhu X, Yang W, Gao Q, Liu X. Comparison of virulence factors and expression of specific genes between uropathogenic *Escherichia coli* and avian pathogenic *E. coli* in a murine urinary tract infection model and a chicken challenge model. *Microbiol* 2009; 155: 1634-1644.
51. Sorsa LJ, Dufke S, Heesemann J, Schubert S. Characterization of an iroBCDEN gene cluster on a transmissible plasmid of uropathogenic *Escherichia coli*: evidence for horizontal transfer of a chromosomal virulence factor. *Infect Immun* 2003; 71: 3285-3293. Mulvey MA. Adhesion and entry of uropathogenic *Escherichia coli*. *Cell Microbiol* 2002; 4: 257-267.
52. Schlager TA, Whittam TS, Hendley JO, Bhang JL, Wobbe CL, Stapleton A. Variation in frequency of the virulence-factor in *Escherichia coli* clones colonizing the stools and urinary tracts of healthy prepubertal girls. *JID* 2003; 8: 188-211.
53. Johnson JR, Kuskowski MA, Gajewski A. Extended virulence genotypes and phylogenetic background of *Escherichia coli* isolates from patients with cystitis, pyelonephritis, or prostatitis. *J Infect Dis* 2005; 191: 46-50.
54. Morin MD, Hopkins WJ. Identification of virulence genes in uropathogenic *Escherichia coli* by multiplex polymerase chain reaction and their association with infectivity in mice. *Urol* 2002; 60: 537-541.
55. Snyder JA, Haugen BJ, Virginia Lockatell C, Maroncle N, Hagan EC, Johnson DE, Welch RA, Mobley HLT. Coordinate expression of fimbriae in uropathogenic *Escherichia coli*. *Infect Immun* 2005; 73: 7588-7596.
56. Kurazono H, Yamamoto S, Nakano M. Characterization of a putative virulence island in the chromosome of uropathogenic *Escherichia coli* possessing a gene encoding an uropathogenic specific protein. *Microb Pathog* 2000; 28: 183-189.
57. Tiba MR, Yano T, Leite DS. Genotypic characterization of virulence factors in *Escherichia coli* strains from patients with cystitis. *Rev Inst Med trop S Paulo* 2008; 50: 255-260.
58. Kanamaru S, Kurazono H, Ishitoya S, Tera A, Habuchi T, Nakano M, Ogawa O, Yamamoto S. Distribution and genetic association of putative uropathogenic virulence factors *iroN*, *iha*, *kpsMT*, *ompT* and *usp* in *Escherichia coli* isolated from urinary tract infections in Japan. *J Urol* 2003; 170: 2490-2493.
59. Krachler AM, Orth K. Targeting the bacteria-host interface. Strategies in anti-adhesion therapy. *Virulence* 2013; 4: 1-11.
60. Kadhum HJ, Finlay D, Rowe MT, Wilson IG, Ball HJ. Occurrence and characteristics of cytotoxic necrotizing factors, cytolethal distending toxins and other virulence factors in *Escherichia coli* from human blood and fecal samples. *Epidem Infect* 2008; 136: 752-760.

Pathogenicity determinants and epidemiology of uropathogenic E. coli (UPEC) strains isolated from children with urinary tract infection (UTI) to define distinct pathotypes

61. Bielaszewska M, Stoewe F, Fruth A, Zhang W, Prager R, Brockmeyer J, Mellmann A, Karch H, Friedrich AW. Shiga toxin, cytolethal distending toxin, and hemolysin repertoires in clinical Escherichia coli O91 isolates. *J Clin Microbiol* 2009; 47: 2061-2066.
62. Timothy S, Shafi K, Leatherbarrow AH, Jordan FTW, Wigley P. Molecular epidemiology of a reproductive tract-associated colibacillosis outbreak in a layer breeder flock associated with atypical avian pathogenic Escherichia coli. *Avian Path* 2008; 37: 375-378.
63. Germon P, Chen YH, He L, Blanco JE, Bree A, Schouler C, Huang SH, Moulin-Schouleur M. *ibeA*, a virulence factor of avian pathogenic Escherichia coli. *Microbiol* 2005; 151: 1179-1186.
64. Obata-Yasuoka M, Ba-Thein W, Tsukamoto T, Yoshikawa H, Hayashi H. Vaginal Escherichia coli share common virulence factor profiles, serotypes and phylogeny with other extraintestinal E. coli. *Microbiol* 2002; 148: 2745-2752.
65. Solberg OD, Ajiboye RM, Riley LW. Origin of class 1 and 2 integrons and gene cassettes in a population-based sample of uropathogenic Escherichia coli. *J Clin Microbiol* 2006; 44: 1347-1351.
66. Ewers C, Li G, Wilking H, Kiebetaling S, Alt K, Antao EM, Laternus C, Diehl I, Glodde S, Homeier T, Bohnke U, Steinruck H, Philipp HC, Wieler LH. Avian pathogenic, uropathogenic, and newborn meningitis-causing Escherichia coli: how closely related are they? *Int J Med Microbiol* 2007; 297: 163-176.
67. Johnson TJ, Kariyawasam S, Wannemuehler Y, Mangiamela P, Johnson SJ, Doetkott C, Skyberg JA, Lynne AM, Johnson JR, Nolan LK. The genome sequence of avian pathogenic Escherichia coli strain O1: K1: H7 shares strong similarities with human extraintestinal pathogenic E. coli genomes. *J Bacteriol* 2007; 189: 3228-3236.
68. Bert F, Johnson JR, Ouattara B, Leflon-Guibout V, Johnston B, Marcon E, Valla D, Moreau R, Chanoine MHN. Genetic diversity and virulence profiles of Escherichia coli isolates causing spontaneous bacterial peritonitis and bacteremia in patients with cirrhosis. *J Clin Microbiol* 2010; 48: 2709-2714.

***Correspondence to**

Masoumeh Navidinia
Medical Laboratory Sciences Department
School of Allied Medical Sciences
Shahid Beheshti University of Medical Sciences
IR Iran