



Prospective Bacteriological study of Surgical Site infections in a tertiary care hospital at Saharsa

Ram Nagina Sinha

Assistant Professor, Department of General Surgery, Lord Buddha Koshi Medical College & Hospital, Saharsa, Bihar, India.

Received:

30th March 2013

Received in revised form: 11th April 2013

Accepted:

15th April 2013 Available

online:

20th April 2013



Online ISSN 2249-22X
<http://www.jbiopharm.com>

ABSTRACT

Background: common post-operative complications. Apart from bacterial contamination of wound, various patient and environment related factors play role in development and outcome of SSI. The present study is undertaken to study the frequency of SSI with reference to factors contributing to it and the antimicrobial susceptibility pattern of the causative organisms.

Methods: This single-observer, cross-sectional, complete-enumeration prospective study was carried out over a period of one year (from March 2012 to Feb 2013) in the department of Surgery at Department of General Surgery, Lord Buddha Koshi Medical College & Hospital, Saharsa, Bihar. 196 pus samples from cases of surgical site infections were processed for gram staining, culture, biochemical identification tests and antimicrobial susceptibility testing. Methicillin-Resistant Staphylococcus aureus (MRSA) strains were detected by using oxacillin and cefoxitin disk diffusion and minimum inhibitory concentration (MIC) of oxacillin was tested by broth dilution technique.

Results: The overall frequency of SSI was 6.17%. Most common isolates were Staphylococcus aureus, coagulase negative Staphylococci (CONS), E. coli and Pseudomonas aeruginosa. The frequency of MRSA was 8.6%. The maximum frequency was among patients operated on emergency basis in surgical department.

Conclusions: The most important determinants for SSI were emergency surgery and presence of co-morbid conditions. The frequency of occurrence was age-dependent, with maximum rate of SSI in males and females in the third and sixth decades of life, respectively.

Keywords: Surgical site infections, Post-operative wound infections, Clean and clean-contaminated wound, MRSA

INTRODUCTION

After urinary tract infection, surgical site infections (SSI) are the most common nosocomial infections in hospitalized patients.¹ As many as 1% of the patients undergoing clean and 11% of patients undergoing clean contaminated surgery experience SSI.² There is substantial burden with increased morbidity and exceeding healthcare costs.³ Despite efforts to

control infection and better understanding of sepsis, wound infection is still a clinical problem and some infections in clean wounds still remain unexplained.⁴ In many SSIs, the responsible pathogens originate from patient's endogenous flora. The causative pathogen depends on the type of surgery; the most commonly isolated organisms are Staphylococcus aureus, coagulase negative Staphylococci,

Enterococcus spp and E. coli.⁵ The present study was undertaken to determine the frequency of surgical site infections with reference to various directly or indirectly related pre-disposing factors. The study also aimed at ascertaining the surgical site infection rate, type of bacteria commonly involved, their antibiotic sensitivity pattern; to determine the frequency of methicillin-resistant Staphylococcus aureus (MRSA) and to calculate the oxacillin minimum inhibitory concentration for MRSA.

METHODS

This single-observer, cross-sectional, complete - enumeration, prospective study was conducted over a one-year period in the in the department of Surgery at Department of General Surgery, Lord Buddha Koshi Medical College & Hospital, Saharsa, Bihar. After obtaining approval from Institutional Ethics Committee, all patients of either sex, admitted in departments of Surgery, who had undergone surgical procedures during the study period, were included. The surgical procedures were classified as planned (elective) surgeries, emergency surgeries and clean, clean-contaminated surgeries and patients were divided accordingly. The surgical sites were inspected at frequent intervals on days 3/5, 7 and 10, and further whenever required, for clinical evidence of infection. Samples from patients with clinical evidence of infection (purulent discharge) were processed for bacteriological examination. Details of patients' age, sex, clinical diagnosis, date of admission, personal history, and date of surgical procedure, type of procedure (clean/ clean-contaminated) and prophylactic antibiotics were recorded in a pre-tested proforma. Sample collection All samples were collected early in the morning before dressing of the wounds. After cleaning the surrounding skin with 70% alcohol, the pus was collected from deep inside the wound in sterile bulbs using syringes or using sterile swab sticks in duplicate.⁶ Sample processing The first swab was used direct smear and gram staining, the second swab was inoculated on blood agar & MacConkey's agar. The plates were incubated overnight at 37° C and the morphology of colonies was observed the following morning. Identification of organisms was done by secondary smears examination and

biochemical tests.⁶ Antimicrobial susceptibility testing Broth cultures of isolated organisms were prepared and matched with McFarland 0.5 turbidity standard. Then these were tested for various antibiotics on Muller Hinton Agar plates by Kirby-Bauer Disk Diffusion Technique. The plates were incubated at 37° C for 18-24 hours.⁶ The antimicrobials were selected for susceptibility testing according to CLSI guidelines. The MIC of oxacillin was tested for the strains resistant to oxacillin and/or cefoxitin. Detection of methicillin resistant staphylococcus aureus (MRSA) was done by cefoxitin disk diffusion testing; Oxacillin disk diffusion testing and minimum inhibitory concentration of oxacillin by Broth dilution method using Muller-Hinton broth. MIC $\geq 4\mu\text{g/ml}$ indicated methicillin resistance.^{7,8}

RESULTS

A total of 3,160 patients admitted in the tertiary care hospital had undergone surgical procedures during the study period. Of these surgeries, 1,328 (904 elective, 424 emergency) were conducted by the Surgery Department. Out of 1,328 surgeries, 700 (57.3%) were "clean", 280 (21.8%) were "clean contaminated" and 272 (20.82%) were "contaminated". The infection rates in clean, clean-contaminated and contaminated wounds were 2.86%, 7.84% and 13.52%, respectively. The occurrence of SSI in elective and emergency surgeries was 4.32% (n=76) and 8.48% (n=119), respectively. The frequency of SSI was 8.35% (n=117, 68 males and 43 females) in the Department of Surgery, 4.68% (n=48 females). Though the rate of SSI was higher in females (male-female ratio 1: 1.11), the gender difference was not statistically significant. The overall rate of SSI (both sexes combined) was in the age group of 21-30 years. However, the highest rates among males and females were seen in age group of 21-30 years and 51-60 years, respectively. Out of 3,160 cases, 195 (6.17%) patients had SSI. Of the 195 samples collected in duplicate, cultures of 11 (5.64%) samples did not show any growth of organisms.

Out of 184 wound infections, 24 were caused by two micro-organisms. In remaining 160 samples, only one organism was isolated. The frequency of mixed infections was higher in patients who had undergone

gastrointestinal surgeries. The commensal flora of the gastro-intestinal tract could be the associated

factor. On cultivation, 208 strains were obtained (Table 1).

Table 1: Isolated organisms from patients.

Organism	Isolates (n = 208)
<i>Staphylococcus aureus</i>	46 (22.1%)
<i>Coagulase negative staphylococci (CoNS)</i>	34 (16.3%)
<i>E.coli</i>	29 (13.9%)
<i>Pseudomonas aeruginosa</i>	22 (10.5%)
<i>Klebsiella pneumonia</i>	16 (7.6%)
<i>Enterococcus spp</i>	13 (6.2%)
<i>Citrobacter freundii</i>	12 (5.7%)
<i>Enterobacter spp</i>	9 (4.3%)
<i>Proteus mirabilis</i>	8 (3.8%)
<i>Proteus vulgaris</i>	6 (2.8%)
<i>Acinetobacter spp</i>	6 (2.8%)
<i>Citrobacter diversus</i>	4 (1.9%)
<i>Streptococcus spp</i>	3 (1.4%)

Table 2: Antibiotic susceptibility of gram positive cocci.

Antibiotic	<i>Staphylococcus aureus</i>	<i>CoNS</i>	<i>Enterococcus spp.</i>	<i>Streptococcus spp.</i>
Penicillin	6 (13%)	9 (26.4%)	6 (46.1%)	2 (66.6%)
Amoxycillin	10 (21.7%)	14 (41.1%)	7 (53.8%)	2 (66.6%)
Oxacillin	28 (60.8%)	24 (70.5%)	10 (76.9%)	3 (100%)
Vancomycin	46 (100%)	34 (100%)	13 (100%)	3 (100%)
Erythromycin	20 (43.4%)	20 (61.2%)	-	3 (100%)
Cotrimoxazole	21 (45.6%)	19 (55.8%)	-	-
Amikacin	26 (56.5%)	26 (76.4%)	-	3 (100%)
Ciprofloxacin	17 (36.9%)	14 (41.1%)	6 (46.1%)	3 (100%)

Table 3: Antibiotic susceptibility of gram negative bacilli (*Enterobacteriaceae*).

Antibiotic	<i>E.coli</i>	<i>Klebsiella spp.</i>	<i>C.freundii</i>	<i>Enterobacter</i>	<i>P.mirabilis</i>	<i>P. vulgaris</i>	<i>C.diversus</i>
Amoxycillin	4 (13.8%)	2(12.5%)	3 (25%)	3 (33.3%)	2 (25%)	1 (16.6%)	1 (25%)
Cephalexin	12(41.3%)	5 (31.2%)	2 (16.3%)	6 (66.6%)	6 (75%)	3 (50%)	0 (0%)
Gentamicin	18(62%)	10 (50%)	8 (66.6%)	5 (55.5%)	5 (62.5%)	4 (66.6%)	3 (75%)
Amikacin	21(72.4%)	12 (75%)	9 (75%)	7 (77.7%)	6 (75%)	4 (66.6%)	4 (100%)
Cefoperazone	16(55.1%)	11(68.7%)	5 (41.7%)	7 (77.7%)	6 (75%)	4 (66.6%)	2 (50%)
Cefepime	21(72.4%)	11(68.7%)	8 (66.6%)	8 (88.8%)	7 (83.5%)	4 (66.6%)	4 (100%)
Ciprofloxacin	12(41.3%)	6 (37.5%)	5 (41.7%)	4 (44.4%)	4 (50%)	3 (50%)	1 (25%)
Cotrimoxazole	15(51.7%)	6 (37.5%)	7 (58.3%)	5 (55.5%)	5 (62.5%)	3 (50%)	2 (50%)

DISCUSSION

SSI The microbiology of SSI largely depends upon the kind of operation performed, the types of work load and the hospital environment.^{9,10} Comparative statistics from hospitals in different parts of the world may be unreliable and even misleading. The prevalence of pathogen varies from place to place and hospital to hospital.¹¹ Periodic studies carried out in the same hospital over a number of years would provide more reliable information.¹⁰ Frequency of infection Surgical site infections remain

a major issue of patient safety despite of improvements in surgical practice and infection control techniques.¹² The overall infection rate in current study is 6.17% whereas that reported by other studies ranges from 4-33%.^{4, 13-18} Department-wise distribution of patients The higher frequency of SSI observed in Department of Surgery could be because of higher number of emergency procedures conducted in the department. Similarly, substantial number of surgeries had dealt with gastrointestinal and urinary systems, which contributed to clean

contaminated wound. Age and sex-wise distribution of patients. The reason for higher rate of SSI in third decade of life in this study could be the greater number of young adults getting operated for exploratory laparotomy for antral or appendicular perforation or appendicitis. Similar observations have been made by other researchers.^{4,18} Organisms isolated Frequency of culture negative SSI in the present study was 5.4%, which compares with 7.28% reported in a tertiary care hospital in Bangalore.¹⁴ The cause of failure to grow any bacterium on culture could either be due to anaerobic nature of the organism or non-viability of organisms in patients on antibiotic therapy. Staphylococcus species The frequency of isolation of Staphylococci is depicted in Table 1. The frequency of isolates of Staphylococcus aureus from infected surgical wounds reported by other researchers was in the range of 19-75%.^{11,14,15,19,20} Enterococcus species Among catalase negative organisms, 6.2% were Enterococci in the present study while that reported by other authors ranged from 0.06-7.1%.^{13,14,19} β haemolytic Streptococci Streptococcus spp was isolated from 3 cases (Table 1). A study from Nagpur has reported its frequency as 3.78%.¹¹ The decreasing frequency of Streptococcus spp as a causative agent of SSI may be ascribed to its susceptibility to even age-old antimicrobial drugs. The emerging multidrug resistant organisms are now taking the place of Streptococci. Enterobacteriaceae In the current study, E.coli was isolated from 29 (13.9%) patients (Table 1). Our finding correlates with the frequencies (7.5-21%) reported by other studies.^{4,11,14,15,19} There were 14 isolates of Klebsiella pneumoniae and 2 of Klebsiella oxytoca (Table 1). This correlates with the findings reported by other studies.^{4,11,14,15} The present study showed 7.6% of isolates were that of Citrobacter spp (12 C. freundii and 4 C. diversus). Other studies from Nagpur and New Delhi have reported frequencies of 2.27% and 4.1%, respectively.^{11,13} Of the isolates of Proteus spp, P. mirabilis and P. vulgaris comprised 8 (3.8%) and 6 (2.8%), respectively (Table 1). Other studies have reported the frequency of Proteus spp. isolates in range of 2.9-6.8%.^{11,14,15,17} The present study showed 9 (4.3%) were Enterobacter spp, whereas a study

from New Delhi has reported the frequency of isolation of the same organism as 2.7%.¹³ Pseudomonas aeruginosa. Twenty-two (10.5%) strains of Pseudomonas aeruginosa were isolated in the current study (Table 1). Other studies have reported the frequency of isolates of this organism in the range of 10.7-19%.^{4,11,13,14,17} Acinetobacter species Out of 208 organisms isolated in the present study, 6 (2.8%) were Acinetobacter spp., (Table 1). Other studies have reported higher frequency of Acinetobacter sp ranging from 10.53-29%.^{4,13,17} Antibiotic sensitivity tests Staphylococcus spp (Table 2). A Mumbai-based study has reported 100% penicillin resistance among S. aureus, whereas ampicillin resistance reported by a Nagpur-based study and a Bangalore-based study was 91.8% and 77.7%, respectively.^{11,14,16} These compare with the findings of our study. MRSA: (Table 2) A seven-year review of MRSA susceptibility has reported a constant susceptibility rate of 98% to co-trimoxazole with significant decrease in susceptibility of quinolones (63%) and also reported that susceptibility to erythromycin has decreased from 15% in the year 2000, to 4% in the year 2006.²¹ The current study obtained comparable susceptibility in relation to erythromycin, but a lower susceptibility to cotrimoxazole. All the MRSA in current study were sensitive to vancomycin, as found in many other studies.^{13,14,16,21} Coagulase negative staphylococci (CoNS) (Table 2) Using disc diffusion method, the frequency of oxacillin resistant CoNS was 29.5% in the current study while other studies have reported frequencies ranging from 14- 40%.²²⁻²⁵ The frequency of co-existing resistance to oxacillin and ciprofloxacin, which was proposed to be an alternative therapy for MRSA infection, has ranged from 23% to 100%.^{22,26} Enterobacteriaceae E. coli with higher resistance against fluoroquinolones and 100% susceptibility to fluoroquinolones and 66.6% susceptibility to Amikacin have been reported.^{13,14} Amikacin remains the antimicrobial with highest susceptibility according to our study. The high degree of resistance reported in Klebsiella spp by many researchers was also observed in the current study (Table 3).^{13,14,17}

CONCLUSION

To conclude, the overall frequency of SSI was 6.17% (2.86% in clean, 7.84% in clean contaminated and 13.52% in contaminated wounds). Maximum frequency of SSI was in the third decade of life for males and sixth decade for females. The most common gram positive and gram negative isolates were *Staphylococcus aureus* (22.1%) and *E. coli* (13.9%), respectively. The frequency of MRSA was 8.6%. All *Staphylococci* and *Enterococci* were vancomycin sensitive. Among Gram-negative bacilli, the highest percentage susceptibility was to amikacin. Imipenem was most effective antipseudomonal drug with 100% susceptibility.

REFERENCES

1. Kirby JP, Mazuski JE. Prevention of surgical site infection. Interventional studies for preventing surgical site infections in sub-Saharan Africa: A systematic review. *Int J Surg*. 2012;10(5):242-9.
2. National Nosocomial Infections Surveillance System. National Nosocomial Infection (NNIS) system report, data summary from January 1992 through June 2004, issues October 2004. *Am J Infect Control*. 2004;32:470-85.
3. Langelotz C, Mueller-Rau C, Terziyski S, Rau B, Krannich A, Gastmeier P, et al. Gender-specific differences in Surgical Site Infections: An analysis of 438, 050 surgical procedures from the German National Nosocomial Infections Surveillance System. *Viszeralmedizin*. 2014;30(2):114-7.
4. Mundhada AS, Tenpe S. A study of organisms causing surgical site infections and their antimicrobial susceptibility in a tertiary care Government Hospital. *Indian J Pathol Microbiol*. 2015;58(2):195-200.
5. Owens CD, Stoessel K. Surgical Site Infections: epidemiology, microbiology and prevention. *J Hosp Infect*. 2008;70 Suppl-2:3-10.
6. Collee JG, Miles RS, Watt B. Tests for identification of bacteria. In: Collee JG, Fraser AG, Marmion BP, Simmons A. (Eds.) *Mackie & McCartney's Practical Medical Microbiology*. 14th Edn. London: Churchill Livingstone. 2006:131-49.
7. Skov R, Smyth R, Clausen M, Larsen AR, Frimodt-Moller N, Olsson-Liljequist B, et al. 2003. Evaluation of a cefoxitin 30 µg disc on Iso-Sensitest agar for detection of methicillin-resistant *Staphylococcus aureus*. *J Antimicrob. Chemother*. 2003;52(2):204-7.
8. Brown DFJ, Edwards DI, Hawkey PM, Morrison D, Ridgway GL, Towner KJ, et al. Guidelines for the laboratory diagnosis and susceptibility testing of methicillin-resistant *Staphylococcus aureus*. *Jour Antimicrob Chemother*. 2005;56:1000-18.

9. Evans HL, Sawyer RG. Preventing bacterial resistance in surgical patients. *Surg Clin North Am*. 2009;89(2):501-19.
10. Barnes BA, Behringer GE, Wheelock FC, Wilkins EW. Postoperative sepsis: trends and factors influencing sepsis over a 20-year period reviewed in 20,000 cases. *Ann Surg* 1961;154(4):585-98.
11. Kurhade A, Akulwar S, Mishra M, Kurhade G, Justiz-Vaillant A, Kurhade K, et al. Bacteriological study of post-operative wound infections in a tertiary care hospital. *J Bacteriol Parasitol*. 2015;6(6):251
12. Barchitta M, Matranga D, Quattrocchi A, Bellocchi P, Ruffino M, Basile G, et al. Frequency of surgical site infections before and after the implementation of a multimodal infection control programme. *J Antimicrob Chemother*. 2012;67(3):749-55.
13. Rajkumari N, Sharma K, Mathur P, Kumar S, Gupta A. A study on surgical site infections after trauma surgeries in a tertiary care hospital in north India. *Indian J Med Res*. 2014;140(5):691-4.
14. Golia S, Kamath ASB, Nirmala AR. A study of superficial surgical site infections in a tertiary care hospital at Bangalore. *Int J Res Med Sci*. 2014;2(2):647-52.
15. Anvikar AR, Deshmukh AB, Karyakarte RP, Damle AS, Patwardhan NS, Malik AK, et al. A one-year prospective study of 3280 surgical wounds. *Indian J Med Microbiol* 1999;17(3):129-32.
16. Lilani SP, Jangale N, Chowdhary A, Daver GB. Surgical site infection in clean and clean-contaminated cases. *Indian J Med Microbiol*. 2005;23(4):249-52.
17. Sharan H, Misra AP, Mishra R. Determinants of Surgical Site Infection in rural Kanpur, India. *J Evol Med Dent Sci*. 2012;1(6):921-928.
18. deSa LA, Sathe MJ, Bapat RD. Factors influencing wound infection (a prospective study of 280 cases). *J Postgrad Med*. 1984;30(4):232-6.
19. Murthy R, Sengupta S, Maya N, Shivananda PG. Incidence of post-operative wound infection and their antibiogram in a teaching and a referral hospital. *Indian J Med Sci*. 1998;52(12):553-5.
20. Suchitra Joyce B, Lakshmidevi N. Surgical site infections: Assessing risk factors, outcomes and antimicrobial sensitivity patterns. *Afr J Microbiol Research*. 2009;3(4):175-9.
21. Awad SS, Elhabash SI, Lee L, Farrow B, Berger DH. Increasing incidence of Methicillin resistant *Staphylococcus aureus* skin and soft tissue infections: reconsideration of empiric antimicrobial therapy. *Am J Surg* 2007; 194(5): 606-10.
22. Majumder D, Bordoloi JS, Phukan AC, Mahanta J. Antimicrobial susceptibility pattern among methicillin resistant staphylococcus isolates in Assam. *Indian J Med Microbiol*. 2001;19(3):138-40.
23. Shobha KL, Rao PS, Thomas J. Survey of *Staphylococcus* isolates among hospital personnel, environment and their antibiogram with special emphasis on MRSA. *Indian J Med Microbiol* 2005; 23(3):186-8.
24. Vijayalaxmi N, Mohapatra LN, Bhujwala RA. Biological characters and antimicrobial sensitivity of *S. epidermidis* isolated from human source. *Indian J Med Res* 1980;72:16-22.
25. Marinello MA, Pierion C, Chenoweth C. The stethoscope, potential source of nosocomial infections? *Arch Inter Med* 1997;157:786-90.
26. Udaya Shankar C, Harish BN, Umesh Kumar PM, Navaneeth BV. Prevalence of MRSA in JIPMER hospital – a preliminary report. *Indian J Med Microbiol* 1997;15 (3):137-8.

Conflict of Interest: None Declared

Cite this article as:

Ram Nagina Sinha. A Prospective Bacteriological study of Surgical Site infections in a tertiary care hospital at Saharsa. *Asian Journal of Biomedical and Pharmaceutical Sciences*, 2013, 3: (18), 64-68.