

## The influence and significance of rapamycin on cardiomyocyte autophagy level of early heart failure rat.

Zhang Shuli<sup>\*#</sup>, Jianhui Zang<sup>#</sup>, Wang Moru

Department of Cardiology, People's Hospital of Laizhou City, Shandong Province, PR China

These authors contributed equally to this work

### Abstract

**Objective:** To explore the influences of rapamycin on cardiomyocyte autophagy level of early Heart Failure (HF) rat.

**Methods:** HF rat model was built by abdominal aorta surgery, and then they were randomly divided into the treatment group and the control group. The treatment group was given 3000 U/kg intraperitoneal injection of rapamycin, 3 times per week, for 4 weeks; the control group given the same amount of saline solution. Respectively 4 weeks, 8 weeks, 12 weeks after surgery (i.e., dosing 4 weeks), heart colour ultrasound examination was given for rat cardiac function. After 24 h fasting of all rats at the end of 12 weeks, rats were killed, then to observe the myocardial tissue cell morphology, cell apoptosis and Apoptosis Index (AI); Using Western blot method to detect myocardial *Beclin 1*, *Cathepsin D* protein expression.

**Results:** Echocardiography showed ventricular hypertrophy at 4 weeks, HF at 8 weeks in the model group; rapamycin intervention in the control group after 4 weeks, Left Ventricular Ejection Fraction (LVEF) increased significantly ( $P < 0.05$ ), interventricular septal thickness at end-systole (IVSs), left ventricular posterior wall thickness at end-systole (LVPWs), Interventricular Septum thickness at end-diastole (IVSd), Left Ventricular Posterior Wall Thickness at end-Diastole (LVPWd) significantly decreased after treatment ( $P < 0.05$ ). Compared with the sham group, *mTOR* expression decreased significantly, *cTnT* and *pmTOR* expression increased obviously in the control group ( $P < 0.05$ ); compared with the control group, *mTOR* expression increased significantly, *cTnT* and *pmTOR* expression decreased obviously in the treatment group ( $P < 0.05$ ). AI of the treatment group was significantly lower than Control ( $25.55\% \pm 2.56\%$  vs.  $34.66\% \pm 1.46\%$ ,  $P < 0.01$ ); compared with the control group ( $0.234 \pm 0.001$  and  $0.001 \pm 0.001$ ), *Beclin 1* and *Cathepsin D* OD value of the control group ( $0.867 \pm 0.005$  and  $0.867 \pm 0.003$ ) increased significantly, the treatment group ( $0.423 \pm 0.002$  and  $0.002 \pm 0.001$ ) was lower than the control group ( $P < 0.01$ ).

**Conclusion:** Rapamycin can effectively improve cardiac function in heart failure rats, inhibit cardiomyocyte autophagy.

**Keywords:** Rapamycin, Heart failure, Autophagy.

Accepted on December 30, 2016

### Introduction

Chronic Heart Failure (CHF) is the ultimate result of most Cardiovascular Disease (CVD), which influences patients' health even life severely, also brings about heavy economic burden on family and society [1]. Therefore, treatment on CHF is still a hot spot of research in cardiovascular field. Recent years, that rapamycin has the function of protecting cardiomyocyte has been found in pharmacological experiment and clinical experiment, and it has been applied to Drug Eluting Stent (DES) [2,3]. At present, a few of reports have been seen about the effects of Rapamycin treatment on CHF. However, the effect on Chinese citizens is still unknown. This paper is aimed at setting up rat model with CHF treated with

Rapamycin through observing Ultrasonic Cardiogram (UCG), autophagic rate of cardiomyocyte and the expression of *Beclin-1* and *Cathepsin D*, then to explore whether Rapamycin has the autophagic function of resisting cardiomyocyte of CHF or not in Chinese people, and its possible mechanism, finally it provides new thinking for treatment on CHF.

### Materials and Methods

#### Animals

Choosing of SPF level of 30 male SD rats, the average age was 8 to 12 weeks, the average weight was  $241.55 \pm 17.70$  g, which were bought from The Animal Center in West China Center of

Medical Sciences of Sichuan University. All rats were fed with food and water freely in same condition. Fodder was provided by The Animal Center in West China Center of Medical Sciences of Sichuan University. This research was approved by the Animal Committee of People's Hospital of Laizhou City.

### **Drug and reagent**

Rapamycin was bought from shenyang 3sBio Inc. TUNEL kits from Sigma company, *Beclin-1* and *Cathepsin D* antibody from Cell Signalling Technology (USA) and GAPDH antibody from ProteinTech Group, Inc. (USA).

### **Grouping and modelling**

30 rats were randomly divided into the model group (n=24) and the sham group (n=6). CHF model was built with abdominal aorta surgery in the model group. The specific methods as follows, water was forbade before 1d of surgery, 10% chloral hydrate (3 ml/kg) was given for anesthesia by intraperitoneal injection, the abdominal cavity was opened and veins in left kidney were exposed, abdominal aorta was separated bluntly about 10 to 15 mm apart from upper part above left kidney veins, then using 4-0 suture passed through abdominal aorta which has been separated, no.7 (diameter 0.7) needle was located beside abdominal aorta, which paralleled to abdominal aorta, after ligation, needle was pulled out, the tightness feeling was proper when needle extraction, then stenosis of abdominal aorta reached to 60% to 70%, finally abdominal cavity was closed. Amikacin was given for anti-infection 3 d by intraperitoneal injection after surgery, rats were observed and fed for 8 weeks, and then model was built. The sham group, the same part with the model group was given threading but no ligation, other groups were same with the model group.

### **Grouping and intervention after modelling**

16 rats survived in the model group after 8 weeks of surgery, then 16 rats were randomly divided into two groups: the treatment group (n=8); 10000 U/ml. Rapamycin was diluted into 1000 U/ml by normal saline, rats were given 3000 U/kg intraperitoneal injection of Rapamycin, 3 times per week, for 4 weeks. The control group (n=8) were given the same amount of saline solution by intraperitoneal injection, 3 times per week, for 4 weeks. The sham group was same with the control group. At the end of 12 weeks, 2 died in the control group, there was no death in the sham group and the treatment group.

### **Examination of heart colour ultrasound**

Respectively 4 weeks, 8 weeks, 12 weeks after surgery (i.e., dosing 4 weeks), heart colour ultrasound examination was given for rat cardiac function. After rats fasting for 24 h, 10% chloral hydrate (3 ml/Kg) was given for anesthesia and skin preparation. Using ultrasound cardiograph (i.e. 33 type, PHILIPS Company) and 7.5 HZ high frequency linear control probe (PHILIPS Company) to detect LVEF, IVSs, LVPWs, IVSd and LVPWd.

### **Intracardiac blood collection and sample disposal of myocardium**

At the end of 12 weeks, all rats were given UCG examination after 24 h fasting, 10% chloral hydrate (3 ml/Kg) was given for anesthesia and thoracotomy by intraperitoneal injection, collecting blood by intracardiac puncture, cutting myocardial tissue in apex of heart about 5 mm × 5 mm × 5 mm, which was put into precooling PBS solution to be washed to no red color, then located into ultra-low temperature freezer of -80°C after quick-freezed in liquid nitrogen for molecular biology experiment; other heart tissues were perfused by precooling saline solution for 3 min, then perfused and fixed by 4% Paraformaldehyde (PA) solution, the heart was took out and located into PA buffer solution to be fixed, and embedded with routine paraffin, then made into 3~4 μm slices of thick paraffin for stand-by use.

### **Pathological examination**

3 slices were randomly selected in each group from paraffin slices has been made, using HE staining and light microscope with 200 folds to observe morphological structure of myocardial tissue and condition of inflammatory cell infiltration in each group.

### **Detection on cell apoptosis by TUNEL**

3 slices were randomly selected in each group from paraffin slices has been made, detecting cardiomyocyte apoptosis according to operational approach of TUNEL kits, fluorescence coloration of cell apoptosis was observed under fluorescent microscope and image was adopted, then with DAB coloration, each slice randomly read 6 underlap visions under high power lens (X400), the positive cell was cell nucleus with brown color. After counting, AI=single vision positive apoptosis cell/single vision total cells × 100%, the average value was counted and analysed.

### **Detection of western blot on beclin-1 and cathepsin D protein expression**

Protein concentration of myocardial tissue was detected by ELISA after grinded and splited, total protein quality of upper sample according to protein concentration was 80 μg, after electrophoretic separation by SDS-PAGE hydrogel, using electro-blotting to transfer protein to PVDF membrane and Ponceaus to stain. Cutting membrane according to required target bands, after skim milk powder with 5% mass concentration of TBS-T solution closed at room temperature for 2 h, rabbit-anti *Beclin-1*, monoclonal antibody *Cathepsin D* and GAPDH antibody were added into TBS-T solution, incubated overnight under 4°C, after membrane washed by TBS-T solution at room temperature, rabbits (rats) secondary antibody incubation were added for 2 h, using TBS-T solution to wash membrane, then it reacted with ECL reagent, film was exposed and scanned, using Image J software to calculate OD value, then using OD ratio between AKT, P-AKT and GAPDH

bands to evaluate expression level of *Beclin-1*, *Cathepsin D* protein.

**Statistical treatment**

Using software spss 17.0 to do statistical analysis on data, all measurement data expressed by  $x \pm s$ , selecting chi-square test for enumeration data comparison and t-test, one-way ANOVA and SNK-q test for between-group average comparison, Spearman Wilcoxon correlation analysis for correlation analysis and dichotomy Logistic regression for multi-factor analysis. There has statistical difference ( $P < 0.05$ ).

**Results**

**Index changes of ultrasonic cardiogram before and after modelling and rapamycin intervention**

After 4 weeks of modelling, compared with LVEF of rats in the model group, there was no statistical difference ( $P > 0.05$ ); LVPWs, IVSs, LVPWd and IVSd increased obviously in the model group ( $P < 0.05$ ), it indicated myocardial hypertrophy of rats in the model group. After 8 weeks of modelling, LVEF of rats in the model group decreased significantly ( $P < 0.01$ ), LVPWs, IVSs, LVPWd and IVSd were still higher than the sham group ( $P < 0.05$ ), it indicated successful modelling of CHF rats in the model group. After 12 weeks of modelling ( the Rapamycin group given drugs for 4 weeks), compared with the sham group, LVEF of rats in the control group decreased furtherly ( $P < 0.05$ ), LVPWs, IVSs, LVPWd and IVSd increased furtherly ( $P < 0.05$ ), it indicated CHF became more severe of rats in the control group; LVEF of rats increased obviously ( $P < 0.05$ ) and LVPWs, IVSs, LVPWd and IVSd decreased obviously ( $P < 0.05$ ) in the treatment group compared with the control group, it indicated Rapamycin can improve cardiac function of rats with heart failure and reverse ventricular hypertrophy (Table 1).

**Table 1.** Comparison of echocardiography 4, 8 and 12 weeks of operation.

Group	Weeks	n	LVPWs (cm)	IVSs (cm)	LVPWd (cm)	IVSd (cm)	LVEF
Sham	4	6	0.277 ± 0.007	0.287 ± 0.005	0.146 ± 0.005	0.177 ± 0.006	0.878 ± 0.001
Model	4	16	0.357 ± 0.007	0.367 ± 0.004	0.246 ± 0.006	0.288 ± 0.007	0.846 ± 0.005
t-value			5.455**	2.565*	3.566*	2.577*	1.576
Sham	8	6	0.277 ± 0.003	0.346 ± 0.007	0.175 ± 0.007	0.178 ± 0.001	0.845 ± 0.006
Model	8	16	0.368 ± 0.001	0.377 ± 0.008	0.277 ± 0.001	0.246 ± 0.004	0.623 ± 0.001
t-value			5.234**	3.245**	3.135**	2.535*	8.355**
Sham	12	6	0.335 ± 0.001	0.345 ± 0.017	0.245 ± 0.013	0.199 ± 0.001	0.824 ± 0.134
Control	12	6	0.445 ± 0.002	0.394 ± 0.013	0.345 ± 0.011	0.245 ± 0.012	0.624 ± 0.041

Treatment	12	8	0.345 ± 0.002	0.374 ± 0.023	0.234 ± 0.025	0.215 ± 0.001	0.714 ± 0.001
t-value			10.355**	5.456*	13.245**	7.345**	32.876**

\* $P < 0.0$ , \*\* $P < 0.01$ ; <sup>a</sup>Compared with Sham group, <sup>b</sup>compared with Control group,  $P < 0.05$ .

**Expression level comparison of cTnT, mTOR and pmTOR**

Compared with the sham group, *mTOR* expression decreased significantly, *cTnT* and *pmTOR* expression increased obviously in the control group ( $P < 0.05$ ); compared with the control group, *mTOR* expression increased significantly, *cTnT* and *pmTOR* expression decreased obviously in the treatment group ( $P < 0.05$ ) (Table 2).

**Table 2.** Comparison of the expression of *cTnT*, *mTOR*, *pmTOR* between three groups.

Group	n	cTnT (ng/ml)	mTOR	pmTOR
Sham group	6	9.554 ± 0.005	1.151 ± 0.001	1.245 ± 0.023
Control group	6	13.367 ± 0.005	1.381 ± 0.003	0.566 ± 0.006
F		11.355**	12.455**	13.565**

**Changes of cardiomyocyte autophagy**

Cell nucleus was clearly visible, relatively big, the border of it was clear and stained to blue in the sham group. There was a large quantity bubble-like structure, which was specific autophagic vacuole in autophagic cardiomyocyte. Phagocytic vacuole and autophagic vacuole decreased obviously in the treatment group compared with the control group. Compared with the sham group, the autophagic rate of cardiomyocyte increased significantly in the control group; compared with the control group, the autophagic rate of cardiomyocyte decreased in the treatment group, but it was still higher than the sham group ( $P < 0.01$ ), Table 3 as follows.

**Detection of western blot on beclin-1 and cathepsin d protein expression of cardiomyocyte**

Compared with the sham group, *Beclin-1* and *Cathepsin D* expression increased obviously in the control group; compared with the control group, *Beclin-1* and *Cathepsin D* expression increased obviously in the treatment group, but it was still higher than the sham group ( $P < 0.01$ ), Table 3 as follows.

**Table 3.** Comparison of the expression of *Beclin-1*, *Cathepsin D* protein (OD) and myocardial Autophagy Index (AI) between three groups.

Group	n	Beclin-1	Cathepsin D	AI (%)
Sham group	6	0.234 ± 0.001	0.331 ± 0.001	8.45 ± 0.89
Control group	6	0.867 ± 0.005	0.821 ± 0.003	34.66 ± 1.46
Treatment group	8	0.423 ± 0.002	0.655 ± 0.001	25.55 ± 2.56

F	13.455**	14.458**	53.541**
---	----------	----------	----------

## Discussion

Heart Failure (HF) is the ultimate result of myocardial damage caused by multiple CVD. Its pathophysiological changes are very complex. Recent years, study finds that cardiomyocyte autophagy plays an important role in occurrence and development of HF [4-6]. Autophagy is a highly conserved cellular mechanism of intracellular protein and organelle recycle mediated by lysosome. Autophagy can be induced by pathological factors of intracellular anoxia, starvation, pressure and stress. But in recent years, the role of uncontrolled factor in occurrence and development of many diseases begin to be taken seriously [7-10]. Cardiomyocyte autophagy can cause reduction of cardiomyocyte, thinness of ventricular wall and enlargement of ventricular cavity, making HF get worse.

Study finds that mTOR is the key regulatory protein of autophagic regulatory pathway of mammals [10-12]. Study shows that when myocardium is hypoxic-ischemic and phosphorylation of *mTOR*, then forming *pmTOR*, which activates cardiomyocyte autophagy and causes cardiomyocyte autophagy furtherly [12-15]. Autophagy can protect cardiomyocyte; however, HF will cause excessive autophagy, finally causing injury of cardiomyocyte [16]. The results show, compare with the sham group, mTOR expression decreases significantly, *cTnT* and *pmTOR* expression increase obviously in the control group ( $P < 0.05$ ); compare with the control group, *mTOR* expression increases significantly, *cTnT* and *pmTOR* expression decrease obviously in the treatment group ( $P < 0.05$ ), it shows that Rapamycin can inhibit excessive autophagy of cardiomyocyte through lowering *pmTOR* expression to a certain degree.

This study shows, cardiac function of HF rats decreases, and autophagic relevant protein of *Beclin-1*, *Cathepsin D* expression of myocardial tissue increase significantly compare with the sham group, after Rapamycin treatment, cardiac function of HF rats improves greatly. And autophagic relevant protein of *Beclin-1*, *Cathepsin D* expression decrease significantly. The results point out autophagy plays an important role in occurrence and development of HF; Rapamycin can relieve cardiomyocyte autophagy of HF and improve cardiac function. Therefore, autophagy may become potential target for HF treatment; regulating myocardial cell autophagy is new strategy for HF treatment. Regulating uncontrollable cardiomyocyte autophagy of HF can reduce cardiac cell death caused by autophagy, which will be benefit for HF treatment. At present, the relevant mechanism of overloaded cardiac pressure causing cardiomyocyte autophagy is still not clear, which needs further study [17-20]. Searching for a kind of drug which can be applied to clinic and regulate cardiomyocyte autophagy, which may be benefit for HF treatment.

## References

- Braunwald E. The war against heart failure: the Lancet lecture. *Lancet* 2015; 385: 812-824.
- Qin YJ, Zhang XL, Yu YQ. Cardioprotective effect of erythropoietin on sepsis-induced myocardial injury in rats. *World J Emerg Med* 2013; 4: 215-222.
- Sanchis-Gomar F, Garcia-Gimenez JL, Pareja-Galeano H. Erythropoietin and the heart: physiological effects and the therapeutic perspective. *Int J Cardiol* 2014; 171: 116-125.
- Bogoyevitch MA. An update on the cardiac effects of erythropoietin cardioprotection by erythropoietin and the lessons learnt from studies in neuroprotection. *Cardiovasc Res* 2004; 63: 208-216.
- Kowalczyk M, Banach M, Mikhailidis DP. Erythropoietin update 2011. *Med Sci Monit* 2011; 17: A240- A247.
- Santhanam A, D'Uscio LV, Katusic ZS. Cardiovascular Effects of Erythropoietin: An Update. *Adv Pharmacol* 2010; 60: 257- 285.
- Gao S, Long CL, Wang RH. K(ATP) activation prevents progression of cardiac hypertrophy to failure induced by pressure overload via protecting endothelial function. *Cardiovasc Res* 2009; 83: 444-456.
- Zhang Y, Zhang L, Yan M. Inhibition of phosphatidylinositol 3-kinase causes cell death in rat osteoblasts through inactivation of Akt. *Biomed Pharmacother* 2007; 61: 277-284.
- Zhu M, Feng J, Lucchinetti E. Ischemic postconditioning protects remodeled myocardium via the PI3K-PKB/Akt reperfusion injury salvage kinase pathway. *Cardiovasc Res* 2006; 72: 152-162.
- Mudalagiri NR, Mocanu MM, Di Salvo C. Erythropoietin protects the human myocardium against hypoxia/reoxygenation injury via phosphatidylinositol-3 kinase and ERK1/2 activation. *Br J Pharmacol* 2008; 153: 50-56.
- Uchiyama T, Engelman RM, Maulik N. Role of Akt signalling in mitochondrial survival pathway triggered by hypoxic preconditioning. *Circulation* 2004; 109: 3042-3049.
- Lee J, Jung K, Kim YS. Diosgenin inhibits melanogenesis through the activation of phosphatidylinositol-3-kinase pathway (PI3K) signalling. *Life Sci* 2007; 81: 249- 254.
- Zakhidova KKh. Correlation between concentration of pathological cytokines and erythropoietin in patients with chronic heart failure with anaemic syndrome. *Vestn Ross Akad Med Nauk* 2014; 32-37.
- Zhou S, Zhuang Y, Zhao W. Protective roles of erythropoiesis stimulating proteins in chronic heart failure with anemia. *Exp Ther Med* 2014; 8: 863-870.
- Zhang J, He Z, Xiao W, Na Q, Wu T, Su K, Cui X. Overexpression of BAG3 attenuates hypoxia-induced cardiomyocyte apoptosis by inducing autophagy. *Cell Physiol Biochem* 2016; 39: 491-500.
- Gui L, Liu B, Lv G. Hypoxia induces autophagy in cardiomyocytes via a hypoxia-inducible factor 1-dependent mechanism. *Exp Ther Med* 2016; 11: 2233-2239.

17. Ji W, Wei S, Hao P, Xing J, Yuan Q, Wang J, Xu F, Chen Y. Aldehyde dehydrogenase 2 has cardioprotective effects on myocardial ischaemia/reperfusion injury via suppressing mitophagy. *Front Pharmacol* 2016; 7: 101.
18. Zeng C, Li H, Fan Z, Zhong L, Guo Z, Guo Y, Xi Y. Crocin-elicited autophagy rescues myocardial ischemia/reperfusion injury via paradoxical mechanisms. *Am J Chin Med* 2016; 44: 515-530.
19. Zhu T, Yao Q, Wang W, Yao H, Chao J. iNOS induces vascular endothelial cell migration and apoptosis via autophagy in ischemia/reperfusion injury. *Cell Physiol Biochem* 2016; 38: 1575-1588.
20. Mo Y, Tang L, Ma Y, Wu S. Pramipexole pretreatment attenuates myocardial ischemia/reperfusion injury through

upregulation of autophagy. *Biochem Biophys Res Commun* 2016; 473: 1119-1124.

**\*Correspondence to**

Zhang Shuli  
Department of Cardiology  
People's Hospital of Laizhou City  
Shandong Province  
PR China