

The relationship between total standardized mini mental state examination (SMMSE) and subscores and D vitamin, Folate and B12 levels in patients with cognitive dysfunction.

Sule Aydin Turkoglu¹, Muhammed Nur Ogun^{1*}, Evgin Micoogullari², Edip Gultekin¹, Serpil Yildiz¹

¹Department of Neurology, Faculty of Medicine, Abant Izzet Baysal University, Bolu, Turkey

²Department of Psychology, Faculty of Sciences, Abant Izzet Baysal University, Bolu, Turkey

Abstract

Introduction: There is evidence that low vitamin levels may be associated with cognitive functions, results from studies on this issue have been incoherent. This study aimed to evaluate whether there is any relationship between the levels of Vitamin D, Folate and B12 and Standardized Mini-Mental Test (SMMT) and subgroups in patients with cognitive dysfunction.

Methods: This retrospective study included 184 women and 89 men with cognitive problems and who had SMMSE scores of 24/30 or less. SMMSE scores were prospectively recorded in the patient's file for each patient by an experienced neuropsychologist. SMMSE scores were prospectively recorded in the patient's file for each patient by an experienced neuropsychologist. Vitamin B12, vitamin D and Folate levels were studied from 10:00 am fasting blood samples of each patient and prospectively recorded in patient file. Patient files were retrospectively reviewed and data were recorded.

Results: No correlation was found between vitamin D and vitamin B12 levels and SMMSE (total and sub scores) in both sexes. Folate levels were significantly correlated with the SMMSE total ($p=0.005$, $r=0.288$), orientation ($p=0.004$, $r=0.290$) and registration ($p=0.04$, $r=0.215$) scores in females. A positive correlation was found between the folate levels and the SMMSE recall sub score in both sexes ($p=0.002$, $r=0.227$).

Conclusions: Folate deficiency is an easily treatable condition. Thus it should be considered in the differential diagnosis of cognitive impairment especially in female patients.

Keywords: Cognitive disjunction, SMMSE, Folate, Vitamin B12, Vitamin D.

Accepted on October 16, 2017

Introduction

There are many studies that point out to the fact that group B vitamins, vitamin D, folic acid, antioxidant vitamins, essential minerals and essential fatty acids play a key role in cognitive function [1,2]. In a pilot study, it was shown that the concentration of many vitamins in AD patients with moderate stage dementia without vascular disease and weight loss were significantly reduced compared to healthy controls. This suggests that regulation of nutrition and specific vitamin intake may prevent brain damage and dementia [3].

The Standardized Mini-Mental State Examination (SMMSE) is one of the main methods to evaluate cognitive functions. SMMSE has been developed primarily to assess delirium and dementia with high sensitivity and specificity [4]. Subsequently, SMMSE has been extended for use in global cognitive impairment. The maximum total score is 30 and the limit value for the cognitive impairment is 24 [5]. The aim of this study is to investigate the relationship between the levels of Vitamin D, Folate and B12 and Standardized Mini-Mental

Test (SMMT) and its sub scores in patients with cognitive dysfunction.

Material and Methods

Study design and participants

This retrospective study was conducted using data of patients who were admitted to Abant Izzet Baysal University, Department of Neurology between January 2015 and March 2017 with cognitive complaints who had SMMSE scores of 24/30 or less were prospectively and consecutively recorded. Of the 184 patients, 95 were females and 89 were males. Patients with vascular dementia, frontotemporal dementia, Lewy body dementia, psychotic findings, and severe depression were excluded from the study. SMMSE scores were prospectively recorded in the patient's file for each patient by an experienced neuropsychologist. Vitamin B12, vitamin D and Folate levels were studied from 10:00 am fasting blood samples of each patient and prospectively recorded in patient file. After obtaining the necessary institutional permission for

the use of patient data, patient files were retrospectively reviewed and data were recorded. The study was approved by the local Ethics Committee (no:2013/136) and conducted in accordance with the Declaration of Helsinki.

Statistical analysis

In the study, the data was analyzed by using SPSS 22.0 for windows package program. The chi-square test was used to evaluate the difference between the categorical variables such as gender and educational status between the groups. Normal distribution of vitamin levels was tested visually (histogram and probability plots) and by analytical (Kolmogorov-Smirnov/ Shapiro-Wilk) methods. Spearman correlation analysis was used for correlations between variables when at least one of the SMMSE and vitamin levels did not meet the normal distribution.




Figure 1. Correlation between folate levels and SMMSE total, orientation, registration and recall sub scores in females.




Figure 2. Correlation between folate levels and SMMSE recall sub score in both sexes.

Results

In the present study, of a total of 184 patients, 89 were males and 95 were females. The educational level of female patients was significantly lower than in males ($p < 0.001$) (Table 1). The mean age was 69 (range: 32 to 91) years in males and 68 (range: 31 to 90) years in females. There was no statistically significant difference regarding age between females and males. ($p = 0.7$). Vitamin D and folate levels were statistically significantly lower in females than males ($p < 0.001$). Vitamin B12 levels were within normal ranges in both sexes. There was no significant difference in vitamin B12 levels between males and females ($p = 0.9$). Vitamin D levels were 12 ng/mL (2-45) below the normal value in both groups, more pronounced in females. There was no significant difference in total SMMT, orientation, record-memory, attention orientation, language and recall scores in males and females (Table 2). In the Spearman's Correlation test, no correlation was found between vitamin D and vitamin B12 levels and SMMSE (total and sub scores) in both sexes. In females, folate levels were significantly correlated with the SMMSE total ($p = 0.005$, $r = 0.288$), orientation ($p = 0.004$, $r = 0.290$) and registration ($p = 0.04$, $r = 0.215$) scores. There was a positive correlation between the folate levels and the SMMSE recall sub score in both sexes ($p = 0.002$, $r = 0.227$) (Figures 1 and 2).

Table 1. Comparison of educational status between males and females.

		Educational Status		
		Educated	Uneducated	p
Sex	Female	37	58	<0.001
	Male	9	80	
Total		46	138	

Table 2. Comparison of vitamin D, Folate and vitamin B 12 levels and Standardized Mini-Mental State Examination sub scores between males and females.

	Female (n=95)	Male (n=89)	p
Vitamin D ng/mL	9 (2-36)*	15(4-45)	$p < 0.01$
Folat ng/mL	6 (2-16)	6 (2-14)	$p = 0.02$
Vitamin B12 pg/mL	282 (111-932)	322(83-732)	$p = 0.9$
SMMT			
SMMT total	22 (8-29)	22(9-29)	$p = 0.9$
SMMT orientation	7 (1-10)	7 (1-10)	$p = 0.08$
SMMT registration	3 (1-3)	3 (1-3)	$p = 0.7$
SMMT attention and calculation	4 (0-5)	4 (0-5)	$p = 0.9$
SMMT recall	1 (0-3)	1 (0-3)	$p = 0.4$
SMMT language	7 (2-9)	7 (3-9)	$p = 0.4$
SMMSE: Standardized Mini-Mental State Examination			

*med (min-max)

Discussion

In this study, correlations between serum Vitamin D, Vitamin B12 and folate levels and standardized mini mental state results in elderly subjects were examined. A significant correlation was found between serum folate level and cognitive impairment, especially in women. SMMSE total score was correlated with the folate levels in both sexes, and correlations were also found with orientation, registration and recall scores from SMMSE subgroups in females.

In several studies it has been shown that folate and Vitamin B12 levels are inversely correlated to the clinical condition in patients with Alzheimer's disease-related dementia [6-11]. Similar correlation has also been demonstrated in elderly individuals with cognitive impairment. Although the underlying mechanism of cognitive dysfunction is not clearly understood, it is considered that low Vitamin B12 and folate levels influence the methylation reactions or homocysteine levels in the brain according to the hypothesis established based on the results of some studies. It is also considered that homocysteine may lead cell death by activating N methyl D aspartate receptors (NMDA) or may cause an exotoxic effect by converting to homocysteine acid [12-16]. In a study conducted by Goodwin et al., nonverbal abstract thinking skills and memory scores in neuropsychological tests were found to be significantly lower in individuals with low folate levels than those in normal folate levels [17]. Ma et al. have shown that a daily oral administration of a 400- μ g folic acid supplement to the patients with mild cognitive impairment (MCI) for 12 months may significantly improve cognitive performance and reduce peripheral inflammatory cytokine levels [18]. Ulstein et al. investigated nutritional factors among home-living patients diagnosed with MCI or Alzheimer's disease-related mild dementia, compared to healthy subjects. They found no associations between vitamin levels and early cognitive impairment in patients with Alzheimer's disease [19]. Riggs et al. reported that individuals who received vitamin C, folate and vitamin B12 supplementation had lower scores than those who did not [14]. It has been reported that low level of vitamin D is associated with low SMMSE scores and low memory testing in subjects with normal vitamin B12 levels [20]. In a prospective cohort study conducted by Schneider et al., no association was found between vitamin D deficiency and neurodegenerative disorders [21]. In our study, we also did not find any significant correlation between vitamin D and SMMSE scores. The patients included in the study were all recruited from the same neuropsychologist and underwent a comprehensive standardized assessment. There were few limitations in our study. Serum homocysteine concentrations were not measured. The number of participants was too small to examine the correlation between vitamin levels and cognitive scores.

Conclusion

In this study, no significant correlation was found between Vitamin B12 and Vitamin D levels and cognitive status, unlike many previous studies. A significant correlation was found between low folate levels and cognitive status. Folate consumption has important implications on cognitive functions. The cognitive impairment due to folate deficiency is a condition that may easily correct. Folate deficiency should be considered in the differential diagnosis of cognitive impairment especially in female patients. Further studies are needed to determine whether folate sufficiency might have a role in cognitive functions.

References

1. Coley N, Vaurs C, Andrieu S. Nutrition and cognition in aging adults. *Clin Geriatric Med* 2015; 31: 453-464.
2. Amraei M, Mohamadpour R, Moayeri A, Abbasi N, Shirzadpour E, Mohamadpour M. Vitamin D and its association with memory and learning: A systematic review and meta-analysis. *Biomed Res* 2017; 28: 7427-7433.
3. Glasø M, Nordbo G, Diep L, Bøhmer T. Reduced concentrations of several vitamins in normal weight patients with late-onset dementia of the Alzheimer type without vascular disease. *J Nutri Health Aging* 2004; 8: 407-413.
4. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975; 12: 189-198.
5. Nys G, Van Zandvoort M, De Kort P, Jansen B, Kappelle L, De Haan E. Restrictions of the Mini-Mental State Examination in acute stroke. *Arch Clin Neuropsychol* 2005; 20: 623-629.
6. Nilsson K, Gustafson L, Fäldt R, Andersson A, Brattström L, Lindgren A. Hyperhomocysteinaemia-a common finding in a psychogeriatric population. *Euro J Clin Invest* 1996; 26: 853-859.
7. Joosten E, Lesaffre E, Riezler R, Ghekiere V, Dereymaeker L, Pelemans W. Is metabolic evidence for vitamin B-12 and folate deficiency more frequent in elderly patients with Alzheimer's disease? *J Gerontol* 1997; 52: M76-M9.
8. Renvall MJ, Spindler AA, Ramsdell JW, Paskvan M. Nutritional status of free-living Alzheimer's patients. *Am J Med Sci* 1989; 298: 20-27.
9. Kristensen MO, Gulmann NC, Christensen JE, Ostergaard K, Rasmussen K. Serum cobalamin and methylmalonic acid in Alzheimer dementia. *Acta Neurol Scand* 1993; 87: 475-481.
10. Ball MJ, Fisman M, Hachinski V, Blume W, Fox A, Kral VA, Kirshen AJ, Fox H, Merskey H. A new definition of Alzheimer's disease: a hippocampal dementia. *Lancet* 1985; 1: 14-16.
11. Karnaze DS, Carmel R. Low serum cobalamin levels in primary degenerative dementia. Do some patients harbor

- atypical cobalamin deficiency states? Arch Intern Med 1987; 147: 429-431.
12. Carmel R. Cobalamin, the stomach, and aging. Am J Clin Nutr 1997; 66: 750-759.
 13. Carmel R, Green R, Jacobsen DW, Rasmussen K, Florea M, Azen C. Serum cobalamin, homocysteine, and methylmalonic acid concentrations in a multiethnic elderly population: ethnic and sex differences in cobalamin and metabolite abnormalities. Am J Clin Nutrition 1999; 70: 904-910.
 14. Riggs KM, Spiro A, Tucker K, Rush D. Relations of vitamin B-12, vitamin B-6, folate, and homocysteine to cognitive performance in the Normative Aging Study. Am J Clin Nutrition 1996; 63: 306-314.
 15. Seshadri S, Beiser A, Selhub J, Jacques PF, Rosenberg IH, D'Agostino RB, Wilson PW, Wolf PA. Plasma homocysteine as a risk factor for dementia and Alzheimer's disease. N Engl J Med 2002; 346: 476-483.
 16. Clarke R, Smith AD, Jobst KA, Refsum H, Sutton L, Ueland PM. Folate, vitamin B12, and serum total homocysteine levels in confirmed Alzheimer disease. Arch Neurol 1998; 55: 1449-1455.
 17. Goodwin JS, Goodwin JM, Garry PJ. Association between nutritional status and cognitive functioning in a healthy elderly population. JAMA 1983; 249: 2917-2921.
 18. Ma F, Wu T, Zhao J, Song A, Liu H, Xu W. Folic acid supplementation improves cognitive function by reducing the levels of peripheral inflammatory cytokines in elderly Chinese subjects with MCI. Sci Rep 2016.
 19. Ulstein I, Bøhmer T. Normal vitamin levels and nutritional indices in Alzheimer's disease patients with mild cognitive impairment or dementia with normal body mass indexes. J Alzheimer's Dis 2017; 55: 717-725.
 20. Przybelski RJ, Binkley NC. Is vitamin D important for preserving cognition? A positive correlation of serum 25-hydroxyvitamin D concentration with cognitive function. Arch Biochem Biophys 2007; 460: 202-205.
 21. Schneider AL, Lutsey PL, Alonso A, Gottesman RF, Sharrett AR, Carson KA. Vitamin D and cognitive function and dementia risk in a biracial cohort: the ARIC Brain MRI Study. Euro J Neurol 2014; 21: 1211.

***Correspondence to**

Muhammed Nur Ogun
Department of Neurology
Faculty of Medicine
Abant Izzet Baysal University
Turkey