

Transstadial transmission and maintenance of *Francisella tularensis* in the Blacklegged Tick *Ixodes scapularis*.

Rinosh J Mani^{1,2*}, Kenneth D Clinkenbeard¹

¹Department of Veterinary Pathobiology, Center for Veterinary Health Sciences, Oklahoma State University, Stillwater, OK 74078 United States of America

²Veterinary Diagnostic Laboratory, Michigan State University, Lansing, MI 48910, United States of America

Abstract

In the south-central United States tularemia endemic region, approximately 10% of human cases occur during the fall-winter season when the questing activity of its primary tick vectors for this region *Dermacentor variabilis* and *Amblyomma americanum* is minimal. It is also been reported that approximately 40% of tularemia infected domestic cats contract the disease during the fall-winter season. To more fully characterize the potential role of *I. scapularis*, whose host seeking activity peaks during the fall-winter season in the south-central United States, as a vector for tularemia, the biology of *F. tularensis* subspecies *holarctica* strain LVS in colony-reared *I. scapularis* artificially infected with LVS by capillary feeding was studied. Capillary-fed *I. scapularis* larva and nymph were initially colonized with 1.9×10^3 and 1.1×10^4 CFU/tick, respectively. For nymphs molted from colonized larvae, the percent of molted nymphs colonized declined at the time of molting, but LVS persisted in 15% of molted nymphs up to 175 days post-capillary feeding. In contrast, only 5% of adults molted from infected nymphs maintained LVS colonization through 150 days post-capillary feeding. For capillary-fed adults, LVS initially colonized the gut and disseminated to hemolymph and salivary glands by 14 day post-feeding. Post-capillary feeding larvae and nymphs maintained LVS colonization initially and persisted although in a low percentage in molted nymphs and molted adults. Based on the feeding habits of *I. scapularis* adults, this tick species has the potential to act as a bridging vector for human tularemia for fall-winter cases of tularemia.

Keywords: *Ixodes scapularis*, *Francisella tularensis*, Tularemia, Transstadial transmission.

Accepted August 16, 2018

Introduction

Francisella tularensis is a highly infectious gram-negative coccobacillus which causes the disease tularemia in humans and a wide range of vertebrate and invertebrate animals [1]. There are two main subspecies of this bacterium, namely subspecies *tularensis* exclusively seen in the North America and subspecies *holarctica* seen in North America as well as in Eurasia [2,3]. Molecular subtyping have identified a number of genotypes for the two subspecies of which genotype A1b seems to be the most virulent with a case fatality rate of 25% and is associated with tick borne infections [4-6]. Historically, human tularemia cases were associated with handling of dead/sick rabbits; however, in recent years it is manifested as a tick borne disease where the ulceroglandular form of the disease predominates [2,7-9].

In the south-central tularemia endemic region of United States, the tularemia agent *Francisella tularensis* is maintained via a spring-summer tick-small mammalian enzootic cycle coincident with questing activity of its two primary tick vectors for this region *Dermacentor variabilis* and *Amblyomma americanum* [2,10]. Likewise, the majority of the sporadic human tularemia cases exhibit a similar spring-summer seasonal pattern; however, approximately 10% of human tularemia in Oklahoma occurs during the months of November through February when its two primary tick vectors are the least active [11,12]. Many

of the cases of human tularemia during this period are attributed to direct transmission from infected rabbits associated with hunting activities [11,12]. However, Hopla noted that some cases of human tularemia during the months when its primary tick vectors are inactive are nevertheless associated with tick transmission [13].

Ixodes scapularis is the third most abundant tick species in the south-central United States. The host preference of adult *I. scapularis* for large mammals including humans and the coincidence of questing activity with the fall-winter incidence of human tularemia supports its potential as a vector [14]. Domestic cats are highly susceptible to tularemia and the disease incidence in cats shows a bimodal pattern of spring-summer and fall maxima [3]. Cats contract the disease by preying on tularemia infected small mammals and the disease during the spring-summer and fall peak incidence is indicative of concurrent tularemia infection cycle in small mammals [3]. Immature stages of *I. scapularis* seeks small mammals including rabbits and white footed mice as hosts; however, the host seeking activity of the immature *I. scapularis* is predominantly during summer months in the south-central tularemia endemic region [14].

To more fully characterize the potential for *I. scapularis* to serve as a vector for tularemia, colony-reared *I. scapularis*

larvae, nymphs, and adults were infected via artificial capillary tube feeding of tick meal containing *F. tularensis* subspecies *holarctica* strain LVS to generate sufficient number of infected ticks to allow assessment of quanta and duration of colonization, transstadial transmission, tissue distribution, and secretion in tick saliva.

Materials and Methods

Ticks, bacterial strain and growth conditions

I. scapularis larvae, nymphs, and adults were obtained from the Tick Rearing Facility, Department of Entomology and Plant Pathology, Oklahoma State University (Stillwater, OK). Larvae and nymphs were collected after they were fed to repletion on rabbits. Unfed flat adult ticks were used for determination of quanta and tissue distribution of LVS. Adult ticks used for salivary induction experiments were partially fed on sheep for five to six days. *Francisella tularensis* subsp. *holarctica* strain LVS was supplied by the Oklahoma State Department of Health. All chemicals used in the study were purchased from Sigma (St Louis, MO) unless indicated otherwise.

Capillary tube feeding of larvae, nymphs and adult ticks

The ticks were surface disinfected by washing in 30% hydrogen peroxide, distilled water and 70% isopropyl alcohol for 5 seconds each. After the washing, adult ticks were immobilized dorsal side up on adhesive part of duct tape placed over double-sided adhesive tape (3M Scotch brand, St Paul, MN) in a 100 by 15 mm Petri dish base. Ticks were then further immobilized by applying single-sided adhesive tape over 1/4th of the caudal region. Larvae and nymphs were immobilized with their dorsal side down on double-sided adhesive tape on dental wax (Electron Microscopy Sciences, Fort Washington, PA). For capillary tube feeding, 10 µL (internal diameter of 0.55 mm) glass capillary tubes (Drummond Scientific Company, Broomall, PA) were used for larvae and nymphs while 35 µL (internal diameter of 0.79 mm) glass capillary tubes were used for adult ticks. The tick meal inoculum contained 1-10 10⁷ CFU/mL *Francisella*

tularensis subsp. *holarctica* strain LVS. Detailed description of the preparation of inocula and growth of LVS has been described earlier [15]. After feeding, ticks were either surface disinfected by washing in 30% hydrogen peroxide, distilled water and 70% isopropyl alcohol for 5 seconds and minced for determination of CFUs or were maintained in 8 oz. paper cup containers with plastic lids and kept in a humidity chamber (relative humidity of > 90%) at 22 ± 1°C (unless specified otherwise) with automated artificial lighting to simulate a 12 h day-night cycle. Determination of CFU per tick was done at one day post-capillary feeding and then at an interval of 7 days. To determine the bacterial numbers in ticks at various times of colonization, individual ticks were minced with a scalpel blade, bacteria extracted by incubation in PBS containing 64 µg/mL ampicillin for 2 h at room temperature on a rotor platform mixer (Boekel Scientific, Feasterville, PA.), serially diluted in PBS containing 64 µg/mL ampicillin, and plated on chocolate agar plates for CFU determination following incubation at 37 °C in 5% CO₂ for 72 h. To determine the bacterial numbers in tissues at varying periods of post-capillary tube feeding, individual ticks were dissected into gut and salivary glands under sterile conditions using a dissecting microscope. The tick tissues were placed in PBS containing 64 µg/mL ampicillin and processed as described above for whole ticks. Hemolymph from ticks was collected into sterile glass capillary tubes from the cut ends of the tick's leg. Figure 1 schematically represents the details of the capillary feeding protocol including the number of ticks used and timeline of feeding. On days one through 175 post-capillary feeding, batches of 15 or more ticks were assayed individually for LVS.

Real-time quantitative PCR

Real-time quantitative PCR was done to determine presence of viable, but non-culturable LVS or LVS DNA in ticks after capillary feeding compared to those which were positive by microbial culture. Real-time PCR setup has been described earlier [16].




Figure 1. Quanta and duration of colonization and transstadial transmission of LVS experimental design.

Larvae, nymphs and adults *I. scapularis* in batches of various sizes were capillary tube fed tick meal containing LVS. The overall rate of ticks positive for LVS for each tick life cycle stage capillary tube fed LVS is given as total number of ticks which were positive for LVS (+LVS)/total number of ticks tested.

Intra-hemocoelic inoculation and salivary induction in ticks

To determine the presence of LVS in tick saliva, partially fed adult ticks were inoculated with 1 μ L of inoculum containing 1×10^8 CFU/mL of *F. tularensis* subsp. *holarctica* strain LVS in PBS injected intra hemocoel in the ventral region of the tick, medial to the caudal most coxa using 10 μ L custom made Hamilton syringe with a 0.5 inch, 33 gauge needle (Hamilton Company, Reno, NV). For inducing salivary secretion in ticks that were colonized with LVS via capillary tube-feeding, partially fed female ticks were fed with LVS (1×10^7 CFU/mL) via capillary tube and were held in humidity chamber at 27°C. Two days post-capillary tube feeding, the ticks were induced to secrete saliva. Partially-fed LVS-colonized adult ticks were immobilized dorsal side up on the sticky part of duct tape placed on a double-sided adhesive tape. Chemical induction and collection of saliva has been described earlier [15,16].

Statistical analysis

F. tularensis colonization in different groups of *I. scapularis* ticks during transstadial transmission from larva to nymphs, and nymph to adult were compared by using 1-way analysis of variance on log-transformed data followed by pairwise multiple

comparison of mean CFU value using Holm-Sidak tests. Overall significance level for Holm-Sidak tests was $P=0.05$. All statistical analyses were performed with SigmaPlot v11.0 software package (Systat Software Inc., Chicago, IL).

Results

LVS colonization of *I. scapularis* larvae and transstadial transmission to molted nymphs

At one-day post-capillary tube feeding, 100% of larvae exhibited colonization as detected by microbial culture with a mean colonization level of $1.9 \pm 1.8 \times 10^3$ CFU/larva (Figure 2; limit of detection was 2 CFU/larva). The colonization level per larvae increased at 7 days post-capillary feeding to $5.3 \pm 7.3 \times 10^4$ CFU/larva (unadjusted $P < 0.01$), suggesting that the bacterium had proliferated following the initial ingestion. For our experimental holding conditions (RH=95% and mean room temperature of $22 \pm 1^\circ\text{C}$), the larvae molted to nymphs between days 35 to 42 post-capillary tube feeding (Figure 2A). A significant decline in the percent of larvae colonized was observed with the onset of molting at 35 days, declining markedly to 0 to 25% for sampling days 56 through 175 post capillary feeding. Although only 7/51 (14%) molted nymphs (Figure 2B) were positive by microbial culture for *F. tularensis*




Figure 2. LVS is transmitted transstadially from larvae to nymphs.

A) Open circles are capillary tube fed larvae and filled circles are molted nymphs. The calculated mean CFU/tick for colonized larvae and molted nymphs for each time point is represented by the horizontal line. For each time point up to day 150, n was 15. For day 175, n was 6. B) Percentage of colonized ticks in the same experiment.

from 100 through 175 days post-capillary tube feeding, using RT-PCR, 17/51 (33%) of these nymphs were positive for *F. tularensis* (Table 1; limit of detection was 5 genome equivalent/nymph). The difference between LVS quanta determined by culture *versus* PCR suggests that either LVS may have assumed a viable but not culturable state or that nymphs had cleared the infection but remnant LVS DNA remained in nymphs which was detected by PCR.

Table 1. Comparison of LVS Detection Using Microbial Culture versus RT-PCR.

Days Post-Capillary Feeding	Molted Nymphs		Molted Adults	
	Microbial Culture Positive	RT-PCR Positive	Microbial Culture Positive	RT-PCR Positive
100	4/15	5/15	1/15	4/15
125	1/15	2/15	0/15	0/15
150	1/15	4/15	0/15	2/15
175	1/6	6/6	ND ^a	ND ^a
Total	7/51 (14%)	17/51 (33%)	1/45 (2%)	6/45 (13%)

^aData not collected.

LVS colonization of *I. scapularis* nymphs and transstadial transmission to molted adults

At one-day post-capillary tube feeding, 100% of nymphs were colonized with a mean level of $1.1 \pm 1.2 \times 10^4$ CFU/nymph (Figure 3A). The bacterial count per tick maintained at the same level until day 7 post-capillary tube feeding and then decline until molting to adults at around 42 days post-capillary tube feeding. Although *F. tularensis* was transstadially transmitted to adult ticks, only 1/45 (2%) of the molted ticks were positive for the bacterium after 100 day post-capillary tube feeding by microbial culture (Figure 3B); however, a slightly higher percentage 6/45 (13%) of adults molted from nymphs were positive for *F. tularensis* by RT-PCR (Table 1).

Tissue localization of LVS in *I. scapularis* adults

To determine the dissemination of *F. tularensis* in adult tick tissues following colonization by ingestion, unfed flat adult ticks were capillary tube fed with tick meal and subsequently dissected on various days post-capillary feeding to determine




Figure 3. LVS is transmitted transstadially from nymphs to adults.

A) Open circles are capillary tube fed nymphs and filled circles are molted adults. The calculated mean CFU/tick for colonized nymphs and molted adults for each time point is represented by the horizontal line. For each time point up to day 150, n was 15. B) Percentage of colonized ticks in the same experiment.

nymphs in the south-central U.S. tularemia endemic region is for reptiles and birds, the immature ticks at a lower prevalence can feed on small mammals thought to be the amplifying hosts for maintaining the enzootic tularemia cycle [14,17]. In contrast to the spring-summer questing period and bird-reptile feeding preference for *I. scapularis* larvae and nymphs, adult *I. scapularis* exhibit a fall-winter questing period with feeding preference for large mammal such as canids and deer [14].

Conclusion

Using the capillary feeding method of infecting ticks, we showed that *F. tularensis* strain LVS multiply and maintain parasitism in *I. scapularis*. Although the number of long-term infected ticks were much lower than what was observed with respect to *D. variabilis* and *A. americanum*, a small percentage of ticks still maintained a high quanta of the bacteria sufficient for transmission to susceptible hosts. From our studies, we confirm the potential for *I. scapularis* adults to serve as bridging vector for transmission of sporadic human tularemia during the fall-winter months in the south-central United States tularemia endemic region and *I. scapularis* larvae and nymphs to serve as a vector for small mammals in the summer enzootic tularemia cycle; however, based on the feeding preference of these tick species *I. scapularis* may not play a prominent role in the ecology of tularemia.

References

1. Keim PS, Wagner DM. Humans and evolutionary and ecological forces shaped the phylogeography of recently emerged diseases. *Nat Rev Microbiol*. 2009;7(11):813-21.
2. Eisen LA. Call for renewed research on tick-borne *Francisella tularensis* in the Arkansas-Missouri primary national focus of tularemia in humans. *J Med Entomol*. 2007;44(3):389-97.
3. Mani RJ, Morton RJ, Clinkenbeard KD. Ecology of tularemia in central US endemic region. *Curr Trop Med Rep*. 2016;3:75-79.
4. Kugeler KJ, Mead PS, Janusz AM, et al. Molecular Epidemiology of *Francisella tularensis* in the United States. *Clin Infect Dis*. 2009;48(7):863-70.
5. Staples JE, Kubota KA, Chalcraft LG, et al. Epidemiologic and molecular analysis of human tularemia, United States, 1964-2004. *Emerg Infect Dis*. 2006;12(7):1113-8.
6. Petersen JM, Molins CR. Subpopulations of *Francisella tularensis* ssp. *tularensis* and *holarctica*: identification and associated epidemiology. *Future Microbiol*. 2010;5:649-61.
7. Snowden J, Stovall S. Tularemia: retrospective review of 10 years' experience in Arkansas. *Clin Pediatr (Phila)*. 2011;50(1):64-8.
8. Sjostedt A. Tularemia: history, epidemiology, pathogen physiology, and clinical manifestations. *Ann N Y Acad Sci*. 2007;1105:1-29.
9. Ellis J, Oyston PC, Green M, et al. Tularemia. *Clin Microbiol Rev*. 2002;15(4):631-46.
10. Jellison WL. Tularemia in North America. 1974, Missoula, Montana: University of Montana.
11. Assal NR, Lindeman RD, Carpenter RL. Epidemiologic study on reported human tularemia in Oklahoma, 1944-65. *J Okla State Med Assoc*. 1968;61(3):120-4.
12. Rohrbach BW, Westerman E, Istre GR. Epidemiology and clinical characteristics of tularemia in Oklahoma, 1979 to 1985. *South Med J*. 1991;84(9):1091-6.
13. Hopla CE. *Ixodes scapularis* as a vector of tularemia organisms in the southern United States. Proceedings of the 11th International Congress of Entomology. 1962 Vienna, Austria.
14. Kollars TM, Oliver JH, Kollars PG, et al. Seasonal activity and host associations of *Ixodes scapularis* (Acari: Ixodidae) in southeastern Missouri. *J Med Entomol*. 1999;36(6):720-6.
15. Mani RJ, Reichard MV, Morton RJ, et al. Biology of *Francisella tularensis* subspecies *holarctica* live vaccine strain in the tick vector *Dermacentor variabilis*. *PLoS One*. 2012;7(4):e35441.
16. Mani RJ, Metcalf JA, Clinkenbeard KD. *Amblyomma americanum* as a bridging vector for human infection with *Francisella tularensis*. *PLoS One*. 2015;10(6):e0130513.
17. Hopla CE. The ecology of tularemia. *Adv Vet Sci Comp Med*. 1974;18(0): 25-53.

*Correspondence to:

Rinosh J. Mani
Section Chief (Bacteriology/Mycology), Academic Specialist
Veterinary Diagnostic Laboratory
College of Veterinary Medicine
Michigan State University
Lansing, MI, USA
Tel: (517) 353-9053
E-mail: manirino@msu.edu