

Treatment of endoscopy associated cerebral gas embolism.

Jeffrey Cooper*

Department of Emergency Medicine, University of Nebraska Medical Center, Omaha, NE, USA

Accepted on November 21, 2017

Description

Arterial gas embolisms, albeit rare [1], can have devastating outcomes. Iatrogenic cases have mortality of 16% at hospital discharge and 21% at one year [2]. Embolization of gas during endoscopy is a rare complication with only a few cases having been reported [3] including three of our own [4]. The inciting pathophysiology remains unclear. It is likely that gas is entrained first in the venous system and then shunts to the arterial system via an arteriovenous malformation or cardiac septal defect such as a Patent Foramen Ovale (PFO). PFO with shunting can occur in up to 25% of the general population [5] making screening for PFOs impractical [6].

Endoscopy requires insufflation for better visualization. Such insufflation creates pressure gradients that have been cited as a factor in creating gas embolisms [7-9]. Apart from occlusion, vascular irritation from large boluses of air cause higher neutrophil and cytokine release leading to further inflammation [10,11].

Carbon dioxide (CO₂) is recommended over air for insufflation. CO₂ can be more readily absorbed and should not cause an occlusion in the vasculature. Several studies have been performed showing superiority of CO₂ over air in endoscopic procedures [12,13].

Diagnosis can be difficult because other causes must be ruled out, and the gas bubbles might be absorbed before the necessary test is completed. An echocardiogram can be useful [3,14,15]. However, no test, including computed tomography (CT), has adequate sensitivity to rule out gas emboli as bubbles may be reabsorbed quickly [3]. The best strategy may be prompt referral for hyperbaric oxygen therapy (HBO) based on clinical suspicion alone, given the low risk of HBO and the rapid increase of morbidity and mortality with delaying treatment [16].

As for treatment, initially, the patient should be given 100% O₂ to help dissolve the gas bubble [3]. Nitrous oxide (N₂O), if used, must be stopped immediately due its likelihood of entering the gas embolism. The use of the Trendelenburg position to reduce the likelihood of bubbles going to the brain, has been disproved and harm secondary to increased intracranial pressure and cerebral edema may result [17-20]. Positioning in the left lateral decubitus position may help remove gas from the right ventricle outflow tract [14,21]. Central venous line placement has been shown to be helpful in removing gas bubbles from the right heart [14,21].

The most beneficial therapy in eliminating the gas bubbles is hyperbaric oxygen [7,11]. HBO acts by a variety of mechanisms leading to its use in many adjunctive indications. One of its primary indications is in the treatment of gas embolism [22]. Its beneficial effects in the setting of gas emboli include: Reduced

bubble volume, increased diffusion gradient out of the bubbles, oxygenation of hypoxic tissues, amelioration of cerebral edema, reduced platelet aggregation and activation of the coagulation cascade due to bubble induced endothelial trauma, decreased endothelial binding of leukocytes and prevention of oxygen free radical release [23]. In a retrospective study of 16 patients with gas emboli, 50% of the patients made complete recoveries after HBO [11].

Time to chamber is a crucial factor. Patients who reach the chamber within five hours are twice as likely to show a full recovery as those receiving HBO after five hours [11]. There are few adjunctive therapies for cerebral gas embolism. These include IV fluids to maintain preload, inotropic support to maintain cerebral perfusion pressure and lidocaine [20]. However, rapid initiation of hyperbaric oxygen therapy cannot be overstressed.

Gas embolisms are a rare complication in endoscopic procedures that can have dire outcomes. Due to the inescapability of the factors that may cause them, one must be vigilant when performing an endoscopic procedure for the possibility of gas emboli, especially in someone with advanced pathology, in long procedures requiring interventions and much insufflation, and when using air insufflation instead of carbon dioxide. While there are few adjunctive treatments, the definitive therapy for gas embolisms is prompt hyperbaric oxygen [24-34].

References

1. Mandelstam P, Sugawa C, Silvis S, et al. Complications associated with esophagogastroduodenoscopy and esophageal dilation. *Gastrointestinal Endoscopy*. 1976;23(1):16-9.
2. Bessereau J, Genotelle N, Chabbaut C, et al. A. Long-term outcome of iatrogenic gas embolism. *Intensive Care Med*. 2010;36:1180-87.
3. Donepudi S, Chavalitdhamrong D, Pu PDL. Air embolism complicating gastrointestinal endoscopy: A systematic review. *World J Gastrointest Endosc*. 2013;16(5):359-65.
4. Cooper J, Thomas J, Singh S, et al. Endoscopic bubble trouble: hyperbaric oxygen therapy for cerebral gas embolism during upper endoscopy. *J Clin Gastroenterol*. 2017;51(6):e48-e51.
5. Tanzi A, Onorato E, Casilli F, et al. Is the search for right-to-left shunt still worthwhile? *Acta Neurol Scand*. 2016;133(4):281-8.
6. Laan M, Totte E, Hulst RV, et al. Cerebral gas embolism due to upper gastrointestinal endoscopy. *Eur J Gastroenterol Hepatol*. 2009;21(7):833-5.

7. Green B, Tendler D. Cerebral air embolism during upper endoscopy: Case report and review. *Gastrointestinal Endoscopy*. 2005;61(4):620-3.
 8. Dion Y, Levesque C, Doillon C. Experimental carbon dioxide pulmonary embolization after vena cava laceration under pneumoperitoneum. *Surgical Endoscopy*. 1995;9(10):1065-9.
 9. Lowdon J, Tidmore T. Fatal air embolism after Gastrointestinal Endoscopy. *Anesthesiology*. 1988;69:622-3.
 10. Herron D, Vernon J, Gryska P, et al. Venous gas embolism during endoscopy. *Surgical Endoscopy*. 1999;13:276-9.
 11. Murphy B, Harford F, Cramer F. Cerebral air embolism resulting from invasive medical procedures treatment with hyperbaric oxygen. *Ann Surg*. 1985;201(2):242-5.
 12. Nakamura K, Yamaguchi Y, Hasue T, et al. The usefulness and safety of carbon dioxide insufflation during endoscopic retrograde cholangiopancreatography in elderly patients: A prospective, double-blind, randomized, controlled trial. *Gastrointestinal Endoscopy*. 2014;61(136):2191-5.
 13. Wang W, Wu Z, Sun Q, et al. Meta-analysis: The use of carbon dioxide unsufflation vs. room air insufflation for gastrointestinal endoscopy. *Alimentary Pharmacol Therap*. 2012;35(10):1145-54.
 14. Tan B, Saunier C, Cotton F, et al. Thoracoabdominal CT scan: A useful tool for the diagnosis of air embolism during an endoscopic retrograde cholangiopancreatography. *Annales Francaises d'Anesthesia et de Reanimation*. 2008;27(3):240-3.
 15. Park Y, Kim H, Kim J, et al. Prolonged paradoxical air embolism during intraoperative intestinal endoscopy confirmed by transesophageal echocardiography- A case report. *Korean J Anesthesiol*. 2010;58(6):560-4.
 16. Tomassoni A. Cardiac problems associated with dysbarism. *Cardiol Clinics*. 1995;13(2):266-71.
 17. Rodriguez R, Cornel G, Weerasena N, et al. Effect of Trendelenburg head position during cardiac deairing on cerebral microemboli in children: A randomized controlled trial. *J Thorac Cardiovasc*. 2001;121(1):3-9.
 18. Dexter F, Hindman B, Marshall J. Estimate of the maximum absorption rate of microscopic arterial air emboli after entry into the arterial circulation during cardiac surgery. *Perfusion*. 1996;11(6):445-50.
 19. Tankisi A, Rasmussen M, Juul N, et al. The effects of 10 degrees reverse Trendelenburg position on subdural intracranial pressure and cerebral perfusion pressure in patients subjected to craniotomy for cerebral aneurysm. *J Neurosurg Anesthesiol*. 2006;18(1):11-7.
 20. Erik S, Muth CM. Gas embolism. *N Engl J Med*. 2000;342(7):476-82.
 21. Chavalitdhamrong D, Draganov P. Acute stroke due to air embolism complicating ERCP. *Endoscopy*. 2013;45.
 22. Grim P, Gottlieb L, Boddie A, et al. Hyperbaric Oxygen Therapy. *JAMA*. 1990;263(16):2216-20.
 23. Bessereau J, Genotelle N, Chabbaut C, et al. Long-term outcome of iatrogenic gas embolism. *Intensive Care Med*. 2010;36:1180-87.
 24. Muth C, Kemmer A, Weslau W. Decompression injuries. *MMW- Fortschritte der Medizin*. 2005;147(27):34-5.
 25. ASGE Standards of Practice Committee. Adverse events of upper GI endoscopy. *Gastrointestinal Endoscopy*. 2012;76(4):707-18.
 26. Demaerel P, Gevers A, Bruecker Y, et al. Stroke caused by cerebral air embolism during endoscopy. *Gastrointestinal Endoscopy*. 2003;57(1):134-5.
 27. Sharma V, Nguyen C, Crowell M, et al. A national study of cardiopulmonary unplanned events after GI endoscopy. *Gastrointest Endosc*. 2007;66:27-34.
 28. Mojadidi M, Winoker J, Roberts S, et al. Diagnostic accuracy of transesophageal echocardiogram for the detection of patent foramen ovale: A meta-analysis. *Echocardiography*. 2014;31(6):752-8.
 29. Katzgraber F, Glenewinkel F, Fischler S, et al. Mechanism of fatal air embolism after gastrointestinal endoscopy. *Int J Legal Med*. 1998;111:154-6.
 30. Swaroop V, Desai D, Mohandas K, et al. Dilation of esophageal strictures induced by radiation therapy for cancer of the esophagus. *Gastrointestinal Endoscopy*. 1994;40(3):311-5.
 31. Dellon E, Hawk J, Grimm I, et al. The use of carbon dioxide for insufflation during GI endoscopy: A systematic review. *Gastrointestinal Endoscopy*. 2014;61(136):2191-5.
 32. Rabe C, Balta Z, Wullner U, et al. Biliary metal stents and air embolism: a note of caution. *Endoscopy*. 2006;38(6):648-50.
 33. Tekle W, Adkinson C, Chaudhry S, et al. Factors associated with favorable response to hyperbaric oxygen therapy among patients presenting with iatrogenic cerebral arterial gas embolism. *Neurocrit Care*. 2013;18:228-33.
 34. Fenster B, Curran-Everett D, Freeman A, et al. Saline contrast echocardiography for the detection of patent foramen ovale in hypoxia: A validation study using intracardiac echocardiography. *Echocardiography*. 2014;31(4):420-7.
- *Correspondence to:**
Jeffrey Cooper, MD
Department of Emergency Medicine
University of Nebraska Medical Center
981150 NMC
Omaha, NE 68198
USA
Tel: 4023059515
E-mail: jeffrey.cooper@unmc.edu