

Value of pleth variability index to guide fluid resuscitation in children with sepsis.

Nan Xu^{1,2}, Lixing Qiao^{1*}, Liping Yin¹, Han Li¹

¹Department of Pediatrics, Zhongda Hospital, School of Medicine, Southeast University, Nanjing, PR China

²Department of Pediatrics, Xishan People's Hospital of Wuxi City, Wuxi, PR China

Abstract

Objectives: The present study was conducted to investigate the value of Pleth Variability Index (PVI) to guide fluid resuscitation in children with sepsis.

Materials and methods: 60 patients with sepsis undergoing Early Goal-Directed Therapy (EGDT) were included as subjects of the present study, and randomly divided into two groups by drawing lots. There were 30 cases in each group. The observation group received PVI-guided fluid resuscitation while the control group received transthoracic echocardiography-guided fluid resuscitation. The mortality was recorded in two groups. The oxygen saturation of central vein (ScvO₂), oxygenation index (PaO₂/FiO₂), APACHE II score and clearance rate of lactic acid were compared between two groups. The Intensive Care Unit (ICU) stay and total volume of fluid input were compared between two groups.

Results: Death occurred in 3 cases (10% mortality) in the observation group and in 7 cases in the control group (23.33% mortality). There was no significant difference in mortality between the two groups ($\chi^2=1.920$, $P=0.166$). The remaining patients attained the criteria for resuscitation. PaO₂/FiO₂ after treatment in the observation group was significantly lower than that before treatment and that in the control group ($P<0.05$). The APACHE II score was significantly lower than that before treatment and that in the control group ($P<0.05$). The clearance rate of lactic acid in the observation group was significantly higher than that in the control group ($P<0.05$). The duration of stay in ICU and the volume of fluid input in the observation group were significantly lower than those in the control group ($P<0.05$).

Conclusion: PVI-guided early fluid resuscitation can achieve satisfactory results, reduce complications and help to improve prognosis.

Keywords: Sepsis, Pleth variability index, Transthoracic echocardiography, Fluid resuscitation.

Accepted on October 16, 2017

Introduction

Sepsis is systemic infection caused by pathogenic bacteria that invades the blood, which occurs frequently in children [1]. If not effectively controlled, it may induce meningitis, endocarditis and osteomyelitis [2], and seriously affect the growth and development of children. Early Goal Directed Therapy (EGDT) is an important method for clinical volume management and fluid resuscitation in recent years [3]. It has been used to reduce organ failure and improve survival rate of critically ill patients by individual rehydration [4]. Ultrasound is an important tool to guide fluid resuscitation, which is noninvasive, but affected by human factors [5]. Pleth Variability Index (PVI) allows continuously noninvasive dynamic monitoring of circulating blood volume, and recently has been reported to guide fluid resuscitation [6]. Thus, 60 children with sepsis were included as subjects in the present study to explore the value of PVI, which was reported as below.

Materials and Methods

General information

60 patients with sepsis undergoing EGDT between October 2013 and March 2017 in our hospital were included in the present study, and randomly divided into two groups by drawing lots. There were 30 cases in each group. In the observation group, there were 19 males and 11 females at the age of 6.28 ± 1.48 y old, with heart rate of 108.15 ± 10.32 beat per min (bpm) and systolic blood pressure of 72.32 ± 10.21 mmHg. There were 17 males and 13 females in the control group at the age of 7.03 ± 1.62 years, with heart rate of 110.23 ± 9.87 bpm and SBP of 68.95 ± 9.79 mmHg. There were no significant differences in sex, age, heart rate and blood pressure between the two groups ($P>0.05$).

Inclusion criteria were as follows [7]: (1) The blood samples were positive for pathogenic antigen or DNA detection; (2) The family members had signed the informed consent form.

Exclusion criteria were as follows: (1) patients complicated with malignant tumors; (2) patients complicated with malignant arrhythmia, acute cerebral infarction and cerebral hemorrhage; (3) patients with severe cardiac insufficiency.

Monitoring method

All children in the observation group were administered with mechanical ventilation by tracheal intubation, oxygen uptake and sedatives, and were connected to the Radical-7 type oxygen saturation monitor from United States, continuously monitored for PVI. The control group was measured for Left Ventricular End-Diastolic Volume (LVEDV) by SSI-5000 color Doppler ultrasonography, and was calculated for the dilatation index of Inferior Vena Cava (dIVC) according to the diameter of the inferior vena cava at the end of inspiration and expiration below the xiphoid process.

Liquid resuscitation protocol

The first dose of 20 ml/kg saline was given by bolus injection over 10-20 min, to a total of 40-60ml/kg within 1h, followed by 1/3 solution maintaining at 5-10 ml/kg/h, and the amount was adjusted according to blood pressure. In the event of shock, 5-10 µg/kg/min dopamine was given by continuously intravenous pumping.

Resuscitation standards [8] were as follows: (1) PVI<14%, SBP ≥ 90 mmHg, ScvO₂ ≥ 70% in observation group; (2) LVEDV>60 ml or dIVC<15.5%, SBP ≥ 90 mmHg, ScvO₂ ≥ 70% in control group.

Statistical analysis

The data were analysed by SPSS19.0 statistical software. The measurement data were expressed as mean ± standard. The comparison between the two groups was conducted with the t-

Table 1. Comparison regarding fluid resuscitation between two groups.

Group	Time	ScvO ₂ (%)	PaO ₂ /FiO ₂ (mmHg)	APACHE II (score)
Observation group (n=27)	Before treatment	58.53 ± 3.12	209.36 ± 22.58	22.03 ± 6.52
	After treatment	74.15 ± 3.26 [#]	295.63 ± 27.73 [#]	16.18 ± 5.02 [#]
Control group (n=23)	Before treatment	59.03 ± 2.97	205.21 ± 20.71	22.86 ± 6.39
	After treatment	73.56 ± 3.11 [#]	274.47 ± 25.31 ^{*#}	19.12 ± 5.18 [#]

Note: compared with that before treatment, [#]P<0.05. Compared with that in the observation group, ^{*}P <0.05.

Table 2. Comparison regarding hospitalization between two groups.

Group	Clearance rate of lactic acid (%)	ICU stay (d)	Volume of fluid input (ml)
Observation group (27)	30.24 ± 9.12	10.37 ± 1.64	1938.56 ± 30.07

Table 3. Comparison regarding complications between two groups (n (%)).

test. The enumeration data were expressed as (%). The comparison between the two groups was conducted with the χ^2 test. P<0.05 indicated significant difference.

Results

Comparison regarding fluid resuscitation between two groups

Death occurred in 3 cases (10% mortality) in the observation group and in 7 cases in the control group (23.33% mortality). There was no significant difference in mortality between the two groups ($\chi^2=1.920$, P=0.166). The remaining patients attained the criteria for resuscitation. PaO₂/FiO₂ after treatment in the observation group was significantly lower than that before treatment and that in the control group (P<0.05). The APACHE II score was significantly lower than that before treatment and that in the control group (P<0.05), as provided in Table 1.

Comparison regarding hospitalization between two groups

The clearance rate of lactic acid in the observation group was significantly higher than that in the control group (P<0.05). The duration of stay in Intensive Care Unit (ICU) and the volume of fluid input in the observation group were significantly lower than those in the control group (P<0.05), as provided in Table 2.

Comparison regarding complications between two groups

The overall incidence of complications in the observation group was significantly lower than that in the control group (P<0.05), as provided in Table 3.

Control group (n=23)	19.36 ± 5.67	14.52 ± 2.35	2050.23 ± 48.51
t	4.959	7.324	9.938
P	0	0	0

Group	n	Coagulation disorder	Multiple organ failure	Bleeding	Reduced oxygen diffusion	Overall incidence
Observation group	27	2 (7.41)	0	1 (3.70)	1 (3.70)	4 (14.81)
Control group	23	4 (17.39)	1 (4.35)	3 (13.04)	2 (8.70)	10 (43.48)
χ^2						5.062
P						0.024

Discussion

Liquid resuscitation is an important measure in the acute phase of sepsis, which plays an important role in saving children's life and improving prognosis [9]. EGDT emphasizes individual rehydration to reduce heart load and reduce complications [10]. PVI mainly reflects the interaction between intra-thoracic pressure and venous return [11]. Peripheral Perfusion Index (PI) can reflect the peripheral tissue perfusion [12]. However, the pulsatile blood flow of finger-tip changes cyclically with heart beat [8]. Therefore, PVI can reflect the body liquid state and the reactivity after liquid infusion. This provides the basis for PVI-guided EGDT. In addition, compared with ultrasound, PVI has the advantage of continuous monitoring. Based on the study of Cannesson et al. [13], the present study regarded PVI>14% as the threshold of resuscitation for EGDT in children with sepsis. The results showed that children in the observation group had significantly shorter ICU stay and lower mortality than the control group, suggesting that PVI-guided EGDT was of higher application value.

A study showed that PVI was sensitive to microcirculation. When the microcirculation was disturbed and the blood flow reduced, the visceral vascular smooth muscle was strongly contracted under stimulation by catecholamines, which increased the amount of peripheral blood and venous return [14], and also changed PVI. In the present study, PVI was used as the resuscitation standard. When PVI met the target, fluid therapy was immediately discontinued to achieve satisfactory volume of fluid input, and hence avoid excessive volume due to the inability of continuous ultrasound monitoring. In the present study, the results also showed that the total volume of fluid input in the observation group was less than that in the control group, which was consistent with the above. Lactic acid is a metabolite of anaerobic glycolysis, which shows a significant change at the early stage of low perfusion or hypoxia in tissues, which can better reflect the tissue oxygenation and metabolism and viscera function. Studies have shown that lactate waters and CVP are significantly negatively correlated with complications [15,16]. The present study showed that the observation group had better the clearance rate of lactic acid and APACHE II than the control group after treatment, suggesting that PVI-guided fluid resuscitation could achieve satisfactory outcomes in children with sepsis and help to improve the prognosis. The present study also showed that the ICU stay in the observation group was also lower than that in the control group, which might be due to the excessive retention of the liquid in the tissue space in the control group, causing edema of the peripheral tissue,

damage of the oxygenation function of tissue, and prolongation of the recovery.

Additionally, in the present study, the overall incidence of complications in the observation group was significantly lower than that in the control group. EGIT under the guidance of PVI avoids excessive fluid, reduce the possibility of pulmonary edema and pulmonary interstitial edema, contributes to oxygen dispersion, helps to improve the oxygen supply to tissues, and prevent the destruction of body compensatory mechanism and the inner environment [17]. Furthermore, the ideal volume state also reduces the possibility of dilution-induced hypoprothrombinemia.

Conclusion

In summary, PVI-guided early fluid resuscitation can achieve satisfactory results, and reduce complications and mortality in children with sepsis, which is of high application value.

References

1. Stewart CE, Radia T, Ghafoor K. Paediatric sepsis, the under-recognised killer: quality improvement initiative of outreach teaching in paediatric sepsis. Arch Dis Child Educ Pract Ed 2017.
2. Huang M, Li KC, Xu X. Retrospective analysis of distribution and antibiotic resistance of pathogenic bacteria in children septicemia over 10 years. Chin J Biochem Pharm 2016; 165-168.
3. Chen X, Zhu W, Tan J, Nie H, Liu L, Yan D, Zhou X, Sun X. Early outcome of early-goal directed therapy for patients with sepsis or septic shock: a systematic review and meta-analysis of randomized controlled trials. Oncotarget 2017; 8: 27510-27519.
4. Kalil AC, Johnson DW, Lisco SJ, Sun J. Early goal-directed therapy for sepsis: a novel solution for discordant survival outcomes in clinical trials. Crit Care Med 2017; 45: 607-614.
5. Tao YK, Zhang GQ, Wang HF, Liu XL, Qi ZW, Yan ST, Lu HT, Gu CD. Ultrasonographic measurement of inferior vena cava internal diameter and respiratory variation index as guidelines in fluid resuscitation of septic shock. Chin J Emerg Med 2013; 22: 1252-1255.
6. Wang YZ, Chen R, Wang DH. Predictive value of pleth variability index on response of fluid resuscitation for patients with hemorrhagic shock. Med Pharm J Chin People Liber Army 2016; 28: 73-79.

7. Lv WX, Ye MK, Liang DW. Value of serum calcitonin in the clinical diagnosis of septicemia. *J Guiyang Med Coll* 2014; 39: 739-740.
8. Liu FF, Tian M, Wu ZF, Jin L, Li WY. Application of Plethysmographic Variability Index for Monitoring Perioperative Fluid Therapy. *Prog in Mod Biomed* 2014; 14: 1194-1196.
9. Li C, Yun D. Improvement effect of early goal-directed therapy on the prognosis in patients with septic shock. *Chin Crit Care Med* 2015; 27: 899-905.
10. Evetts GE, Birch K. Early-goal-directed therapy and protocolised treatment in septic shock. *J Intensive Care Soc* 2015; 16: 164-168.
11. Lu N, Zheng R, Lin H, Yu J, Shao J, Wu X, Wang H. A prospective clinical study of pleth variability index in prediction of volume responsiveness in patients with septic shock. *Chin Crit Care Med* 2015; 27: 17-21.
12. Manevska N, Stojanoski S, Pop Gjorceva D, Todorovska L, Miladinova D, Zafirova B. Normal values of tissue-muscle perfusion indexes of lower limbs obtained with a scintigraphic method. *Pril (Makedon Akad Nauk Umet Odd Med Nauki)* 2017; 38: 55-61.
13. Cannesson M, Delannoy B, Morand A, Rosamel P, Attof Y, Bastien O, Lehot JJ. Does the Pleth variability index indicate the respiratory-induced variation in the plethysmogram and arterial pressure waveforms? *Anesth Analg* 2008; 106: 1189-1194.
14. Wang JL, Luo ZB, Yang JZ. Correlation of pleth variability index with central venous pressure. *J Reg Anat Oper Surg* 2015; 24: 643-646.
15. Li SH, Liu F, Zhang YT, Zhao WH. Initial serum lactate level as predictor of morbidity after major abdominal surgery. *Nation Med J Chi* 2008; 88: 2470-2473.
16. Lin ZP, Zhou Y, Chen WH. Effect of voluven and lactate ringers solution on hemodynamics and acid-base balance in patients with general anesthesia. *Chin J Bioch Pharm* 2014; 142-144.
17. Lei M, Bao Q, Xu LY, Zhong TD. Efficacy of pleth variability index in guiding volume therapy in patients undergoing thoracoscopic surgery. *Chin J Anesthesiol* 2015; 35: 987-989.

***Correspondence to**

Lixing Qiao
Department of Pediatrics
Zhongda Hospital
School of Medicine
Southeast University
PR China