

## Vestibular influences on respiratory endurance.

Teenu mol M J<sup>1</sup>, Kumar Sai Sailesh<sup>2\*</sup>, Archana R<sup>3</sup>, Srilatha Bhasetti<sup>4</sup>, Mukkadan J K<sup>5</sup>

<sup>1</sup>Department of Physiology, Little Flower Institute of Medical Sciences and Research, Angamaly, Kerala, India

<sup>2</sup>Department of Physiology, DMWayanad Institute of Medical Sciences, Naseera Nagar, Meppadi, Wayanad, Kerala, India

<sup>3</sup>Department of Physiology, Saveetha Medical College, Saveetha University, Thandalam, Chennai, Tamil Nadu, India

<sup>4</sup>Department of Biochemistry, Apollo Institute of Medical Sciences and Research, Hyderabad, Telangana, India

<sup>5</sup>Little Flower Medical Research Centre, Angamaly, Kerala, India

Accepted September 29, 2017

### Editorial

The vestibular system, which is comprised of tiny receptors in the inner ear plays a major role in maintenance of posture and equilibrium. Apart from its regular function, vestibular system influences all the body systems and contributes to homeostasis. Hence stimulation of vestibular system, which begins in the intrauterine life, must be continued throughout the life. Vestibular regulation of respiratory functions is mediated by the reflexes that from medial and inferior vestibular nucleus. Vestibular projections are identified throughout the brain and in fact certain brain structures like hippocampus undergoes atrophy followed by vestibular lesions. Postural changes activate of pulmonary receptors and stretch receptors in the respiratory muscles which will affect the respiratory muscle [1,2]. Numerous studies have shown that vestibular nerve stimulation produces a reflex response in phrenic nerve, intercostal nerve and abdominal muscle nerve [3-5]. Vestibular system maintaining posture through its action on limb muscle and these influences also extend to the respiratory postural muscles [6]. The stable venous return to the heart during movements and changes in the posture may provide by the vestibulo-respiratory reflexes in concert with vestibulo-sympathetic reflexes. The augmented contraction of the diaphragm increases negative intrathoracic pressure this would result in more blood being pulled in the heart. The blood from abdomen into the thorax is forced by the increased activities of the abdominal muscles which also enhances venous return [7,8].

The regions of the vestibular nuclei which participates in vestibulo-sympathetic reflexes and the same is the mediator for vestibular influences on respiration. the brainstem neurons have the responsibility for central respiratory drive. the vestibulo-respiratory reflexes have considered the same role of brainstem neurons here. The dorsal respiratory and the ventral root ganglion are the two groups of respiratory neurons in the medulla. The dorsal group ganglion is located in the ventrolateral portion of the nucleus of the solitary tract. It contains predominately inspiratory neurons, including bulbo-spinal neurons that convey descending respiratory drive to motor neurons supplying the diaphragm and inspiratory intercostal muscles. The ventral respiratory group (VRG) is located in the ventrolateral medulla and extends from the nucleus retro-ambiguus at the border with the first cervical

spinal segment (caudal end) to the nucleus ambiguus and retro-facial nucleus (rostral end). The VRG contains heterogeneous cell types, including bulbo-spinal inspiratory and expiratory neurons. Medullary interneurons, and laryngeal and pharyngeal motor neurons. A third group of respiratory neurons is located in the parabrachial region of the pons. Shaping the property of respiratory activity might also the responsibilities of other vestibular connection. Recent neuroanatomical studies have demonstrated a projection from the superior vestibular nucleus to the vicinity of the pontine respiratory group [9]; however, vestibular inputs to pontine respiratory neurons are yet to be evidenced neuro-physiologically. Neurons in the raphe nuclei also receive vestibular inputs [9]. Role of vestibular stimulation has been extensively studied in recent years which is evident by published articles on vestibular stimulation and registered clinical trials. This is the need of time to understand the essentiality of stimulation of the vestibular system in our everyday life style to improve the quality of life.

### Reference

1. Miller AD, Yates BI. Vestibular effects on respiratory activity. In: Yates BI, Miller AD, editors. Vestibular autonomic regulation. Boca Raton (FL): CRC Press; 1996. pp: 13-25.
2. Yates BI, Miller AD. Vestibular respiratory regulation. In: Miller AD, Bianchi AL, Bishop BP, editors. Neural control of the respiratory muscles. Boca Raton (FL): CRC Press; 1996. pp: 271-283.
3. Siniaia MS, Miller AD. Vestibular effects on upper airway musculature. Brain Res 1996;736:160-164.
4. Yates BJ, Jakus J, Miller AD. Vestibular effects on respiratory outflow in the decerebrate cat. Brain Res. 1993;629:209-17.
5. Danon J, Druz WS, Goldberg NB, Sharp JT. Function of the isolated paced diaphragm and the cervical accessory muscles in CI quadriplegics. Am Rev Respir Dis 1979;119:909-19.
6. Lloyd TC. Effect of inspiration on inferior vena caval blood flow in dogs. J Appl Physiol 1983;55:1701-8.
7. Youmans WB, Tjioe DT, Tong EY. Control of involuntary activity of abdominal muscles. Am J Physical Med 1974;53:57-74.

8. Yates BJ, Goto T, Bolton PS. Responses of neurons in the caudal medullary raphe nuclei of the cat to stimulation of the vestibular nerve. *Exp Brain Res* 1992;89:323-32.
9. Yates BI, Goto T, Kennan I, Bolton PS. Responses of caudal medullary raphe neurons to natural vestibular stimulation. *Neurophysiol* 1993;70:938-46.

**\*Correspondence to:**

Kumar SS  
Assistant Professor  
Department of Physiology  
DMWayanad Institute of Medical Sciences  
Naseera Nagar, Meppadi, Wayanad, Kerala, India  
E-mail: [saisailesh.kumar@gmail.com](mailto:saisailesh.kumar@gmail.com)