

RESHAPE FACE FOR PATIENT WITH SILICONE INJECTION CASE

Beatrix Lee

Accreditation University Indonesia, Indonesia

Reshape Face for patient with Silicone Injection Case with the right type of Filler. Patients who meet me they has performed the Surgery to remove silicone on their faces and they are so helped with the procedure and they very satisfied with the results of removal of Silicone on their face . They come to me complaining about how their facial shape can be reproportioned after performed surgical removal of silicon. Because as we know silicon is very dangerous, the content of cytokines TNF-a (alpha), IFN-g (gamma), IL-10, TREG (CD4 + CD25 +) and indoleamine 2,3-dioxygenase (IDO) that has been banned by the FDA since 1992, can cause granuloma and migration of the injected facial area causing severe tissue damage. I already help more than 50 patients who has become Victims of Silicone Injection , for patients who want to do a filler procedure ,I advise patients for facial reshaping should have been six months recovery from surgery, because before six months the patient's face is still not revised from inflammation. After the downtime is over I will analyze the damage on the patient's face, I will determine the type of filler used based on highly advanced hyaluronic acid gel, because the right type of filler will be very decisive. My first Consideration and my diagnosis for patient is because Silicone causes Tissue Damage and after performed removal patient will have lost they Tissue Area Volume. This condition made me decide to use the Biphasic Filler.