allied Joint Event on

19th International Conference on

OCULAR PHARMACOLOGY AND EYE CARE

&

World Congress on

PUBLIC HEALTH, EPIDEMIOLOGY AND NUTRITION

September 03-04, 2018 | Lisbon, Portugal

Amila Sampath Chandrasekera, Arch Gen Intern Med 2018, Volume 2 | DOI: 10.4066/2591-7951-C4-011

CO-RELATION BETWEEN THE PERIPAPILLARY MICROVASCULATURE AND OCULAR PULSE AMPLITUDE IN GLAUCOMATOUS OPTIC NEUROPATHY

Amila Sampath Chandrasekera

Vision Care Optical Services PVT Ltd, Sri Lanka

Introduction: Although elevated intraocular pressure (IOP) is the main risk factor for glaucoma, other risk factors and vascular risk factors have been implicated in the pathogenesis of glaucoma. Impaired microcirculation in the optic nerve head may contribute to the initiation and progression of glaucomatous neuropathy. It has been proposed that the main pathologic changes in glaucoma are in the deep vascular areas in optic nerve head region.

Objective: To find out the co-relation between ONH microvasculature perfusion and ocular pulse amplitude and their effect on retinal nerve fiber layer thickness in glaucomatous optic neuropathy.

Procedure: Study was conducted with 1000 subjects who were diagnosed as glaucomatous optic neuropathy (based on the Hodapp-Parrish-Anderson criteria) and grouped as normal tension glaucoma, primary open angle glaucoma and non-glaucomatous. In every subject peripapilary vascular perfusion (PVP), ocular pulse amplitude (OPA) and retinal nerve fiber layer thickness (RNFLT) was measured.

Results: In every group, glaucomatous stages and glaucoma suspects showed significantly lower blood perfusion index compared with normal eyes (P \leq 0.0015). Blood perfusion showed a direct correlation with ocular perfusion pressure calculated with OPA (P \leq 0.0123). Similar discrimination capability PVP compared with RNFL thickness was found in both disease groups.

Conclusion: Impaired blood supply to the optic nerve head peripapillary area, may cause to develop RNFL thinning which directly leads to glaucomatous optic neuropathy. Investigations on PVP and OPA will provide a very early diagnosis and a repeatable follow up baseline for the disease, beyond the existing methods.



BIOGRAPHY

Amila Sampath Chandrasekera has completed the certificate of ophthalmic assistance in 2011 and completed diploma in Optometry in 2015 from Academy of Vision Care Optical Services Sri Lanka. He has presented oral and poster presentations in academic and international level poster and won the second place for best poster award in International Conference in Clinical and Experimental Ophthalmology 2015. Currently, he is practicing at the retinal and glaucoma diagnostic unit at Vision Care Optical Services PVT Ltd., head office Sri Lanka.

emmysampath2@gmail.com