




ISSN Print/Online: 2320-9577/2320-9585  
INTERNATIONAL JOURNAL OF PURE AND APPLIED ZOOLOGY  
Volume 1, Issue 2, June 2013  
Available online at: <http://www.ijpaz.com>  
RISHAN PUBLICATIONS



RESEARCH ARTICLE

OPEN ACCESS

## A STUDY ON THE TOXICITY OF CHROMIUM ON THE HISTOLOGY OF GILL AND LIVER OF FRESHWATER FISH *LABEO ROHITA*

<sup>1</sup>K. MUTHUKUMARAVEL\* AND <sup>2</sup>P. RAJARAMAN

<sup>1</sup>PG And Research Department of Zoology, Khadir Mohideen College,  
Adirampattinam – 614 701. Tamilnadu, India

<sup>2</sup>Thiru Vi Ka Govt. Arts College, Tiruvarur-610 003, Tamilnadu, India

\*Corresponding Author Email: [kumar\\_phd\\_2003@yahoo.co.in](mailto:kumar_phd_2003@yahoo.co.in), Tel: +91 9791387363

Article History: Received: 06.06.2013, Accepted: 19.06.2013, Published: 26.06.2013

### ABSTRACT

The toxic effect of chromium on the histology of gill and liver of *Labeo rohita* fingerlings was studied. The fingerlings were exposed for 10, 20 and 30 days in 10% sublethal concentration of 96 h LC<sub>50</sub> of chromium (3.5 ppm). The gills exposed to sublethal concentration of chromium showed mild histological alterations during 10 days of exposure. However after 30 days, fusion of gill lamellae, hypertrophy and degeneration of epithelium were prominent. Liver lesions consisted of vacuolation, degeneration of hepatocytes and disintegration of cell boundaries of hepatocytes. These changes occurred predominantly in the 30 days exposure.

**Keywords:** Heavy metal, chromium, fish, histology, gills and liver.

### INTRODUCTION

Fish is an excellent source of protein in human diet. The unique feature which differentiates fish food other animal protein sources is the presence of omega-3 fatty acids such as Linolenic acid, Decosa hexaenoic acid (DHA) and Eicosa pentaenoic acid (EPA). DHA promotes learning ability in children and improved memory in adults. DHA is also essential for the foetal growth and development. Omega fatty acid is also good for heart and helps to control diabetics by improving insulin action (Lee and Reasner, 2000 and Lemos *et al.*, 2005). That's why it must be included in human diet at least 1.3 kg per week (FAO, 1989). However, the fish habitats are being contaminated alarmingly through a number of aquatic pollutants. Among these pollutants heavy metals are most injurious to fish. These pollutants have not only depleted

the fish stock but also have threatened the human health by incorporating into food chain (Pip, 1995). Heavy metal chromium is widely used in metallurgic, refractory, chemical and tannery industries. Tanneries are the major industries that use chromium for the treatment of leather and nearly 40% of the chromium used is released into the environment. In the present work, an attempt was made to evaluate the long term exposure effect of chromium on the histology of gill and liver of the freshwater fish *Labeo rohita*.

### MATERIALS AND METHOD

The fish, *L. rohita* fingerlings (Wt: 8 0.5g; Length 7.1 cm) were collected from the Katherasan Aqua Farm near Thanjavur, Tamil Nadu. They were acclimatized for 15 days in large cement tanks (Temperature 24.2°C, pH 7.2 ±0.3 and Dissolved Oxygen 7.9±0.7 mg/l)

previously washed with 1% potassium permanganate. The water was renewed every 24 h. The LC<sub>50</sub> of potassium dichromate for 96h was found out by using Probit method (Finney, 1971). Stock solution of chromium was prepared and the toxicity tests were conducted to be 3.5 ppm. For histological studies *L. rohita* were reared in sublethal concentration (10% of 96 hours LC<sub>50</sub>) for a period of 10, 20 and 30 days. The gill arches and liver were dissected out and fixed in Bouin's fluid. The tissues were embedded in paraffin (58°C) and sectioned at 6µ thickness. The sections were stained with haematoxylin and eosin (Gurr, 1959). The stained slides were examined for histopathological changes and were photomicrographed.

## RESULTS AND DISCUSSION

In control fish, the secondary gill lamellae (SGL) appeared as finger-like structures. The SGL was thin, slender and attached on either side of the primary gill lamellae (PGL). The secondary gill lamellae are highly vascularised and surrounded by a thin layer of epithelial cells (Figure 1).


The overall observed results in the present investigation indicates that marked histopathological changes have been found in the gill and liver of fish *L. rohita* under sublethal concentrations of chromium in chronic exposure. Fusion and shortening lamellae, hypertrophy, degeneration of epithelium and necrosis were found in the gills of chromium treated *L. rohita* (Figures 2 & 3). Higher degree of hypertrophy and fusion of gill lamellae were prominent in the gills of fish exposed to 30 days. Hemalatha and Banerjee (1977) and Gupta and Kumar (2006) noted similar types of gill lesions in zinc treated *Heteropneustes fossilis* and mercury treated *Cirrhinus mrigala* respectively. Kaoud and El-Dahshan (2010) observed severe hyperplasia in secondary gill lamellae which lead to complete embedding in adjacent lamellae in copper, cadmium, lead and mercury treated *Oreochromis niloticus*. In the present study, hypertrophy and degeneration of secondary lamellae were apparent in *L. rohita* exposed to chromium (Figure 4). These observations are quite comparable to pathological lesions induced in gills by mercuric chloride in *Acipenser persicus* fry (Khoshnood *et al.*, 2011), by lead and cadmium treatment in *Cyprinus carpio* (Patnaik *et al.*, 2011), *Lates calcarifer* (Thophon *et al.*,

2003), *Brachydanio rerio* and *Salmo gairdneri* (Karlson-Norgren *et al.*, 1985). Patel and Bahadur (2010) also noted severe gill lesions in copper treated *Catla catla*. In the present investigation the gill epithelium of chromium treated fish was completely desquamated, fusion and shapeless secondary lamellae and were broken at several places (Figure 4). Daoust *et al.* (1984) also observed similar pathological lesions in the gill of copper treated rainbow trout, *Salmo gairdneri*. Further, Hemalatha and Banerjee (1997) and Al-Attar (2007) also observed such gill damages in zinc chloride and nickel treated *Heteropneustes fossilis* and *Oreochromis niloticus*.

Liver of fish is responsible for digestion, filtration and storage of glycogen. The liver also produces many enzymes that stored in the gall bladder. The liver functions to store food energy. The normal liver is made up of continuous mass of hepatocytes with large number of blood sinusoids (Figure 5). The fish exposed to the sublethal concentrations of chromium showed the vacuolation, loose arrangement of hepatic cells, histolysis and disintegration of cell boundaries (Figures 6-8). The damage was more severe and progressive after 30 days exposure. Histological changes in the liver of fishes have been extensively reported. Athikesavan *et al.* (2006) reported the histological lesions in the liver of *Hypophthalmichthys molitrix* exposed to nickel. They observed marked changes like degeneration of blood vessels hypertrophy, vacuolisation necrosis and pyknotic nuclei of the exposed fish. The histological lesions due to cadmium and zinc poisoning were reported by Van Dyk *et al.* (2007) in *Oreochromis mossambicus*. Loganathan *et al.* (2006) and Radhakrishnan and Hemalatha (2010) have also observed the histological changes in the liver of zinc treated *L. rohita* and cadmium chloride treated *Channa striatus*. They observed that the cytoplasmic vacuolization of hepatocytes, congestion of blood vessel, leucocytic infiltration and necrosis. The results of the present observations in *L. rohita* exposed to chromium were in agreement with those of the earlier workers especially in the vacuolization and necrosis in hepatic tissue. Intracellular vacuolation, necrosis and shrinkage of nuclei were also apparent in the present study in chromium treated *L. rohita*. Similar changes are observed by Loganathan *et al.* (2006) in zinc treated *L. rohita*. The development of necrosis,


congestion of hepatic blood vessels and vacuolization in chromium treated *L. rohita* were mainly due to large scale accumulation of these metals in liver. Liver is the vital organ for

detoxification of unwanted and toxic substances. Accumulation and elimination processes of metals ions in the liver may lead to hepatic lesions.



**Figure 1.** Control fish gill.

PGL - Primary Gill Lamellae  
SGL - Secondary Gill Lamellae




**Figure 2.** 10 % SLC of Chromium treated fish gill 10 days.

ESL - Erosion of Secondary Lamellae  
FSL - Fusion of Secondary Lamellae




**Figure 3.** 10 % SLC of Chromium treated fish gill after 20 days.

DE - Degeneration of Epithelium, H - Hypertrophy  
FSL - Fusion of Secondary Lamellae




**Figure 4.** 10 % SLC of Chromium treated fish gill after 30 days.

DE - Degeneration of Epithelium, HT - Hypertrophy,  
FSL - Fusion of Secondary Lamellae



**Figure 5.** Control fish liver.

NH - Normal hepatocytes,  
BS - Blood Sinus.



**Figure 6.** 10 % SLC of Chromium treated fish Liver after 10 days.

DH - Degeneration of hepatocytes,  
V - Vacuolization.






Fig.7: 10 % SLC of Chromium treated fish Liver after 20 days.

V – Vacuolization

DCB - Disintegration of Cell Boundaries




Fig.8: 10% SLC of Chromium treated fish Liver after 30 days.

DH - Degeneration of Hepatocytes

V - Vacuolization

## Conclusion

The results in the present study showed that the exposure of rohu to chromium caused pathology in their organs such as gill and liver and they were associated with the exposure period. Histological alterations in rohu under the toxicity of heavy metal chromium can be used as a sensitive model to monitor the aquatic pollution.

## CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest associated with this article.

## ACKNOWLEDGEMENT

The authors wish to acknowledge the Principal and HOD of Zoology, Khadir Mohideen College, Adirampattinam for making available all the necessary facilities.

## REFERENCES

- Al-Attar, A.M. 2007. The influences of nickel exposure on selected physiological parameters and gill structure in the teleost fish, *Oreochromis niloticus*. *J. Biol. Sci.*, **7** (1): 77-85.
- Athikesavan, S., Vincent, S., Ambrose, T. and Velmurugan, B. 2006. Nickel induced histopathological changes in the different tissues of freshwater fish, *Hypophthalmichthys molitrix* (Valenciennes). *J. Environ. Biol.*, **27**: 391-395.
- Daoust, P.Y., Wobeser, G. and Newstead, J.D. 1984. Acute pathological effects of inorganic mercury and copper in gills of rainbow trout. *Vet. Pathol.*, **21**(1): 93-101.
- FAO (Food and Agriculture Organization), 1989. Evaluation of certain food additives and contaminants. 33<sup>rd</sup> report of joint FAO/WHO Expert Committee on food additives. Technical report series, p. 776.
- Finney, D.J. 1971. Probit Analysis, 3<sup>rd</sup> Ed. Cambridge University Press, Cambridge, p.333.
- Gupta, A.K. and Kumar, A. 2006. Histopathological lesions in the tissues of *Cirrhinus mrigala* (Ham.) fingerlings exposed to a sublethal concentration of mercury. *J. Environ. Biol.*, **27**(2): 235-239.
- Gurr, E. 1958. Methods of analytical histology and histochemistry. Leonard Hill Books Ltd, London.
- Hemalatha, S. and Banerjee, T.K. 1997. Histopathological analysis of sublethal toxicity of zinc chloride to the respiratory organs of the air breathing catfish *Heteropneustes fossilis* (Bloch). *Biol. Res.*, **30**: b11-21.
- Kaoud, H.A. and El-Dahshan, A.R. 2010. Bioaccumulation and histopathological alterations of the heavy metals in *Oreochromis niloticus* fish. *Nature Sci.*, **8**(4): 147-154.
- Karlsson-norrgren, L., Runn, P., Haux, C., Forlin, L. 1985. Cadmium induced changes in gill morphology of zebra fish *Brachydanio rerio* (Hamilton-Buchanan) and rainbow trout *Salmo gairdneri* Richardson. *J. Fish Biol.*, **27**: 81-95.

- Khoshnood Z., Khodabandeh S., Shahryari Moghaddam M., Mosafer Khorjestan S. 2011. Histopathological and pathomorphological effects of mercuric chloride on the gills of *Persian sturgeon*, *Acipenser persicus*, fry. *Int. J. Nat. Resour. Mar. Sci.*, **1**(1): 23-32.
- Lee, N A and Reasner C A. 2000. Beneficial effect of chromium supplementation on serum triglycerides levels in *Cyprinus carpio*. *Diab. Care.*, **17**: 1449-1452.
- Lemos, N G, Dias A L, Silva S T and Mantovani M S. 2005. Evaluation of environmental water using the comet assay in *Tilapia rendalli*. *Environ. Toxicol. Pharmacol.*, **9**: 197-201.
- Loganathan K, Velmurugan B, Howrelia J H, Selvanayagam M and Patnaik B B. 2006. Zinc induced histological changes in brain and liver of *Labeo rohita* (Ham). *J Environ Biol.*, **27**(1): 107-110.
- Patel, J.M. and Bahadur, A. 2010. Histopathological alterations in *Catla catla* induced by chronic exposure of copper ions. *J. Cell. Tissue Res.*, **10**(3): 2365-2370.
- Patnaik, B.B., Howrela, H.J., Mathews, T. and Selvanayagam, M. 2011. Histopathology of gill, liver, muscle and brain of *Cyprinus carpio communis* L exposed to sublethal concentration of lead and cadmium. *African. J. Biotech.*, **10**(57): 12218-12223.
- Pip, E. 1995. Cadmium, lead and copper in freshwater mussel from Assiniboine river, Manitoba, Canada. *J. Moll. Stud.*, **61**: 295-302.
- Radhakrishnan, M.V and Hemalatha S. 2010. Sublethal toxicity effects of cadmium chloride to liver of freshwater fish *Channa striatus* (Bloch) Am-Emuras. *J. Toxicol. Sci.*, **2**(1): 54-56.
- Thophon S, Kruatrache M, Upatham E.S, Pokethitiyook P, Sahaphong S. and Jaritkhuan S. 2003. Histopathological alterations of white seabass *Lates calcarifer* in acute and sub chronic cadmium exposure. *Environ. Poll.*, **121**: 307-320.
- Van Dyk. J., Pieterse, G. and Van Vuren, J. 2007. Histological changes in the liver of *Oreochromis mossambicus* (Cichlidae) after exposure to cadmium and zinc. *Ecotoxicol. Environ. Saf.*, **66**: 432-440.

---

**Cite this article as:**

Muthukumaravel, K. and Rajaraman, P. 2013. A study on the toxicity of chromium on the histology of gill and liver of freshwater fish *Labeo rohita*. *J. Pure Appl. Zool.*, 1(2): 122-126.

---