

## An audit of sequential coronary registered tomography angiography (CTA) to evaluate plaque movement and restorative effect of anti-atherosclerotic drugs.


Isabella Gomez\*

Allied Academies, 40 Bloomsbury Way, Lower Ground Floor, London, United Kingdom

Accepted February 18, 2021

Figure 1. Qualitative and quantitative analysis of coronary CTA. Panels (a-c) Standard qualitative analysis of coronary CTA data for the presence of obstructive CAD (stenosis $\geq$ 50%): a 3D volume-rendered image of the coronary tree with potential narrowing in the proximal (arrow) and mid LAD (arrowhead). b multiplanar reformatted long and c short axis images of the proximal LAD area (white arrows) demonstrating luminal narrowing $<$ 50%. This patient would be classified as having non-obstructive CAD unlikely to be hemodynamically significant as a result of routine diagnostic assessment. Panels d, e Advanced

quantitative analysis of another patient demonstrating a plaque with high-risk features. (d)-Cross section of a vessel depicting central low attenuation core in red (asterisk) with ring-like peripheral high attenuation in light green (arrow)-the napkin ring sign. (e)-Output of quantitative analysis with derived minimal luminal area, degree of stenosis, remodeling index, plaque burden, noncalcified and low HU volume, and mean plaque attenuation.



\*Correspondence to:

Isabella Gomez

E-mail: [cardiothoracic@escienceopen.com](mailto:cardiothoracic@escienceopen.com)