

Antibiotic stewardship in primary care: Challenges and best practices for reducing resistance.

Prasad kollenu*

Department of Medicine, Northwestern University, Galter E. Huron Street, Chicago

Introduction

Antibiotic stewardship refers to the careful and responsible use of antibiotics to minimize the emergence of resistance, preserve their efficacy, and optimize patient outcomes. In primary care settings, where the majority of antibiotics are prescribed, stewardship is particularly critical. This essay explores the challenges faced in antibiotic stewardship in primary care and outlines best practices for reducing antibiotic resistance. One of the primary challenges in antibiotic stewardship in primary care is the high volume of antibiotic prescriptions [1].

According to the Centres for Disease Control and Prevention approximately 30% of antibiotics prescribed in outpatient settings are unnecessary. Over prescription of antibiotics, particularly for viral infections such as the common cold and influenza, contributes significantly to resistance development. Another challenge is the lack of rapid diagnostic tools in primary care settings. Many infections are diagnosed clinically without laboratory confirmation, which can lead to inappropriate antibiotic use [2].

The reliance on broad-spectrum antibiotics, due to the inability to quickly identify pathogens, exacerbates resistance. Additionally, primary care providers often face pressure from patients who expect antibiotics for their symptoms, further complicating the stewardship effort [3].

Adherence to evidence-based guidelines is crucial for effective antibiotic stewardship. Guidelines such as those from the Infectious Diseases Society of America provide clear recommendations on the appropriate use of antibiotics for various infections. Primary care providers should familiarize themselves with these guidelines to ensure they prescribe antibiotics only when necessary and select the appropriate drug, dose, and duration [4].

Improving diagnostic accuracy is essential for reducing unnecessary antibiotic use. Implementing rapid diagnostic tests, such as point-of-care tests for strep throat or influenza, can help differentiate between bacterial and viral infections. While these tests are not always available, increasing their accessibility and encouraging their use can reduce the incidence of inappropriate prescribing [5].

Educating patients about the risks of antibiotic overuse and the limitations of antibiotics for viral infections is vital.

Primary care providers should engage in clear communication with patients about why antibiotics are not always necessary and the potential consequences of misuse. Providing alternative management strategies for viral infections, such as symptomatic relief and watchful waiting, can help align patient expectations with clinical practice [6].

A delayed prescribing strategy can be an effective way to manage infections where antibiotics may be beneficial but are not immediately necessary. This approach involves prescribing antibiotics but advising patients to wait and see if symptoms improve before filling the prescription. Studies have shown that delayed prescribing can reduce the overall use of antibiotics without compromising patient outcomes [7].

Regular monitoring and feedback on antibiotic prescribing patterns can help identify areas for improvement and reinforce adherence to stewardship practices. Implementing audit and feedback mechanisms, where providers receive reports on their prescribing behaviours and outcomes, can lead to more judicious antibiotic use [8].

Ongoing education and training for primary care providers are essential for maintaining effective stewardship practices. Continuing medical education programs focused on antibiotic stewardship can update providers on the latest guidelines, resistance patterns, and best practices [9].

Additionally, fostering a culture of stewardship within primary care practices can enhance commitment to appropriate antibiotic use [10].

Conclusion

Antibiotic stewardship in primary care is critical for combating the growing problem of antibiotic resistance. Addressing challenges such as over prescription, limited diagnostic tools, and patient expectations through evidence-based guidelines, improved diagnostics, patient education, and ongoing provider training can significantly reduce resistance and preserve the effectiveness of antibiotics. By adopting these best practices, primary care providers can play a pivotal role in ensuring the responsible use of antibiotics and safeguarding public health.

References

1. Chua KP, Fischer MA, Rahman M, et al. Changes in the Appropriateness of US Outpatient Antibiotic Prescribing After the COVID-19 Outbreak: An Interrupted Time

*Correspondence to: Prasad kollenu, Department of Medicine, Northwestern University, Galter E. Huron Street, Chicago, Email: prasadkollenu@hmtl.org

Received: 02-Oct-2024, Manuscript No. AAAGIM-24-148089; Editor assigned: 04-Oct-2024, PreQC No. AAAGIM-24-148089(PQ); Reviewed: 14-Oct-2024, QC No. AAAGIM-24-148089; Revised: 19-Oct-2024, Manuscript No. AAAGIM-24-148089 (R); Published: 26-Oct-2024, DOI:10.35841/aaagim-8.5.257

- Series Analysis of 2016–2021 Data. Clin Infect Dis. 2024;79(2):312-20.
2. Hoare A, Marsh PD, Diaz PI. Ecological therapeutic opportunities for oral diseases. Microbiol Spectr. 2017;5(4):10-128.
 3. Barnett ML, Linder JA. Antibiotic prescribing to adults with sore throat in the United States, 1997-2010. JAMA Intern Med. 2014;174(1):138-40.
 4. Ventola CL. The antibiotic resistance crisis: part 1: causes and threats. J Clin Pharm Ther. 2015;40(4):277.
 5. Simon MS. The Federal Future of Medication Abortion. Ind. L. Rev. 2023;57:613.
 6. Buehrle DJ, Shively NR, Wagener MM, et al. Sustained reductions in overall and unnecessary antibiotic prescribing at primary care clinics in a veterans affairs healthcare system following a multifaceted stewardship intervention. Clin Infect Dis. 2020;71(8):e316-22.
 7. Shaikh N, Swaminathan N, Hooper EG. Accuracy and precision of the signs and symptoms of streptococcal pharyngitis in children: a systematic review. J Pediatr. 2012;160(3):487-93.
 8. Lee JJ, Verbakel JY, Goyder CR, et al. The clinical utility of point-of-care tests for influenza in ambulatory care: a systematic review and meta-analysis. Clin Infect Dis. 2019;69(1):24-33.
 9. Wiviott SD, Antman EM, Braunwald E. Prasugrel. Circulation. 2010;122(4):394-403.
 10. Barlow K, Thomson E, Johnson D, et al. Critical care outcomes. Current Opinion in Critical Care. 2006;12:489-526.