

# Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study.

Rashed Alam\*

Department of Population Science and Human Resource Development, University of Rajshahi, Rajshahi, Bangladesh

## Abstract

**Background:** Malnutrition is a significant public health issue in Bangladesh, particularly impacting women and children. Rajshahi, marked by socio-economic disparities, offers a distinctive context to explore the nutritional status and health outcomes of these vulnerable groups.

**Objectives:** This study aims to assess the nutritional status of women and children in Rajshahi and investigate associated health outcomes. Additionally, it seeks to identify socio-economic and cultural factors that influence nutrition.

**Methods:** A mixed-methods approach was utilized, incorporating a cross-sectional survey of 460 households and in-depth interviews with mothers and caregivers. Anthropometric measurements were taken to evaluate the nutritional status of women and children, while dietary assessments measured nutrient intake and diversity. Logistic regression analysis was performed to determine the likelihood of malnutrition based on socio-economic characteristics, thereby identifying key risk factors.

**Results:** The findings indicate a troubling prevalence of malnutrition, with 36% of children under five classified as stunted and 25% as underweight. The analysis highlights critical factors contributing to chronic undernutrition, including maternal education, employment and dietary diversity. Notably, mothers aged 27-37 exhibit a lower risk of undernutrition and urban households with secure food access demonstrate better nutritional outcomes. Moreover, frequent antenatal care visits and a diverse diet are vital in mitigating undernutrition risks among children.

**Conclusion:** This study emphasizes the urgent need for targeted interventions to combat malnutrition in Rajshahi. Recommendations include implementing community-based nutrition education programs and improving access to healthcare services. By addressing the socio-economic and cultural determinants of nutrition, stakeholders can enhance health outcomes for women and children in the region, ultimately contributing to broader public health objectives in Bangladesh.

**Keywords:** Socioeconomic inequalities, Nutritional status, Women, Children, Rajshahi district.

## Introduction

Bangladesh faces significant challenges in addressing the poor nutritional status of women and children, which is exacerbated by socioeconomic inequalities. The nutritional status of women, who are primarily responsible for food preparation and caregiving, directly impacts the health of their children. Socioeconomic disparities, particularly in rural and marginalized areas, contribute to substantial gaps in nutrition, influenced by factors such as gender, religion, culture and economic status. Gender inequality remains a key driver, with women historically facing systemic discrimination that limits their access to education, employment and healthcare. In Bangladesh's patriarchal society, traditional gender roles restrict women's social and economic mobility, making them economically dependent on men first their fathers, then their husbands and later their sons. This dependence, combined with

cultural norms that prioritize male well-being, results in gender-based nutritional disparities, with girls often receiving less nutritious food than boys, further deepening these inequalities [1].

The status of women in Bangladesh is also shaped by a patrilineal kinship system, which enforces their social and economic dependence on men. This system perpetuates the subordination and inequality of women, limiting their opportunities and health outcomes. Women are primarily seen as caregivers and mothers, with little attention paid to their own nutritional needs. As a result, poor maternal nutrition, including high rates of anemia, zinc deficiency and lack of vitamin B<sub>12</sub>, contributes to poor pregnancy outcomes, low birth weight and higher infant mortality rates. Poor maternal nutrition not only impacts fetal development but also contributes to intergenerational cycles of malnutrition, as malnourished

\*Correspondence to: Rashed Alam, Department of Population Science and Human Resource Development, University of Rajshahi, Rajshahi, Bangladesh; E-mail: mrasps29@ru.ac.bd

Received: 21-Nov-2024, Manuscript No. AAJFSN-24-153058; Editor assigned: 26-Nov-2024, AAJFSN-24-153058 (PQ); Reviewed: 10-Dec-2024, QC No. AAJFSN-24-153058; Revised: 17-Jun-2025, Manuscript No. AAJFSN-24-153058 (R); Published: 24-Jun-2025, DOI: 10.35841/aaifsn-8.3.299

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

mothers are more likely to give birth to undernourished children [2].

Undernutrition remains a major public health issue in Bangladesh, contributing to millions of deaths annually. Globally, undernutrition is linked to 3.5 million deaths each year, accounting for 35% of the disease burden in children under five. Bangladesh is among the 20 countries where 80% of the world's undernourished children live. High rates of undernutrition, especially among women of childbearing age, contribute to poor maternal and child health, leading to adverse birth outcomes and higher infant mortality. However, rising obesity rates are also emerging as a concern, with 16% of ever-married women and 10% of unmarried women aged 15-19 classified as overweight or obese. This dual burden of undernutrition and overnutrition reflects complex dietary shifts, particularly in urban areas [3].

Gender-based food allocation preferences play a significant role in malnutrition, particularly in Bangladesh, where boys often receive preferential treatment in food distribution, leaving girls with inadequate nutrition. This bias exacerbates the nutritional challenges faced by female children, who are more vulnerable to food insecurity and malnutrition. Such gender-based disparities in food allocation contribute to higher malnutrition rates among girls, reinforcing broader gender inequalities in health outcomes. Beyond food distribution, the nutritional status of women and children is also influenced by the availability and quality of healthcare services. In rural areas like Rajshahi, limited access to healthcare compounds the problem, preventing timely diagnosis and treatment of nutritional deficiencies. Poor healthcare infrastructure further hinders the ability of women and children to access essential services, while low-income households often struggle to afford balanced diets, relying on cheaper, less nutritious food options that exacerbate food insecurity and malnutrition [4].

Education, particularly among women, is crucial for improving nutritional outcomes. Educated mothers are more likely to adopt health-promoting behaviors, such as breastfeeding and providing nutritious meals for their children. However, in areas with limited access to education and healthcare, like rural Rajshahi, these benefits are less pronounced, perpetuating nutritional disparities. Addressing these gaps requires targeted interventions that enhance access to nutritious food, healthcare and education. Improving maternal nutrition, especially during pregnancy, is a key intervention to reduce poor pregnancy outcomes and promote better health for mothers and children. Studies highlight the need for region-specific strategies in districts like Rajshahi, where socioeconomic disparities and limited resources exacerbate health challenges [5].

The nutritional status of women and children in Bangladesh is strongly shaped by socioeconomic inequalities, including income, education, employment and healthcare access. Despite economic progress, significant disparities persist, especially in rural and marginalized areas. Addressing these inequalities through targeted policies is essential for improving health and nutrition and for advancing toward the SDGs. This study aims to assess the impact of socioeconomic status and healthcare

access on the nutritional outcomes of women and children in Rajshahi district, offering evidence-based guidance for future interventions (SDGs) [6].

## Materials and Methods

### Data

The study employs a combination of primary and secondary data to provide a comprehensive view of the nutritional landscape in Rajshahi district, focusing on three areas: Chorghat, Mohonpur, and Rajshahi city. Primary data is collected through field surveys and interviews, using a structured questionnaire designed to gather quantitative information on socioeconomic factors, dietary intake, and nutritional practices. Separate questionnaires are used for women and caregivers of children to address gender-specific and age-related issues. A stratified random sampling method ensures diverse representation across socioeconomic strata. Field enumerators conduct face-to-face interviews, ensuring accuracy and confidentiality, while additional qualitative data is gathered through focus group discussions and in-depth interviews with key stakeholders, such as community health workers and local policymakers. The data collection process is designed for reliability and clarity, with all questionnaires reviewed and translated into Bengali for ease of understanding. The study will also incorporate a cross-sectional design, recruiting healthcare professionals to interview respondents and ensuring high response rates through interviewer training and multiple household visits. A representative sample of 600 households will be selected from two Upazilas and Rajshahi city, with a focus on gathering data from 460 women [7].

### Variables

This study analyzes Body Mass Index (BMI) for women and the prevalence of undernutrition in children as dependent variables. For children, three growth indices are used: The Height-for-Age Z-score (HAZ), which reflects linear growth impairment and cumulative growth deficits; the Weight-for-Height Z-score (WHZ), which indicates recent or severe weight loss; and the Weight-for-Age Z-score (WAZ), a composite measure combining height-for-age and weight-for-height. WAZ can indicate both acute weight loss (wasting) and chronic growth failure (stunting) but does not differentiate between the two. Growth indices were calculated using the World Health Organization (WHO) 2006 growth standards, derived from the WHO multicentre growth reference study. Children with Z-scores below -2 Standard Deviations (SD) from the reference median were classified as undernourished.

The study includes three broad categories of independent variables: characteristics of children, mothers and households. Children's characteristics include sex, age (in months), birth weight (in grams) and whether the child has suffered from diarrhea. Maternal characteristics include age, education, working status, Antenatal Care (ANC) attendance, marital status and exclusive breastfeeding. Household characteristics

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

include religion, place of residence, family type, food security, water sources and minimum dietary diversity [8].

### Statistical analysis

Both univariate and multivariate techniques were used for data analysis. In the univariate analysis, categorized independent variables were cross-tabulated with dichotomized BMI to examine their association with severe malnutrition. Logistic regression was performed with dichotomized BMI and chronic undernutrition as dependent variables and all independent variables as categorized variables. A forward stepwise selection method was used to identify significant variables and construct the final model.

### Results

This study results Table 1 analyzed the associations between maternal and child characteristics and the nutritional status of mothers in Rajshahi, Bangladesh, as indicated by Body Mass Index (BMI). Several factors were found to significantly influence maternal BMI, including maternal age, education, employment, Antenatal Care (ANC), food security and water

access. Younger mothers (17-26 years) were more likely to be underweight, indicating vulnerability due to early marriage and limited education. Maternal education and employment were also linked to better nutritional outcomes, with educated and working mothers more likely to have a normal BMI. The number of ANC visits was a key factor, as mothers with regular check-ups had better nutritional status. Household food security emerged as a critical determinant; secure food access was associated with higher BMI, while food insecurity led to higher rates of underweight mothers. Safe water access and adequate dietary diversity were also correlated with better nutritional outcomes. Additionally, factors such as child birth weight, exclusive breastfeeding and childhood diarrhoea significantly impacted maternal BMI. The study emphasizes the need for targeted interventions in Rajshahi, focusing on improving education, healthcare access, food security and water quality to reduce maternal malnutrition, particularly among younger, rural and food-insecure populations [9].

**Table 1.** Association between maternal and child characteristics and nutritional status of women (BMI) in Rajshahi, Bangladesh (n=460).

Characteristics	Level	BMI			p-value
		Underweight	Normal weight	Total	
Maternal age	17-26	47 (35.1%)	87 (64.9%)	134 (29.1%)	0.009
	27-37	106 (37.1%)	180 (62.9%)	286 (62.2%)	
	≥ 38	18 (45.0%)	22 (55.0%)	40 (8.7%)	
Religion	Muslim	67 (23.2%)	222 (76.8%)	289 (62.8%)	0.039
	Non-Muslim	35 (20.5%)	136 (79.5%)	171 (37.2%)	
Education	Illiterate	26 (59.1%)	18 (40.9%)	44 (9.6%)	0.05
	Primary	32 (32.3%)	67 (67.7%)	99 (21.5%)	
	Secondary	97 (38.5%)	160 (61.5%)	257 (58.8%)	
	Higher	16 (26.6%)	44 (73.4%)	60 (13.1%)	
Respondents working status	Working	43 (40.9%)	62 (59.1%)	105 (22.8%)	0.04
	Not working	128 (36.1%)	227 (63.9%)	355 (77.2%)	
Place of living	Rajshahi city	49 (31.6%)	106 (68.4%)	155 (33.6%)	0.057
	Charghat	60 (39.5%)	92 (60.5%)	152 (33.0%)	
	Mohonpur	62 (40.5%)	91 (59.5%)	153 (33.3%)	
Maternal ANC	No visit	13 (30.9%)	29 (69.1%)	42 (9.1%)	0
	1-3 visit	87 (36.6%)	151 (63.4%)	238 (51.8%)	
	≥ 4 visit	71 (39.4%)	109 (60.6%)	180 (39.1%)	
Marital status	Married	158 (36.5%)	275 (63.5%)	433 (94.1%)	0.464
	Widowed	5 (45.5%)	6 (54.5%)	11 (2.3%)	
	Divorced	8 (50.0%)	8 (50.0%)	16 (3.6%)	
Type of family	Single-family	112 (32.7%)	230 (67.3%)	342 (74.3%)	0.069

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

	Joint family	59 (50.0%)	59 (50.0%)	118 (25.7%)	
Household food security status	Unsecure	111 (63.4%)	64 (36.6%)	175 (38.0%)	0.016
	Secure	60 (21.1%)	225 (78.9%)	285 (62.0%)	
Water sources	Safe	26 (9.4%)	251 (90.6%)	277 (60.2%)	0.005
	Unsafe	145 (79.2%)	38 (20.8%)	183 (39.8%)	
Minimum dietary diversity	Adequate	36 (12.9%)	243 (87.1%)	279 (60.7%)	0.025
	Inadequate	135 (74.6%)	46 (25.4%)	181 (39.3%)	
Child age, months	12-23	67 (32.8%)	137 (67.2%)	204 (43.3%)	0.057
	24-35	25 (33.8%)	49 (66.2%)	74 (16.1%)	
	36-47	36 (39.6%)	55 (60.4)	91 (19.8%)	
	48-59	43 (47.3)	48 (52.7%)	91 (19.8%)	
Child gender	Male	104 (42.4%)	141 (57.6%)	245 (53.3%)	0.008
	Female	67 (31.2%)	148 (68.8%)	215 (46.7%)	
Birth weight, gm	≥ 2500	20 (35.1%)	37 (64.9%)	57 (12.4%)	0.049
	<2500	151 (37.5%)	252 (62.5%)	403 (87.6%)	
Exclusive breastfeeding	Yes	67 (34.4%)	128 (65.6%)	195 (42.4%)	0.004
	No	104 (39.2%)	161 (60.8%)	265 (57.6%)	
Child suffering from diarrhea	Yes	15 (44.1%)	19 (55.9%)	34 (7.4)	0.001
	No	156 (36.6%)	270 (63.4%)	426 (92.6%)	

Table 2 presents the adjusted Odds Ratios (OR) for associations between sociodemographic, anthropometric and health-related characteristics and child chronic undernutrition in a study of 460 children. Maternal age was not significantly associated with child chronic undernutrition, with the odds ratios for mothers aged 27-37 years (OR=1.05, 95% CI: 0.66–1.67) and ≥ 38 years (OR=0.84, 95% CI: 0.47-1.49) showing no statistical significance (p=0.748 and p=0.556, respectively). Older child age (24-59 months) was significantly associated with higher odds of chronic undernutrition, with children aged 24-35 months (OR=2.40, 95% CI: 1.20-4.98), 36-47 months (OR=2.65, 95% CI: 1.29-5.92), and 48-59 months (OR=2.79, 95% CI: 1.34-5.96) having significantly higher odds compared to children aged 12-23 months (p=0.004, p=0.002 and p=0.001, respectively). No significant association was found between child gender (OR=0.91, 95% CI: 0.66-1.78, p=0.448) or water source safety (OR=0.87, 95% CI: 0.71-1.22, p=0.281) and chronic undernutrition. However, maternal education was protective, with children of mothers with secondary education (OR=0.78, 95% CI: 0.56-0.87, p=0.009) or higher education (OR=0.59, 95% CI: 0.43-0.74, p<0.001) having significantly lower odds of chronic undernutrition compared to children of mothers with pre-primary or no education. Employed mothers also had lower odds of child chronic undernutrition (OR=0.69,

95% CI: 0.43-0.85, p=0.030). Maternal Antenatal Care (ANC) visits showed that women with ≥ 4 visits had lower odds of chronic undernutrition in their children (OR=0.67, 95% CI: 0.53-0.89, p<0.001). Children from food-secure households had significantly lower odds of chronic undernutrition (OR=0.77, 95% CI: 0.64-0.91, p<0.001). Low birth weight (<2500 gm) was associated with higher odds of chronic undernutrition (OR=1.23, 95% CI: 1.08-1.44, p=0.008), as was maternal chronic undernutrition (OR=1.81, 95% CI: 1.61-2.51, p=0.007). Adequate dietary diversity was also protective (OR=0.68, 95% CI: 0.55-0.94, p=0.006). Exclusive breastfeeding showed no significant effect on child chronic undernutrition (OR=0.97, 95% CI: 0.78-1.87, p=0.654), while diarrhea in children was a significant risk factor (OR=1.61, 95% CI: 1.31-1.95, p=0.015). These findings underscore the importance of maternal education, employment, ANC visits, food security, dietary diversity and birth weight in mitigating child chronic undernutrition, while highlighting the risks posed by maternal undernutrition and child diarrhea.

**Table 2.** Intensity of the effects of odds ratio of various child, maternal and household characteristics on child chronic undernutrition: A logistic regression analysis (n=460).

Characteristics	Child chronic undernutrition	P-value
	Adjusted odds ratio (95% CI)	

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

<b>Maternal age, yrs (ref=17-26)</b>		
27-37	1.05 (0.66-1.67)	0.748
≥ 38	0.84 (0.47-1.49)	0.556
<b>Place of residence (ref=Charghat)</b>		
Rajshahi city area	0.55 (0.42, 0.78)	0.005
Mohonpur	0.97 (0.38, 0.99)	0.001
<b>Child age, months (ref=12-23)</b>		
24-35	2.40 (1.20, 4.98)	0.004
36-47	2.65 (1.29, 5.92)	0.002
48-59	2.79 (1.34, 5.96)	0.001
<b>Child gender (ref=female)</b>		
Male	0.91 (0.66, 1.78)	0.448
<b>Maternal education (ref=pre-primary or none)</b>		
Primary	0.98 (0.78, 1.77)	0.361
Secondary	0.78 (0.56, 0.87)	0.009
Higher	0.59 (0.43-0.74)	<0.001
<b>Maternal current job (ref=no)</b>		
Yes	0.69 (0.43, 0.85)	0.03
<b>Maternal ANC (ref=no visit)</b>		
1-3 visit	0.93 (0.79, 1.94)	0.312
≥ 4 visit	0.67 (0.53, 0.89)	<0.001
<b>Household food security status (ref=unsecure)</b>		
Secure	0.77 (0.64, 0.91)	<0.001
<b>Water sources (ref=unsafe)</b>		
Safe	0.87 (0.71, 1.22)	0.281
<b>Birth weight, gm (ref= ≥ 2500)</b>		
<2500	1.23 (1.08, 1.44)	0.008
<b>Maternal chronic undernutrition (ref=no)</b>		
Yes	1.81 (1.61, 2.51)	0.007
<b>Minimum dietary diversity (ref=inadequate)</b>		
Adequate	0.68 (0.55, 0.94)	0.006
<b>Exclusive breastfeeding (ref=no)</b>		
Yes	0.97 (0.78, 1.87)	0.654
<b>Child suffering from diarrhea (ref=no)</b>		
Yes	1.61 (1.31, 1.95)	0.015
<b>Note:</b> CI: Confidence Interval		

Table 3 presents the adjusted Odds Ratios (OR) for sociodemographic, anthropometric and health-related characteristics associated with maternal chronic undernutrition (BMI<18.5) in a study of 460 women. Maternal age was a

significant factor, with women aged 27-37 years having significantly lower odds of chronic undernutrition (OR=0.15, 95% CI: 0.03-0.93, p=0.041) compared to women aged 17-26 years. However, no significant association was found for

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

women aged  $\geq 38$  years (OR=0.87, 95% CI: 0.46-1.94,  $p=0.739$ ). Regarding place of residence, women in Rajshahi City had lower odds of chronic undernutrition compared to those in Charghat (OR=0.95, 95% CI: 0.82-0.98,  $p=0.005$ ), while those in Mohonpur had significantly higher odds (OR=0.57, 95% CI: 0.28-0.89,  $p=0.021$ ). Maternal employment did not show a significant association with chronic undernutrition (OR=0.87, 95% CI: 0.39-1.94,  $p=0.740$ ). Women with  $\geq 4$  Antenatal Care (ANC) visits had significantly lower odds of chronic undernutrition (OR=0.37, 95% CI: 0.16-0.86,  $p=0.021$ ), while those with 1-3 visits showed no significant difference (OR=0.65, 95% CI: 0.22-1.91,  $p=0.445$ ). Household food security was a protective factor, with food-secure women having significantly lower odds of chronic undernutrition (OR=0.34, 95% CI: 0.17-0.67,  $p=0.002$ ). No significant association was found between access to safe water and maternal chronic undernutrition (OR=0.98,

95% CI: 0.33-2.88,  $p=0.138$ ). Women with adequate dietary diversity had significantly lower odds of chronic undernutrition (OR=0.18, 95% CI: 0.04-0.76,  $p=0.020$ ). In summary, maternal chronic undernutrition is influenced by factors such as maternal age, place of residence, ANC visits, household food security, and dietary diversity. Protective factors include older maternal age (27-37 years), living in Rajshahi city and having adequate ANC visits, food security and dietary diversity. In contrast, living in Mohonpur and inadequate dietary diversity were associated with higher odds of undernutrition, while maternal employment and access to safe water did not significantly affect chronic undernutrition.

**Table 3.** Intensity of the effects of odds ratio of various child, maternal and household characteristics on maternal chronic undernutrition: A logistic regression analysis (BMI<18.5) (n=460).

Characteristics	Maternal chronic undernutrition	P-value
	Adjusted odds ratio (95% CI)	
<b>Maternal age, yrs (ref=17-26)</b>		
27-37	0.15 (0.03-0.93)	0.041
$\geq 38$	0.87 (0.46-1.94)	0.739
<b>Place of residence (ref=Charghat)</b>		
Rajshahi city area	0.95 (0.82, 0.98)	0.005
Mohonpur	0.57 (0.28, 0.89)	0.021
<b>Maternal current job (ref=no)</b>		
Yes	0.87 (0.39, 1.94)	0.74
<b>Maternal ANC (ref=no visit)</b>		
1-3 visit	0.65 (0.22, 1.91)	0.445
$\geq 4$ visit	0.37 (0.16, 0.86)	0.021
<b>Household food security status (ref=unsecure)</b>		
Secure	0.34 (0.17, 0.67)	0.002
<b>Water sources (ref=unsafe)</b>		
Safe	0.98 (0.33, 2.88)	0.138
<b>Minimum dietary diversity (ref=inadequate)</b>		
Adequate	0.18 (0.04, 0.76)	0.02
<b>Note:</b> CI: Confidence Interval		

## Discussion

This study explores several key factors influencing maternal and child nutrition in Bangladesh, with a particular focus on maternal age, education, employment, antenatal care, food security, dietary diversity, birth weight and childhood diarrhea. Below is an analysis of the findings, with supporting references that highlight global trends and context-specific challenges.

The study finds that mothers aged 27-37 years have significantly lower odds of chronic undernutrition compared to

younger mothers (AOR=0.15,  $p=0.041$ ). This suggests that women in this age group are better equipped to handle the nutritional demands of pregnancy. In contrast, mothers aged 38 years or older do not show a significant difference in undernutrition (AOR=0.87,  $p=0.739$ ). This aligns with research in Bangladesh, where younger maternal age (17-26 years) is linked to higher risks of undernutrition due to biological, socioeconomic and experiential factors. Similarly, studies in India and other South Asian countries have highlighted that younger mothers are more vulnerable to poor health outcomes,

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

including low birth weight and maternal anemia. These findings emphasize the need for targeted nutrition interventions for younger mothers, as their age and related vulnerabilities significantly impact their nutritional status and child health.

The study highlights the urban advantage in reducing child undernutrition, with Rajshahi city outperforming rural areas like Charghat. Urban areas generally offer better healthcare, nutrition programs and socioeconomic resources, all of which contribute to improved health outcomes. This is consistent with global trends, where urban children often have better access to healthcare and nutrition, resulting in lower rates of undernutrition. However, the study also finds that children in Mohonpur, another rural area, do not show significant differences in undernutrition compared to those in Charghat, suggesting that some rural communities may benefit from local resources, community networks or agricultural activities that help mitigate nutritional risks. While urbanization generally improves access to services, it can also introduce challenges, such as pollution and dietary shifts, which could negatively affect nutritional outcomes. This complex interplay between urban and rural health highlights the need for context-specific interventions that consider local resources and disparities in healthcare access and food security.

Maternal education is a powerful determinant of child nutritional status. The study reveals that mothers with secondary and higher education are significantly less likely to have chronically undernourished children. Specifically, mothers with secondary education have an adjusted Odds Ratio (OR) of 0.78 ( $p=0.009$ ) and those with higher education show even lower odds ( $OR=0.59$ ,  $p<0.001$ ). This indicates that educated mothers are more likely to make informed decisions about nutrition and healthcare. Educated mothers are also more likely to secure better employment, increasing their ability to afford nutritious food and healthcare. Studies from Bangladesh, India and other countries have consistently shown that maternal education reduces child undernutrition and improves health outcomes.

However, primary education did not show a significant effect on reducing child chronic undernutrition ( $OR=0.98$ ,  $p=0.361$ ). This suggests that while primary education offers some benefits, it may not be as impactful as secondary or higher education in shaping maternal and child nutrition outcomes. Investing in maternal education is thus critical not only for individual families but also for public health improvements across society.

Both maternal education and employment significantly reduce the likelihood of child chronic undernutrition. The study finds that employed mothers are 31% less likely to have undernourished children compared to non-employed mothers ( $OR=0.69$ ,  $p=0.030$ ). Employed mothers tend to have higher incomes, which enhances food security and enables them to purchase nutritious food and access healthcare services more readily. They are also more likely to attend Antenatal Care (ANC) and ensure their children receive necessary health services, including vaccinations.

Similar results have been observed in India and Ethiopia, where maternal employment is associated with better nutritional outcomes for children. Employment also provides economic empowerment, which helps alleviate food scarcity and increases dietary diversity for children. These findings highlight the importance of increasing maternal employment opportunities, especially in low-income settings like Bangladesh, where such opportunities can help break the cycle of malnutrition.

The study also highlights the role of antenatal care in reducing maternal chronic undernutrition. Attending four or more ANC visits is associated with significantly lower odds of maternal undernutrition ( $AOR=0.37$ ,  $p=0.021$ ). In Bangladesh, ANC is crucial in providing essential nutritional guidance, healthcare, and medical support during pregnancy. Women who attend fewer than four ANC visits (1-3 visits) do not show significant improvements in nutritional status ( $AOR=0.65$ ,  $p=0.445$ ), indicating that more frequent and comprehensive ANC visits are necessary to address maternal nutrition adequately. This finding is supported by studies in India, where women with more ANC visits experienced better nutritional outcomes and health.

The study finds that households with secure food status significantly reduce the likelihood of maternal chronic undernutrition ( $AOR=0.34$ ,  $p=0.002$ ). This is a critical finding in Bangladesh, where food insecurity is a major barrier to maternal and child nutrition, leading to poor health outcomes. Secure food environments are crucial in providing the necessary nutrients for pregnant and lactating women, which in turn supports their health and the health of their infants. Research in other countries, such as Ethiopia, supports the direct link between food security and maternal health outcomes. Targeted interventions to improve food security can play a critical role in reducing maternal undernutrition globally.

Maternal undernutrition is a significant factor influencing child nutritional status. The study finds that undernourished mothers are 81% more likely to have chronically undernourished children ( $OR=1.81$ , 95% CI: 1.61-2.51,  $P=0.007$ ). In Bangladesh, where maternal undernutrition affects over 30% of women, this relationship is especially concerning, as it perpetuates intergenerational cycles of malnutrition. Undernourished mothers are more likely to give birth to Low Birth Weight (LBW) infants, who are at higher risk for chronic undernutrition during childhood. Addressing maternal undernutrition through comprehensive interventions is essential for breaking the cycle of malnutrition and improving child health outcomes.

The study reveals that dietary diversity is significantly associated with reduced chronic undernutrition. Individuals with more diverse diets have a 32% lower likelihood of experiencing chronic undernutrition ( $OR=0.68$ ,  $p=0.006$ ). In Bangladesh, where malnutrition is a significant public health challenge, promoting dietary diversity is essential for improving nutrition, particularly for children. Increased dietary diversity ensures adequate intake of essential vitamins and

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

minerals that are often missing in less varied diets. Studies in India and Nigeria show similar findings, where children with more diverse diets are less likely to suffer from malnutrition.

The study also finds that dietary diversity significantly reduces maternal chronic undernutrition (AOR=0.18,  $p=0.020$ ). Women with diverse diets are more likely to meet their nutritional needs during pregnancy and lactation, reducing the risk of undernutrition and promoting maternal and fetal health. Similar studies in Ethiopia have found that pregnant women with higher dietary diversity had better nutritional outcomes and lower rates of anemia. These findings support the need for global interventions to improve dietary diversity as a strategy to reduce undernutrition and improve maternal and child health outcomes.

Birth weight is another critical determinant of child nutrition. The study found that Low Birth Weight (LBW) is significantly associated with higher odds of chronic undernutrition (OR=1.23, 95% CI: 1.08-1.44,  $P=0.008$ ). In Bangladesh, where approximately 28% of infants are born with LBW, addressing the determinants of LBW is crucial for improving child nutrition and health. Globally, LBW is a well-established risk factor for developmental delays, malnutrition and mortality. Research from India and sub-Saharan Africa supports this link between LBW and chronic undernutrition. Targeted maternal health interventions to improve nutrition during pregnancy can help improve birth outcomes and reduce the incidence of LBW.

Childhood diarrhoea is a significant public health issue that is closely linked to malnutrition, with an odds ratio of 1.61 (95% CI: 1.31-1.95,  $P=0.015$ ) indicating that children suffering from diarrhea are 61% more likely to experience chronic undernutrition compared to those who do not. In Bangladesh, diarrhea remains a leading cause of morbidity and mortality among children under five, exacerbated by inadequate access to clean water, sanitation and hygiene practices. The frequent loss of nutrients due to diarrheal illness can lead to a vicious cycle of malnutrition, where undernourished children are more susceptible to infections and the resulting illnesses further compromise their nutritional status. Globally, this relationship is well-documented. For example, studies in sub-Saharan Africa and South Asia have shown that children with diarrhea experience higher rates of stunting and wasting, with recurrent episodes contributing to long-term growth deficits. Interventions aimed at reducing diarrheal diseases, such as improved sanitation, access to clean water and education on hygiene, are essential for breaking this cycle and improving child health outcomes. Addressing the dual challenges of diarrhea and malnutrition is crucial for enhancing child survival and development, particularly in resource-limited settings like Bangladesh.

### **Policy implication**

The study's findings highlight critical areas for policy intervention to address nutritional disparities in Rajshahi, Bangladesh. Policymakers should prioritize educational advancement, especially for mothers, as higher education is linked to improved nutritional practices and health outcomes.

Expanding healthcare accessibility, particularly maternal and child health services, is essential for providing timely interventions. Strengthening food security strategies through government and community programs is necessary to ensure consistent access to nutritious food for all households. Additionally, investments in sanitation and hygiene infrastructure are crucial to mitigating health risks associated with undernutrition, particularly among children. The recommendations emphasize targeted educational programs to empower women with knowledge, enhancing healthcare services in underserved areas, supporting local food production initiatives, fostering community engagement in health programs and establishing robust monitoring and evaluation frameworks to track progress. By implementing these strategies, policymakers can reduce socioeconomic inequalities in nutritional status and promote better health outcomes for women and children in the region, contributing to a more equitable society.

### **Conclusion**

This study investigates the socioeconomic factors impacting the nutritional status of women and children in Rajshahi District, Bangladesh, highlighting persistent malnutrition despite public health progress. The study, employing a mixed-methods approach, finds that women from lower socioeconomic backgrounds face higher malnutrition rates, adversely affecting their children's health. Maternal education was identified as a key factor in improving child nutrition, while cultural norms and gender roles were barriers to accessing adequate nutrition and healthcare. Disparities in healthcare access, exacerbated by financial constraints and poor infrastructure, further worsen the situation. The study calls for policies focused on improving women's education, healthcare access, and community-based nutrition programs. It advocates for a multi-sectoral approach to address both immediate and underlying factors contributing to malnutrition, offering a framework for tackling these issues and promoting health equity.

### **References**

1. Abdulahi M. Maternal age and its association with undernutrition in Ethiopia: A population-based study. *BMC Public Health*. 2019;19:1204.
2. Ahmed T, Mahfuz M, Ireen S, et al. Nutrition of children and women in Bangladesh: Trends and directions for the future. *J Health Popul Nutr*. 2012;30:1.
3. Bhandari N. Maternal education and child nutrition in Bangladesh: A cross-sectional study. *BMC Publ Health*. 2016;16:673.
4. Bhutta ZA, Das JK, Rizvi A, et al. Evidence-based interventions for improvement of maternal and child nutrition: What can be done and at what cost?. *Lancet*. 2013;382:452-77.
5. Black RE, Allen LH, Bhutta ZA, et al. Maternal and child undernutrition: Global and regional exposures and health consequences. *Lancet*. 2008;371:243-60.

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299

6. Bryce J, Coitinho D, Darnton-Hill I, et al. Maternal and child undernutrition: Effective action at national level. *Lancet*. 2008;371:510-26.
7. Burgard SA, Lin IF. The importance of maternal education for children's health in Bangladesh. *Soc Sci Med*. 2013;79:1-8.
8. Checkley W, Buckley G, Gilman RH, et al. Childhood malnutrition and infection network. Multi-country analysis of the effects of diarrhoea on childhood stunting. *Int J Epidemiol*. 2008;37:816-30.
9. Fagbule D. The impact of diarrheal diseases on the nutritional status of children under five in Nigeria. *Nigerian J Clin Prac*. 2018;21:431-6.

**Citation:** Alam R. Assessing the nutritional status and health outcomes of women and children in Rajshahi, Bangladesh: A comprehensive study. *J Food Sci Nutr* 2025;8(3):299