

## Cantharidin content determination and its inhibitory effect on human hepatoma HepG2 cells.

Yisheng Lin<sup>1,2</sup>, Yanhao Lia<sup>1\*</sup>, Zhenpeng Peng<sup>3</sup>, Shilong Yu<sup>4</sup>, Shuangfeng Lin<sup>5</sup>, Xiaoshu Huc<sup>3\*</sup>

<sup>1</sup>Department of Interventional Radiology, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong Province, PR. China

<sup>2</sup>Department of Interventional Radiology, the First Affiliated Hospital Of Guangzhou University Of Traditional Chinese Medicine, Guangzhou, Guangdong Province, PR. China

<sup>3</sup>Department of Radiology, the First Affiliated Hospital, Sun Yat-Sen University, Guangzhou, Guangdong, China

<sup>4</sup>Jilin Cancer Hospital, Changchun, Jilin province, PR. China

<sup>5</sup>Laboratory Animal Center, the First Affiliated Hospital Of Guangzhou University Of Traditional Chinese Medicine, Guangzhou, Guangdong Province, PR. China

### Abstract

To establish a method for determination of cantharidin content in *Mylabris phalerata. Pallas*, and to study its inhibitory and apoptosis-inducing effects on human hepatoma cell line HepG2. Agilent column (TC-C18, 250 mm × 4.6 mm, 5 μm); mobile phase: methanol-water (volume ratio: 20:80); flow rate: 1.0 ml/min; UV detection wavelength: 230 nm; and column temperature: 30°C. MTT assay was used to determine cell viability, and flow cytometry was used to detect the effects of cantharidin on cell cycle arrest and apoptosis of HepG2 cells. Cantharidin had a good linearity (R=0.9993) within a 0.056~0.504 μg range, and had an average recovery of 98.55%. Different concentrations of cantharidin all markedly induced the apoptosis of human hepatoma HepG2 cells. After treatment for 24, 36 and 48 h, respectively, HepG2 cells were inhibited to varying degrees in a dose and time dependent manner. PI staining with flow cytometry showed that the test concentrations of cantharidin could induce HepG2 cell apoptosis, and impact the cell cycle to varying degrees by increasing the proportion of G0/G1 phase cells. Compared with the control group, apoptosis was positively correlated with drug concentration, presenting a marked dose-dependence. The method proposed is simple, fast, accurate and reproducible, which can be used for the determination of cantharidin content in *Mylabris phalerata. Pallas*. Cantharidin has an inhibitory effect on HepG2 cells.

**Keywords:** Cantharidin, Flow cytometry, HepG2 cell.

Accepted on February 16, 2016

### Introduction

Banmao is the dried body of Meloidae insects *Mylabris phalerata. Pallas* or *Mylabris cichorii. Linnaeus* [1]. It is pungent, hot, and strongly toxic, which enters the liver, stomach and kidney meridians, and has blood stasis-eliminating, mass-resolving, swelling-subsiding, toxicity-counteracting and sore-removing effects [2]. Originally recorded in the Shen Nong's Herbal Classic, Banmao belongs to the family Meloidae of the order Coleoptera, which is one of the earliest discovered medicinal materials with anti-tumor action. Medicinal value of Banmao lies in its ability to effectively remove dead skins, and be applied on malignant sores and scabies [3].

Modern research has shown that cantharidin contained in Banmao has a strong anti-tumor activity, which can be used for carbuncles, gangrenes, malignant sores, scrofula, rabies, as well as abortion; domestic and foreign scholars have done extensive research on the mechanism of action of cantharidin [4-8]. As the major active constituent of Banmao [9-10], cantharidin also has strong toxicity, so rational application of cantharidin and strict control of its dose has become the focus of research and development of cantharidin and its derivatives.

### Material and Methods

#### *Instruments and reagents*

Agilent 1200 HPLC system; ELX 800 microplate reader (BioTEK, USA); FACSAria flow cytometer (BD, USA);

CO-150 CO<sub>2</sub> incubator (NBS, USA); RPMI1640 (Gibco); FBS (Hangzhou Sijiqing Bioengineering Materials Co., Ltd.); MTT (Sigma).

### Drug and cell line

Banmao medicinal material was purchased from Anguo market, which was identified as *Mylabris phalerata*. Pallas. Human HepG2 cell line was purchased from the Cell Bank at SIBCB.

### Chromatographic conditions

Agilent column (TC-C18, 250 mm × 4.6 mm, 5 μm); mobile phase: methanol-water (volume ratio: 20:80); flow rate: 1.0 ml/min; UV detection wavelength: 230 nm; and column temperature: 30°C [11,12]. The results are shown in figure 1.

### Linearity range

**Preparation of reference solution:** Cantharidin reference substance, which was dried to constant weight under vacuum at 45°C, was accurately weighed, and added with methanol to prepare a 0.056 mg/ml reference solution.

**Preparation of test solution:** About 2 g of *Mylabris phalerata* Pallas medicinal powder (sifted through a 20 mesh sieve) was accurately weighed, placed in a stoppered conical flask, added precisely with 50 mL of HCl/acetone solution, weighed, ultrasonicated for 45 min, then removed and let cool. After replenishing the lost weight, the solution was filtered through a 0.45 μm microporous membrane to give the test solution.

**Linearity range:** 1, 3, 5, 7 and 9 μL of cantharidin reference solution was accurately drawn, and injected into HPLC system for measurement of peak area. Standard curve was plotted with mass concentration (X) versus peak area (Y), and regression equation was  $y = 7853.9 \times -254.346$ ,  $r = 0.9993$ , suggesting good linearity of cantharidin within a 0.056~0.504 μg range.




Figure 1. HPLC chromatograms (reference substance; test sample).

### Stability test

100 μL of the same test solution was taken, and injected separately at 0, 1, 2, 3, 4, 5 and 24 h for measurement. The results showed that RSD=1.12%, indicating that the sample solution was stable within 24 h.

### Accuracy test

The same test solution was precisely drawn and injected six times repeatedly for measurement of peak area. RSD was

found to be 1.14%, demonstrating good accuracy of the instrument.

### Recovery test

*Mylabris phalerata* Pallas medicinal materials with known contents of samples were accurately weighed, added separately with appropriate amount of cantharidin, prepared as per the method under preparation of test solution, and determined by the method under content determination, followed by calculation of recovery. The results revealed an average recovery of 98.55%, indicating that the method has good recovery. The results are shown in table 1.

Table 1. Cantharidin recovery test results.

No.	Sampling amount (g)	Sample content (mg)	Reference amount (mg)	Measured amount (mg)	Recovery (%)
1	0.5563	5.248	5.215	10.38	99.2
2	0.5542	5.244	5.215	10.35	98.9
3	0.5528	5.221	5.215	10.33	98.9
4	0.5531	5.232	5.215	10.28	98.4
5	0.5519	5.228	5.215	10.31	98.7
6	0.5538	5.243	5.215	10.17	97.2

### Reproducibility test

Five aliquots of the same batch of medicinal materials were accurately weighed, and prepared into samples as per the method under preparation of test solution. 5 μL of test solution was precisely drawn, and injected repeatedly into HPLC for determination under chromatographic conditions described above. The results found that RSD was 0.38%, indicating good reproducibility of the method.

### Content determination

Three batches of *Mylabris* medicinal materials with different origins were taken, prepared into samples as per the test solution preparation method, and determined under the above chromatographic conditions, followed by calculation of cantharidin contents. The results are shown in table 2.

Table 2. Cantharidin content.

Batch No	Sample amount (g)	Cantharidin content (mg/g)
20150301	5.548	10.12
20150302	5.539	10.08
20150303	5.562	10.24

## Results

Inhibitory effect of cantharidin on HepG2 cell growth

**Cell culturing**

HepG2 cells were cultured in a 37°C, 5% CO<sub>2</sub> incubator with 10% FBS-containing DMEM medium. The medium was replaced once in every 3 days. Logarithmic phase cells were harvested for the experiment.

**MTT assay of HepG2 cell growth inhibition rate by cantharidin**

Well grown exponential phase cells were prepared into a 5 × 10<sup>4</sup>/ml cell suspension, and seeded in a 96 well plate at 200 µl per well. The cells were treatment with different concentrations of cantharidin (10, 20, 40 µg/ml), and three replicate wells were set up for each concentration. After culturing for 24, 36 and 48 h, each well was added with 20 µL of 5 mg/ml MTT solution, and incubated for an additional 4 h then supernatant was discarded. Each well was added with 150 µL of DMSO, shaken, and then A value was measured at 570 nm. The experiment was repeated three times, and growth inhibition rate was calculated.

**Growth inhibition rate**

$$= \frac{A \text{ value of negative control group} - A \text{ value of treatment group}}{A \text{ value of negative control group}} \times 100\%$$

Different concentrations of cantharidin could all markedly induce HepG2 cell apoptosis. After treating for 24, 36 and 48 h, HepG2 cells were inhibited to varying degrees in a dose and time dependent manner.




Figure 2. HepG2 cell growth inhibition rate by cantharidin.

**Determination of cell cycle and apoptosis rate**

Logarithmic phase HepG2 cells were taken, trypsinized, prepared into a 5 × 10<sup>6</sup> cell suspension, added with different concentrations of cantharidin samples (10, 20, 40 µg/ml), and cultured for 24 h. After trypsinization, centrifugation and supernatant removal, the cells were added with 2 mL of sodium citrate buffer, water bathed at 37°C for 30 min, then stained with 1.5 mL of PI, and kept in a dark place for 30 min. Afterwards, the cells were passed through 300-mesh nylon net six times, and analyzed for cell cycle distribution by flow cytometry.

The results are shown in table 3. PI staining with flow cytometry revealed that the test concentrations of cantharidin

could induce HepG2 cell apoptosis. Cell cycle was impacted to varying degrees, with relative increase in the proportion of G0/G1 phase cells. Compared with the control group, apoptosis was positively correlated with the drug concentration, presenting a marked dose-dependence.

Table 3. Effect of cantharidin on cell cycle of HepG2 cells (n=6,  $\bar{x} \pm s$ ).

Group	Concentration (µg/ml)	G0/ G1 phase	S phase	G2/ M phase
Control group	-	72.34 ± 0.35	13.73 ± 0.44	13.93 ± 0.49
	10	76.64 ± 0.41	11.48 ± 0.56	11.88 ± 0.62
Cantharidin	20	78.86 ± 0.52	10.46 ± 0.36	10.68 ± 0.55
	40	82.42 ± 0.74	10.27 ± 0.48	07.31 ± 0.64

**Discussion**

Liver cancer is a major killer threatening human health among malignancies. Most liver cancers are diagnosed at advanced stages, where the best time for treatment has been lost, thus tremendously affecting the patients and their families. At present, the major treatment for cancer is chemotherapy, especially for advanced cancer, where chemotherapy has almost become a mandatory clinical therapy. However, the majority of chemotherapeutics have strong cytotoxicity on normal cells, which suppress the development of cancer cells while causing enormous damage to the patients' bodies.

Primary liver cancer ranks top among malignancies in terms of mortality. How to effectively prevent liver cancer is a crucial issue faced by the medical and pharmaceutical industries. Chinese medicine generally believes that one major cause of liver cancer is the pathogenic toxins, how to effectively dispel the pathogenic toxins is an important way of treating cancer.

Pang Jie et al., [13] confirmed by experimental study that the disodium cantharidinate vitamin B6 injection directly inhibited H22 xenograft tumor, killed H22 cells in vitro, induced apoptosis and prolonged the survival of ascetic tumor-bearing mice; furthermore, these effects were enhanced with increasing dose. The anti-hepatoma mechanism of disodium cantharidinate vitamin B6 injection is believed to be exerted not only by inhibiting and killing tumor cells, but also by regulating the immune function.

Cao Yongyan et al., [14] prepared compound Banmao capsule serum by serum pharmacological approach, and used it to treat human hepatoma SMMC-7721 cells. They determined the proteins in cancer cells influenced by compound Banmao capsule using proteomic techniques such as two-dimensional electrophoresis, image analysis and mass spectrometry, and found that the anti-cancer mechanism of compound Banmao capsule is a process jointly participated by multiple channels, which involves multiple targets in the proliferative, immunologic and apoptotic aspects of tumors.

## Conclusion

In this study, the anti-cancer activity of Chinese medicine Banmao is explored centering on cantharidin. MTT assay and flow cytometry results demonstrate that the test concentrations of cantharidin can induce HepG2 cell apoptosis, and influence cell cycle to varying degrees, indicating that cantharidin has a growth inhibitory effect on tumor cells. But whether it can regulate immune function still needs further confirmation.

## References

1. Chinese Pharmacopoeia Commission. Pharmacopoeia of the People's Republic of China (2010 edition, Vol. I). Beijing: China Medical Science Press 2010; pp. 331-332.
2. Hu CY. Research Progress on the Quantitative Determination of Cantharidin. Guangzhou Chemical Industry 2011; 39: 49-51.
3. Liu J, Gao JH, Liu XQ. Progress in the research of cantharidin and its derivatives. J Chinese Med Materl 2003; 26: 453-455.
4. DjehaH A, Todryk SM, Pelech S, Wrighton C, Irvine A. Antitumor immune responses mediated by adenoviral GDEPT using nitroreductase/CB1954 is enhanced by high-level coexpression of heat shock protein 70. Cancer Gene Ther 2005; 12: 560-571.
5. Bradley G, Juranka PF, Ling V. Mechanism of multidrug resistance. Biophys Acta 1988; 948: 87-128.
6. King TA, Ghazaleh RA, Juhn SK, Adams GL, Ondrey FG. Induction of heat shock protein 70 inhibits NF-kappa-B in squamous cell carcinoma. Otolaryngol Head Neck Surg 2005; 133: 70-79.
7. Tchounwou PB, Yedjou CG, Dorsey WC. Arsenic trioxide-induced transcriptional activation of stress genes and expression of related proteins in human liver carcinoma cells (HepG2). Cell Mol Biol 2003; 49: 1071-1079.
8. Calderwood SK, Theriault JR, Gong J. Message in a bottle: role of the 70-kDa heat shock protein family in anti tumor immunity. Eur J Immunol 2005; 35: 2518-2527.
9. Nikbakhtzadeh MR, Tirgari S. Cantharidin Component of Iranian Blister Beetles (Col: Meloidae) and their differences between Iranian and Exotic Species. Iranian J Pub Health 2002; 31: 113-117.
10. Yang JH, Chen JW, Xu JG, Lu LL, Li X. Comparative Analysis of Constituents of Hydrolysis Product of Mylabris phalerata Pallas in Different Producing Areas. Nat Product Res Dev 2009; 21: 449-452.
11. Ge XZ, Du XX, Yu Y. Extraction of Cantharidin from Mylabris phalerata Pallas with Compound Enzyme and HPLC Analysis. Asian J Chem 2013; 25: 123-126.
12. Dang YJ, An LN, Hu CH, Zhu CY. Inclusion Complex of Cantharidin with beta-Cyclodextrin: Preparation, Characterization, In Vitro and In Vivo Evaluation. J Appl Poly Sci 2012; 123: 1557-1562.
13. Pang J, Cao Y, Chen RS. Effect of disodium cantharidinate and vitamin B6 injection on hepatocarcinoma 22 cell line. J Practical Oncol 2010; 25: 693-697.
14. Cao YY, Wang M, Huang KS, Yang Q, Ding M. Proteomic analysis of differentially expressed proteins in human hepatoma SMMC-7721 cells induced by Fufang Banmao capsule serum. J Chinese Materia Medica 2007; 32: 831-834.

### \*Correspondence to:

Yanhao Lia  
 Department of Interventional Radiology  
 Nanfang Hospital  
 Southern Medical University  
 PR. China  
 Xiaoshu Huc  
 Department of Radiology  
 The First Affiliated Hospital  
 Sun Yat-Sen University  
 PR. China