

Dermatological evaluation of counter irritant and anti-inflammatory effect of ethanolic extract of seed of *Mangifera Indica* in rabbits.

Ameen Abbas^{1*}, Amna Ramzan¹, Eaman Ateeq¹, Usama Fiyaz¹, Mashhud-ul-Hasan², Nazish Ramzan³, Alisha Rehman⁴

¹Department of Biochemistry, Muhammad Institute of Medical and Allied Sciences, Multan, Pakistan

²Department of Biochemistry, Baha Uddin Zakariya University, Multan, Pakistan

³Department of Chemistry, Baha Uddin Zakariya University, Multan, Pakistan

⁴Department of Physiotherapy, Muhammad Institute of medical and Allied Sciences, Multan, Pakistan

Received: 10-October-2019, *Manuscript No.* AABB-19-3432; **Editor assigned:** 16-October-2019, *PreQC No.* AABB-19-3432; **Reviewed:** 30-October-2019, *QC No.* AABB-19-3432; **Revised:** 25-August-2022, (*PQ*) *Manuscript No.* AABB-19-3432; **Published:** 22-September-2022; DOI: 10.3451/aabb.5.5.120.

Abstract

Objective: The assessment of the effect of ethanolic Mangiferin in the treatment of irritation and inflammation on skin.

Methods: The study was conducted at Muhammad Institute of Medical and Allied Sciences, Multan, in October 2021. Four groups of 12 rabbits; of either gender, were structured in this way that each group contained 3 rabbits. Anti-inflammatory activity was measured against betamethasone and potency of anti-inflammatory agents was examined by inducing inflammation through four different inducers phenol, formalin, acetic acid and sand paper. To treat ear distilled water was used as control group.

Results: Ethanolic extract of seed of *Mangifera indica* showed excellent counter irritant activity when compared with betamethasone and control group water. Maximum tolerated dose and minimum tolerated dose was calculated at an interval of 2 hours in (75, 50 and 25) µg/ml pattern. These doses with same pattern showed anti-inflammatory activity ranging between 93.01% to 75.13% in phenol 92.01% to 79.31% in acetic acid 94.02% to 79.21% in formalin and 93.12% to 60.21% in sand paper respectively.

Conclusion: Ethanolic Mangiferin has the ability to lighten the effect of inflammation.

Keywords: Rabbits' ear, Counter-irritant, Ethanolic Mangiferin, Betamethasone.

Introduction

In case of tropic inflammation, increase of histamine level has been noted in the skin. Histamine is demonstrated as one of the major inflammatory mediators released by mast cells which are normally present in the connective tissue adjacent to blood vessels. Histamine is also found in blood basophils and platelets [1]. It is stored in mast cell granules and is released by degranulation in response to a variety of stimuli, including physical injury, such as trauma, cold or heat by unknown mechanisms or binding of antibodies to mast cell which underlies immediate hypersensitivity reactions. Histamine causes dilation of arterioles and increases the permeability of venules. Histamine is considered the principal mediator of the immediate transient phase of increased vascular permeability, producing inter endothelial gaps in postcapillary venules [2]. Its vasoactive effects are mediated mainly *via* binding to receptors, called H1 receptors, on microvascular endothelial cells. The anti-histamine drugs that are commonly used to treat some inflammatory reactions, such as allergies, are H1 receptor antagonists that bind to and block the receptor H1 [3].

Mangiferin (1,3,6,7-tetrahydroxyxanthone C2-B-D-glucoside) is a polyphenolic compound extracted from various parts of *M. indica* like roots, kernels, bark and flesh. Mangiferin shows various pharmacological activities like anti-diabetic, anti-oxidant, anti-tumor, anti-inflammatory and anti-microbial [4]. Mango seed is rich in Mangiferin component and showed excellent anti-inflammatory activity when compared with mango peel. Mangiferin has the ability to reduce Ig E and suppress anaphylactic reactions. Expectedly, Mangiferin reduce the activity of release of histamine by different ways

- Suppress mast cells, basophils and plasma cells to release histamine.
- Prevent the arachidonic acid to releases prostaglandin E which release higher concentration of histamine.
- Block the activity of H1 receptor by its antagonists and keeping them inactive in cytoplasm.

Ethanolic Mangiferin is used to treat tropical inflammation as well as due to ethanolic extract it is also beneficial as anti-septic drug. The current study was designed to evaluate the

effect of ethanolic extract of seed of *M. indica* on experimentally- induced irritation in rabbits [5].

Materials and Methods

The study was conducted at Muhammad Institute of Medical and Allied Sciences, Multan, in October 2021 and consisted of 12 rabbits of either gender. The rabbits were taken from Multan pet market, Punjab. The standard betamethasone was purchased from Ethical Laboratories (Pvt. Ltd.) Pakistan, while all chemicals were purchased from Solex Chemical (Pvt) Ltd-Multan. The 12 rabbits (with a mean weight of 1.5 ± 03 kg) were kept under two-hour observation for initially assessment of dermatological and allergic behavior before analyzing the counter irritant activity [6]. Animals were kept at Animal house of Muhammad Institute of Medical and Allied Sciences, Multan. They were kept in stainless steel cages and provided with good diet that's commercially available. The experiments were carried out according to the rules of Institute of Laboratory Animal Resources, Commission of life sciences, Natural research council and approved by The Animal Ethical Committee of Muhammad Institute of Medical and Allied Sciences, Multan., Pakistan [7].

The fresh seeds of the plant *M. indica* were collected from Khand Mangoes Farm, Multan. Seeds were authenticated by the cooperation of expert taxonomist at the Department of Botany, Baha Uddin Zakariya University, Multan and seed specimen were subjected for record. For the preparation of extract, first of all the fresh seeds of mango were subjected for shade drying [8]. All the adulterants and vegetative wastes were removed from vegetative material by manual picking before grinding into coarse powder with the help of special herbal grinder. The solution of coarse powder was prepared with ethanol and stored in the air tight jars. The filtrate was evaporated at room temperature under reduced pressure on rotary evaporator. The obtained extract was stored at -20°C in amber color airtight jars in laboratory refrigerators [9, 10].

The anti-inflammatory activity was determined by crude extracts against betamethasone (standard). The potency of anti-inflammatory agents was measured by inducing the inflammation in the experimental animal. Maximum tolerated dose and minimum tolerated dose was calculated at an interval of 2 hours in (75, 50 and 25) $\mu\text{g}/\text{mL}$ pattern. All rabbits were divided into four (G1, G2, G3 and G4) groups. Each group contained 3 rabbits. Irritation was induced in G1 by anti-clock wise application of sand paper on both ears [11, 12]. Induction of redness and erythema were observed and recorded. Standard betamethasone was applied on one ear while the other ear was treated with ethanolic Mangiferin extract of 75 $\mu\text{g}/\text{mL}$ dose in first rabbit, 50 $\mu\text{g}/\text{mL}$ in second rabbit and 25 $\mu\text{g}/\text{mL}$ in third rabbit. The time, dosage and degree of counter-irritancy were noted. Same method with remaining three inducers (phenol, formalin, acetic acid) was reexamined in G2, G3 and G4 groups. Decrease in redness and erythema was observed and recorded [13] (Figures 1 and 2).




Figure 1. Shows Preparation of extract.



Figure 2. Counter-irritant effect of normal, control, ethanolic *m indica* and betamethasone.

Results

All three doses showed moderate to excellent counter-irritant effect in dose-dependent manner on the rabbit's ear. The dose of 75 $\mu\text{g}/\text{mL}$ showed maximum anti-inflammatory activity in sandpaper irritation model by 93.12%. The maximum inflammatory healing response was examined by 75 $\mu\text{g}/\text{mL}$ dose (Table 1 and Figures 3-6).

Table 1. Shows counter-irritant activity.

Extract conc. mg/ml	Phenol irritation inhibition (%)	Acetic acid irritation inhibition (%)	Formalin irritation inhibition (%)	Sand paper irritation inhibition (%)
<i>M. indica</i> 25 $\mu\text{g}/\text{mL}$	75.13	79.31	79.21	60.21
	75.1	78.9	79.14	60.12
	74.21	78.01	78.99	59.99
<i>M. indica</i> 50 $\mu\text{g}/\text{mL}$	89.11	85.91	87.11	78.13
	89.09	85.19	87.03	78.31
	89.02	85.02	87	78.01
<i>M. indica</i> 75 $\mu\text{g}/\text{mL}$	93.21	92.12	94.29	93.29
	93.12	92.2	94.92	93.92
	93.01	92.01	94.02	93.12

Betamethasone	98.31	97.21	98.21	97.38
	98.13	97.12	98.12	97.31
	98.23	97.09	98.09	97.09




Figure 3. Acetic acid Irritation model inhibition. Note: ■ Betamethasone ■ M.indica 75µg/ml ■ M.indica 50µg/ml ■ M.indica 255µg/ml




Figure 5. Formalin irritation model inhibition. Note: ■ Betamethasone ■ M.indica 75µg/ml ■ M.indica 50µg/ml ■ M.indica 25µg/ml.




Figure 5. Sand Paper irritation model inhibition. Note: ■ Betamethasone ■ M.indica 75µg/ml ■ M.indica 50µg/ml ■ M.indica 25µg/ml.




Figure 6. Phenol irritation model inhibition%. Note: ■ Betamethasone ■ M.indica 75µg/ml ■ M.indica 50µg/ml ■ M.indica 255µg/ml

Maximum counter-irritancy time was noted in formalin inducer by 25µg/mL dose while minimum counter-irritancy time was observed in all inducers by 75µg/mL dose [14] (Table 2).

Table 2. Counter irritant activity of groups in minutes.

Groups	Phenol (min)	Acetic acid (min)	Formalin (min)	Sand paper (min)
M.indica 25 µg/mL	13	12	14	13
M.indica 50 µg/mL	10	9	11	10
M.indica 75 µg/mL	3	3	3	3
Betamethasone	2	2	1	1

Discussion

Different pharmacological aspects are responsible for anti-inflammatory activity. Mangiferin from mango seed has been identified which is demonstrated as to be involved in several activities. For instance, it prevents tumor formation, extends life span and is possible cure for diabetes, asthma and all inflammatory responses. It has advantage over the betamethasone as its multiple uses do not cause any harm rather than stinging and dryness as does betamethasone. Xanthoid structure of Mangiferin; rich source of polyhydroxy components, contributes to its free radical scavenging ability leading to multiple biological activities [15]. The presence of Mangiferin was bound to react with the irritated and inflamed cell membrane. Subsequently, the irritated and damaged superficial and deeper layers healed. Furthermore, anti-septic activity of ethanol is effective at killing microbes, bacteria and other microorganisms on the surface of damaged skin. Moreover, there is no evidence of adverse side effects of Mangiferin. Hitherto, Mangiferin could be a promising candidate to development of multiple drugs [16].

Conclusion

Ethanol extract of kernel of *M. indica* countered the effect of irritation in experimental animal also showed a significant

effect in terms of dose when compared with controls and standard betamethasone.

References

1. Akerman S, Williamson DJ, Kaube H, et al. The role of histamine in dural vessel dilation. *Brain Res.* 2002;956:96-102.
2. Asako H, Kurose I, Wolf R, et al. Role of H1 receptors and P-selectin in histamine-induced leukocyte rolling and adhesion in postcapillary venules. *J Clin Investig.* 1994;93:1508-1515.
3. Willis AP, Leffler CW. NO and prostanoids: Age dependence of hypercapnia and histamine-induced dilations of pig pial arterioles. *Am J Physiol Heart Circ Physiol.* 1999; 277:299-307.
4. Togias A. H1-receptors: Localization and role in airway physiology and in immune functions. *J Allergy Clin Immunol.* 2003;112:60-68.
5. Cowden JM, Zhang M, Dunford PJ, et al. The histamine H4 receptor mediates inflammation and pruritus in Th2-dependent dermal inflammation. *J Invest Dermatol.* 2010;130:1023-1033.
6. Khare P, Shanker K. Mangiferin: A review of sources and interventions for biological activities. *BioFactors.* 2016;42:504-514.
7. Stohs SJ, Swaroop A, Moriyama H, et al. A review on antioxidant, anti-inflammatory and gastroprotective abilities of mango (*Mangifera indica*) leaf extract and mangiferin. *J Nutr Health Sci.* 2018;5:302-303.
8. Rivera DG, Hernandez I, Merino N, et al. Extract (Vimang) and mangiferin reduce the airway inflammation and Th2 cytokines in murine model of allergic asthma. *J Pharm Pharmacol.* 2011;63:1336-1345.
9. Branco AC, Yoshikawa FS, Pietrobon AJ, et al. Role of histamine in modulating the immune response and inflammation. *Mediat Inflamm.* 2018.
10. Higgs GA, Flower RJ, Vane JR. A new approach to anti-inflammatory drugs. *Biochem Pharmacol.* 1979;28:1959-1961.
11. Saha S, Sadhukhan P, Sil PC. Mangiferin: A xanthonoid with multipotent anti-inflammatory potential. *BioFactors.* 2016;42:459-474.
12. Kumaran MS, Kaur I, Kumar B. Effect of topical calcipotriol betamethasone dipropionate and their combination in the treatment of localized vitiligo. *J Eur Acad Dermatol Venereol.* 2006;20:269-273.
13. Rivera DG, Balmaseda IH, Leon Aa, et al. Anti-allergic properties of *Mangifera indica* L. extract (Vimang) and contribution of its glucosylxanthone mangiferin. *J Pharm Pharmacol.* 2006;58:385-392.
14. Mei S, Ma H, Chen X. Anticancer and anti-inflammatory properties of mangiferin: A review of its molecular mechanisms. *Food Chem Toxicol.* 2021;149:111997.
15. Blech MF, Hartemann P, Paquin JL. Activity of non antiseptic soaps and ethanol for hand disinfection. 1985;181:496-512.
16. Telang M, Dhulap S, Mandhare A, et al. Therapeutic and cosmetic applications of mangiferin: A patent review. *Expert Opin Ther Pat.* 2013;23:1561-1580.

*Correspondence to

Dr. Ameen Abbas

Department of Biochemistry

Muhammad Institute of Medical and Allied Sciences

Multan

Pakistan

E-mail: rammyali01@gmail.com