

Evaluation of diagnostic potential of recombinant D-erythrose 1-phosphate dehydrogenase using indirect enzyme-linked immunosorbent assay for diagnosis of bovine brucellosis

W S A A Shell

Central Laboratory For Evaluation Of Veterinary Biologics, Egypt National Research Centre Dokki, Egypt Cairo University, Egypt, E-mail: tarikwaleedshell@hotmail.com

Abstract

Serological tests used for diagnosis of bovine brucellosis are usually depending on smooth lipopolysaccharides (S-LPS) as a diagnostic antigen which usually gives false positive reactions. So, our study aims to produce and evaluate a diagnostic kit for accurate diagnosis of bovine brucellosis differentiating between vaccinated and infected cattle and exclusion of false positive cases. Idea of this kit depends upon the fact that The EryC gene is absent in *Brucella abortus* S19 only but it is present and functional in all other *Brucella* strains and isolates so according to these facts, the use of ELISA kit coated with single subunit (recombinant) EryC protein may be useful, rather than S-LPS, as an alternative diagnostic antigen in diagnosis of bovine brucellosis and differentiation between S19 vaccinated and *Brucella* infected cattle. The present study evaluated antibody responses of brucellosis infected and S19 vaccinated cattle to purified recombinant EryC protein in an indirect enzyme-linked immunosorbent assay (I-ELISA). Cattle sera were screened using Rose Bengal Plate test (RBPT). 114 samples of naturally infected cattle (Rose Bengal test positive), 78 sera from S19 vaccinated cattle and 25 sera samples from *Brucella* free cattle were used in this study. I-ELISA using S-LPS and periplasmic proteins as a coating antigen were used as a gold standard test. The results revealed that in case of sera of naturally infected cattle, sero-positivity was 94.7%, 100%, 100% and 100% with EryC-ELISA, LPS-ELISA, periplasmic-ELISA and Rose Bengal test respectively. Where in case of sera of S19 vaccinated cattle, all samples were negative when tested with EryC-ELISA while in case of LPS-ELISA, periplasmic-ELISA and rose Bengal test, sero-positivity was 92.3%, 84.6% and 100% respectively. It could be concluded that the EryC protein could be used in serological tests for diagnosis of bovine brucellosis and differentiation between infected and *Brucella abortus* S19-vaccinated cattle but more studies are needed to be done on large cattle populations accompanied with bacteriological isolation to detect the sensitivity and specificity of this protein as a diagnostic antigen and also for validate this test. Brucellosis is a major bacterial zoonosis of global importance affecting a wide range of different mammals including cattle, sheep, goats, swine, rodents, marine mammals and man worldwide.

In food animals, the disease primarily affects the reproductive system with concomitant loss in fertility and productivity of affected animals. In man, infection is characterized by recurrent febrile episodes that lead to the description of this disease as undulant fever (economic and public health importance). The severity of this disease and lack of vaccines suitable for use in man has led to the investigation of *Brucella* as agents for bioterrorism (Bio-risk importance). Vaccines to be used for human are not yet available, and so eradication of human brucellosis largely depend on the eradication of the disease in animals

This work is partly presented at

21st European Biotechnology Congress on October 11-12, 2018, held in Moscow, Russia.

