

Length of Hospital Stay in Critical Care Units.

Maria Brenner *

School of Nursing and Midwifery, Trinity College Dublin, Ireland

Accepted on 14 April, 2020

Editorial Note

I am pleased to mention that during the year 2019, all issues of volume 1 were published online well within the time and the print issues were also brought out and dispatched within 30 days of publishing the issue online. AAICCN during this year also brought out a supplement to Jan-Feb 2019 issue. Critical Care Nursing is an important part of Nursing with a focus of complete care of the patients. Journal of Intensive and Critical Care Nursing is a newly launched Open access, Peer-reviewed scientific journal which will be dedicated to promote the scientific community dealing with utmost care of critically ill patients in the area of Critical Care Nursing, Intensive Care Medicine, Emergency and Critical Care.

Ahmedul Abdul Alhari explained the need for well-trained health care providers is paramount. Undoubtedly, in mass casualty events, the scene of a real incident is not an appropriate place to learn. Thus, clinical care for mass casualty incidents (MCIs) victims and scene management procedures must be preplanned and practiced preincident to be able to achieve the optimum outcomes. Hence, training must be a priority in any planning and response strategy. Around the globe exercises and simulations are the most accepted training methodologies to train emergency responders for mass casualty incidents [1].

Alfateh Sayed Mohammed Noor explains the evidence of acceptable performance of the ICU with regards to mortality rate. We still need to improve our average LOS for all patients. There is improvement in 2018 performance compared to 2017. Insignificant changes in expenditure between the two comparison periods [2].

References

1. Alharbi AA. Educational approaches to mass casualty incidents. *J Intensive Crit Care Nurs.* 2019; 2(1):1-3.
2. Mohammed Noor AS, Al-Harthy AM, Ramadan, et al. Evaluation of performance in intensive care unit. *J Intensive Crit Care Nurs.* 2019; 2(1): 1-6.

*Correspondence to

Dr. Maria Brenner

School of Nursing and Midwifery

Trinity College Dublin

Ireland

E-mail: aphelan1@tcd.ie