



Liver toxicity induced by an isoniazid prophylaxis therapy in an hiv infected single women in namakkal: a case report

P.REVINSELVAN

Naval AIDS Research Centre, Namakkal, India

Abstract:

Chemoprophylaxis will be initiated among HIV patients to prevent co-infection with tuberculosis. Many studies reveal that Anti-Retroviral Therapy (ART) will itself causes its side effects and toxicity. In immune-compromised individuals, there are chances of decreased metabolic and cellular functions. When an Isoniazid Prophylaxis Therapy (IPT) is introduced along with ART in HIV patients some individuals are found to have Drug-Induced Liver Injury (DILI). we report a case of liver toxicity induced by IPT in an ART undergoing HIV patient.

Biography:

P.Revinselvan, I am working in HIV Cure research with my Prakasine Complexes. Now this complexes working better than ART, very good remedy for HIV patients. I have started this research in 1993 with Tamil Traditional Medicine (TTM). Now, I have applied TTM, bio and nano technologies to develop this new molecules. This molecules not only give functional and permanent cure for HIV people, it seems this molecules act as cell proliferators in vivo. This can be useful for cancer and diabetic patients too..

Recent Publications:

- 1. Thiyagarajan. Gunapadam (Materia Medica). Department of Indian Medicine and
- 2. Homeopathy, Chennai; 1992: 167-199.



- 3. 2. Agasthiyar. Agasthiyar paripooranam 400 (Agasthiyar total cure 400 formulations).
- Rathina Nayakar & sons, Thirumagal vilakku press, Chennai; 1964: 40.
- 5. 3.Matchamuni. Matchmuni perunool 700 (Matchmuni big therapeutic book 700
- 6. formulations).CCRAS publications, Chennai;1985: 113-114.

Webinar on HIV, AIDS Infectious Disease & Healthcare | November 23, 2020 | Dubai, UAE

Citation: P.Revinselvan, Liver Toxicity Induced By An Isoniazid Prophylaxis Therapy In An Hiv Infected Single Women In Namakkal: A Case Report

J Bacteriol Infec Dis 2020 Volume: and Issue: S(10)