# **On-farm** potential of improved sorghum varieties in dry lowland areas of Ethiopia.

# Fitsum Miruts, Belay Roba<sup>\*</sup>, Derara Sori

Department of Agricultural Extension and Communication Process, Ethiopian Institute of Agricultural Research, Melkassa Agricultural Research Centre, Melkassa, Ethiopia

## Abstract

Feeding the world's growing population has become a global issue due to climate change, which has resulted in low agricultural productivity. Although an attempt has been made to feed this growing population by introducing improved agricultural technologies that boost productivity, the sustainability of these introduced technologies is not promising as farmers insist to use traditional way of production. The study was conducted in the districts of Gololcha and Shanana Kolu to investigate the on-farm potentials of Melkam, Tilahun, and Argiti varieties for further scaling out. Melkam had a higher grain yield (3628 kg/ha) than Argiti (3234 kg/ha), Tilahun (3050 kg/ha), and the district average sorghum productivity (2320 kg/ha). The mean yield comparison results of yield performances revealed a significant difference at (p-0.05). The higher gross margin (97,432 Ethiopian birr per hectare) was received from Melkam followed by Argiti (84,824 Ethiopian birr per hectare). Melkam variety had a marginal beneficial advantage of 57,306 over the local, Argiti (44,698) and Tilahun (38,810 Ethiopian birr) per hectare. The improved varieties were also preferred by farmers because of their high yield, early maturity, and drought-tolerant traits. The mean weight score of the selected traits revealed that grain yield and injera-making quality were the top priorities for women farmers, while grain yield and earliness were the top priorities for men farmers. The preference of the two groups differs, indicating that gender matters in the scaling out of improved varieties introduced to the farming community. As a result, it is recommended that Melkam varieties be adopted and scaled out with the recommended production practices in the study areas and areas with similar agroecology to improve food availability and farmers' income.

Keywords: On-farm Yield, Cost-benefit, Demonstration, Improved varieties.

## Introduction

Sorghum (Sorghum bicolour (L.) Moench) is one of the major cereal crops produced in the world [1]. It is a major food security crop in sub-Saharan Africa, supporting some 300 million people. It is a relatively drought-tolerant crop adapted to grow under harsh production conditions. In Ethiopia, sorghum is ranking third after teff and maize in the total production area. It ranks fourth after maize, teff, and wheat in grain production. It is grown in almost all regions, covering a total area of 1.6 million ha. It is grown in three major agroecologies. In the dry lowland environment, it is the major crop cultivated in more than 60% of the area [2].

The crop is preferred for the preparation of various traditional foods (Injera and porridge) and beverages (Tella and Borde). The stalk is an important source of animal feed. In addition, the biomass is used for construction, fuel wood, and fencing. In the study areas, sorghum takes first rank in production. It is largely produced by smallholder farmers. Sorghum cultivars cultivated in the study areas are characterized by their long height (3 meters and above) and late maturing (more than 8 months) with low productivity. The red-coloured sorghum (grain) is the dominant one.

Ethiopian Institute of Agricultural Research (EIAR) developed and released more than 25 drought-tolerant, yielder, and early maturing sorghum varieties suitable for dry lowland areas with improved production management practices to improve the productivity level [3]. However, the cultivation of landraces/ local cultivars persists among smallholder farmers across the country. The number of farmers growing the improved sorghum varieties is 28%, which is very low [4]. The low level of improved sorghum varieties cultivation is attributed to the low access to improved seeds or information and availability of farmer-preferred varieties [5-7].

\*Correspondence to: Belay Roba, Department of Agricultural Extension and Communication Process, Ethiopian Institute of Agricultural Research, Melkassa Agricultural Research Centre, Melkassa, Ethiopia; E-mail: robabelay1984@gmail.com

Received: 20-Jan-2024, Manuscript No. AAASCB-24-125491; Editor assigned: 23-Jan-2024, AAASCB-24-125491 (PQ); Reviewed: 06-Feb-2024, QC No. AAASCB-24-125491; Revised: 11-Feb-2025, Manuscript No. AAASCB-24-125491 (R); Published: 18-Feb-2025, DOI: 10.35841/aaascb-9.1.277

To boost sorghum productivity, increasing the availability of improved varieties through demonstration, promotion, and scaling out will assist the farming community for further adoption. To accomplish this, many varieties have been demonstrated in certain areas, and work to address the wider farming community is currently underway. The goal of this study is to find a promising variety that meets farmers' demand for further expansion.

## **Materials and Methods**

#### Description of the study area

The experiments have been conducted in the districts of Gololcha and Shanan Kolu, the districts, which represent lowland and midland agroecologies, are situated in northeastern Ethiopia. Mixed crop and livestock farming is the main form of agricultural production in both districts. The main crops grown are sorghum, coffee, maize, teff, and khat. The main rainy season for the district was from May to September, with 400–1112 mm, and was followed by a prolonged dry period of seven months. According to GDOoANR and SKOoANR, the temperature ranges from 26 to 40 degrees Celsius. The main cultivars that farmers cultivate are the locally popular cultivars known as Arkabas (matures in 5–6 months) and Alewalem (matures in 9 months) [8,9].

Different stakeholders introduce improved and early matured sorghum varieties like Gobiye, Abshir, Teshale, and Dekeba as local cultivars in use are late mature. Due to the farmers' preference for certain traits and the lack of strong extension services, the varieties were not scaled up to the larger sorghum producer farmers. Farmers insist on using their local cultivars at large (Figure 1).



#### Site and farmers' selection

Based on their sorghum production volume, the Gololcha and Shanan Kolu districts were specifically selected from the Arsi Zone. The Arsi Zone agricultural and natural resource office was consulted before making the choice. Ten kebeles from the Gololcha district and four kebeles from Shanan Kolu were the targets. Additionally, the kebeles were chosen at random from the study area. Farmers who are interested in the technology, willing to manage the experimental field, willing to allocate land for the experiment, and willing to share the results with other farmers are just a few of the selection criteria for experiment host farmers.

The improved sorghum management practices were introduced, and practical training was given, before the start of the experiment. In total, 93 host farmers' fields (fifty-seven in 2020 and thirty-six in 2021) were established on 23.25 hectares where each farmer allocated 0.25 hectares on average for three varieties. Farmers who established the experiments were considered as a replication. The replication of the experiment in two consecutive years was to increase the reliability of the data and conclusions to be drawn.

#### Materials and management used

Three released sorghum varieties were evaluated in 2020 and 2021 during the main cropping seasons in both districts. Improved sorghum varieties; namely Tilahun, Argiti, and Melkam were used for the experiment. These varieties are adapted to dry lowland areas. The varieties were planted side-by-side to compare and evaluate their performance. During planting, the seeds were manually drilled at a seed rate of 10 kg/ha.

Each variety was planted on a plot size of  $400 \text{ m}^2$  (20 m long by 20 m width), with an intra and inter-row spacing of 20 cm and 70 cm, respectively. Land preparation was done according to the recommended practice. Fertilizer, 100 kg/ha of the NPS was applied at the time of planting and 50 kg/ha of Urea was applied in the form of split application; half of it was at planting and the rest was top-dressed at the knee-height stage (Table 1).

Figure 1. Study area map.

 Table 1. Description of improved sorghum varieties used for the experiment.

Variety	Flowering date	Plant height (cm)	Productivity	Maturity date (days)	Туреѕ
Melkam	76-82	126-163	3500-5800	118	Semi-compacted head and white grain color

Argiti	79	200	3750-6000	125	Compacted head and white grain color
Tilahun	78	190	3400-6000	120	Semi-compacted head and white grain color

## Quantification of yield gaps

The researchers recorded qualitative and quantitative data during scheduled farm visits and follow-ups. Data were collected using a data sheet. Yield gaps were calculated using the definition and concepts provided by Lobell, et al. [10]. These are research station yields (Maximum possible yield with improved management practices, under controlled field conditions, taken from the crop variety registry book MoAL, for this case), on-farm demonstration yields (obtained yields with improved management practices under on-farm situations) and district-level average yields (yield obtained under farmers condition with traditional management practices) [11].

The yield gaps were divided into two groups. The yield gap between the research station yields and on-farm demonstration field yield. The second gap is the demonstration yield and district-level average yields. The difference between the research station yields and the district-level average yield is the total yield gap.

## Data collection and data analysis

Data were collected on grain yield, farmers preferred treats, and cost of production. For simplicity, the yield obtained from the plot was converted to a kg/ha. The data were analyzed using descriptive statistics, preference ranking, yield gap, and profitability (gross margin, net return, total return, and the benefit-cost ratio). In profitability analysis land was taken as a fixed asset in the assessment of its opportunity cost. Costs of land preparation, seed, fertilizer, and hired labor were

*Table 2.* Demographic characteristics of farmers (N=93).

calculated based on the existing market price of inputs during the experimental period. The total cost was estimated by adding variable costs and fixed costs. To estimate the cost of sorghum cultivation, BCR methods were used [12].

## **Results and Discussion**

#### Description of sorghum producer households

Of the total demonstration host farmers, 51 are from the Shanan Kolu district while 42 are from the Gololcha district. The number of female demonstration host farmers was lower than male farmers (Table 2). Male farmers represented 81%, whereas female farmers were at 19%. A majority of sorghum farmers have no formal education (86%) and only 12% attended a primary level of education. More than half of the sorghum producers (76%) are aged between 31 to 50 years with an average year of 38. The sorghum farmers' family size ranged from 1 to 10, with an average size of less than two. The majority, 60% had a family size of 1-5. The farmers' experience in sorghum production ranges from 5 to 38 years, with an average farming experience of 18 years. The total farm size owned by the farmers ranged from 0.25 ha to 11 ha. Most farmers had a farm size of 1-3 ha (45%) followed by less than 1 ha (43%). This shows that the study area is dominated by smallholder farmers. Most farmers (73.1%) grow sorghum on a plot size of less than 0.5 ha.

Variables		Number of farmers	Percent	Mean	SD
Education level	No formal education	80	86	-	-
	Primary education	11	11.8		
	Secondary	2	2.2		
Age (years)	≤ 30	13	14	38	6.88
	31-50	76	81.7		
	51-70	4	4.3		
Sex	Male	75	80.6	-	-
	Female	18	19.4		
Experience (years)	≤ 10	20	21.5	18	6.89
	11-20	49	52.7		
	21-30	19	20.4		
	>30	5	5.4		
Farm size (ha)	<1	43	46.2	1.35	1.47

*Citation:* Miruts F, Roba B, Sori D. On-farm potential of improved sorghum varieties in dry lowland areas of Ethiopia. J Agric Sci Bot. 2025;9(1):277.

	1-3	45	48.4		
	>3	5	5.4		
Area allocated to sorghum	≤ 0.5	68	73.1	0.47	0.26
(na)	0.5-1	23	24.7		
	>1	2	2.2		
Family size	1-5	56	60.2	1.35	1.47

#### **On-farm yield performance of sorghum varieties**

The mean grain yield of the improved sorghum varieties demonstrated was 3050 kg/ha, 3234 kg/ha, and 3628 kg/ha for Tilahun, Argiti, and Melkam varieties respectively. The highest mean yield was recorded by the Melkam variety in 3682 kg/ha in Shanan Kola district. This means that the Melkam variety has a grain yield advantage over Tilahun and Argiti with 15% and 11% respectively. Comparing the improved varieties with the local control (district average yield), a 36% grain yield advantage was recorded by the Melkam variety, the highest mean grain yield gap.

The average yield of Argiti and Tilahun was also more than the district average yield which was 2320 kg/ha. According to EIAR, the average sorghum productivity at farmers' fields is about 2800 kg/ha, while it is 3000 kg/ha to 5000 kg/ha at experimental plots [13]. The mean yield obtained from the improved varieties in this study is therefore superior and promising to smallholder farmers in the study area and similar agroecology. The high yield potential of the improved varieties therefore will provide to improve food security and income of sorghum-producing households in the study area. The yield difference observed was due to the suitability of the environment for the Melkam variety in the study area.

The result in Figure 2 is evident that the performance of the Melkam variety was found better than Tilahun, Argiti varieties,

Table 3. Mean yield gap of sorghum varieties (kg/ha).								
Variety	Experimental station yield	District averaged yield	On-farm demonstration Yield	Technology gap	Total yield gap			
Tilahun	4700	1840	3017.88	1177.88	2,860.00			
Argiti	4800		3378.64	1538.64	2,960.00			
Melka	5100		3562.79	1722.79	3,260.00			
Tilahun	4700	2800	3077	277	1,900.00			
Argiti	4800		3115.31	315.31	2,000.00			
Melkam	5100		3682.06	882.06	2,300.00			
Tilahun	4700	2320	3050.3	730.3	2,380.00			
Argiti	ti 4800	3234.24	914.24	2,480.00				
Melkam	5100		3628.19	1308.19	2,780.00			
	ld gap of sorghum Variety Tilahun Argiti Melka Tilahun Argiti Melkam Tilahun Argiti Melkam	Id gap of sorghum varieties (kg/ha).VarietyExperimental station yieldTilahun4700Argiti4800Melka5100Tilahun4700Argiti4800Melkam5100Tilahun4800Melkam5100Tilahun4700Melkam5100Tilahun4700Melkam5100Tilahun4800Melkam5100	VarietyExperimental station yieldDistrict averaged yieldTilahun47001840Argiti4800100Melka51002800Tilahun47002800Argiti5100100Melkam51002800Melkam51002800Melkam51002800Tilahun47002800Melkam51002900Melkam51002000Melkam51002000Melkam51002000Melkam51002000Melkam51002000	VarietyExperimental station yieldDistrict averaged yieldOn-farm demonstration YieldTilahun470018403017.88Argiti48003378.64Melka51003562.79Tilahun470028003077Argiti48003115.31Melkam51003682.06Tilahun47003230Argiti48003050.3Tilahun51003234.24Melkam51003628.19	VarietyExperimental station yieldDistrict averaged yieldOn-farm demonstration YieldTechnology gapTilahun470018403017.881177.88Argiti48002003378.641538.64Melka51007122.791722.79Tilahun470028003077277Argiti48003115.31315.31Melkam51003203682.06882.06Tilahun470023203050.3730.3Melkam51003124.24914.24Melkam51003024.24914.24			

1 1

and the local cultivars. This result clearly showed that new varieties have a higher grain yield. The result conforms with that of EIAR and Girma, et al. (Table 3) [14,15].



Figure 2. On-farm yield performance of sorghum varieties.

The same letters in the same group are not significantly different at 5% using the Dunkan test.

#### Gender-based preference analysis

A group of women and men farmers were formed from the host farmers as the preference may vary between them. Grain

yield, good injera-making quality, earliness, higher market price, white grain color, large grain size, grain thresh ability, Striga resistance, Stalk sweetness, diseases and pest tolerance,

Citation: Miruts F, Roba B, Sori D. On-farm potential of improved sorghum varieties in dry lowland areas of Ethiopia. J Agric Sci Bot. 2025;9(1):277.

large biomass, tallest plant height, and birds attack tolerant are the criteria taken into account.

Accordingly, the mean weight score of the selected traits result showed that in the case of women farmers grain yield and injera-making quality ranked as their top priority while grain yield and earliness for the men farmers. Although both genders have the same demand for grain yield, the preference between the two groups is different showing that genders matter for the

 Table 4. Farmers prioritized traits and mean score.

scaling out of improved variety introduced to the farming community.

The analysis shows that the Melkam variety was preferred for most of the characteristics except large grain size, stalk sweetness, large biomass, and plant height. Argiti variety was ranked as the second most preferred variety (Table 4).

Farmers prioritized trait	Women score	Rank	Male score	Rank	Mean score	Rank
Grain yield	29	1	30	1	29.5	1
Good injera-making quality	29	1	25	2	27	2
Earliness	16	6	30	1	23	3
Highest market price	23	2	19	5	21	4
Grain colour (white)	19	5	22	4	20.5	5
Grain size (large)	21	3	19	5	20	6
Grain threshes ability	19	5	16	7	17.5	7
Striga resistance	8	10	23	3	15.5	8
Stalk sweetness	9	8	18	6	13.5	9
Diseases and pest- tolerant	16	6	10	8	13	10
Biomass (large)	2	11	18	6	10	11

#### Sorghum production costs

The costs of sorghum production are categorized under materials (fertilizer and seed) and operations incurred by farmers. From the total variable costs, more than 80% of the expenditure is operation costs. The costs of operation include, labor for ploughing, planting, weeding, harvesting, threshing and transport, and post-harvest handling formed the largest cost item and accounted for 72% of the total variable costs.

The average variable cost of sorghum production was 18,664 ETB per hectare for improved varieties and 17,874 ETB per

hectare for local cultivars. The cost difference between the improved and the local cultivar was due to the price difference of a seed price and the fertilizer amount used. The lion's share of the cost was incurred for ploughing (24%), followed by threshing, and transport costs (20%). As the result shows all costs of sorghum production are the same except costs of seed and fertilizer (Table 5).

Table 5. The operational cost of sorghum production.

Type of expenses	Average cost (ETB/ha) T		Total cost (%)		
	Local	Improved	Local	Improved	
Cost of input	2610	3400	14.6	18.2	
Fertilizer	2,250	3,000	12.6	16.1	
Seed	360	400	2	2.1	
Operation cost	15264	15,264	85.4	81.8	
Ploughing (three times)	4,466	4,466	25	23.9	
Planting and fertilizer application	2,200	2,200	12.3	11.8	
Total wedding	3,332	3,332	18.6	17.9	
Harvesting	1,666	1,666	9.3	8.9	

*Citation:* Miruts F, Roba B, Sori D. On-farm potential of improved sorghum varieties in dry lowland areas of Ethiopia. J Agric Sci Bot. 2025;9(1):277.

Threshing and transport	3,600	3,600	20.1	19.3
Total production cost (variable)	17,874	18,664	100	100

#### Cost benefit analysis of sorghum production

The total costs for sorghum production vary from farmer to farmer and location to location. The costs and benefits for each variety were calculated by considering variable costs. The net benefit from Melkam, Tilahun, and Argiti varieties production was 97,432, 78,936, and 84,824 ETB per hectare respectively. However, the net benefit for the local sorghum cultivars was 40,126 ETB per hectare. This is far lower than the improved varieties of sorghum. Melkam variety had the highest benefit-to-cost ratio. For everyone ETB invested in the production of sorghum, the farmers earned a higher net income from the

Melkam variety. Melkam has a higher benefit-cost ratio (4.4) followed by Argiti (3.9) and Tilahun (2.2).

The benefit-cost ratio for local cultivars (2.2) is lower than the improved varieties. The profitability of sorghum production relates directly to the productivity and grain price. Thus, farmers will be at a high level of profit if they cultivate the Melkam variety using associated production practices followed by the Argiti variety in the study areas (Table 6).

Table 6. Cost and Benefit analysis of the demonstrated varieties per hectare.

Operations	Local	Melkam	Tilahun	Argiti
Total variable cost	17,874	18,664	18,664	18,664
Fixed cost (land cost/ha)	8,000	8,000	8,000	8,000
Total cost (fixed and variable cost)	25,874	26,664	26,664	26,664
Grain yield (kg/ha)	2,320	3,628	3,050	3,234
Grain price (ETB/kg)	25	32	32	32
Gross income	58,000	116,096	97,600	103,488
Net benefit/total revenue	40,126	97,432	78,936	84,824
Total net benefit (including fixed)	32,126	89,432	70,936	76,824
Marginal benefit (ETB)	Base	57,306	38,810	44,698
BCR	2.2	4.4	3.7	3.9

#### Marginal analysis

Farmer's net benefit is raised by cultivating the improved varieties. Based on the finding, the highest marginal yield was earned by Melkam and followed by Argiti. The marginal yield of Melkam, Argiti, and Tilahun was 73%, 57% and 49% respectively. This shows that Melkam was the first chic and Argiti was the second one. Since the marginal yield of the Tilahun variety was under 50%, it is not recommended to be cultivated by the farmer. The improved varieties gave an

additional one kg of grain yield with additional costs of 0.6, 0.9, and 1.1 ETB for Melkam, Argiti, and Tilahun respectively. In the rule of marginal cost analysis, the lowest cost that increases a unit yield is the best option. Accordingly, the Melkam variety is the best choice for sorghum producer farmers (Table 7).

 Table 7. Marginal analysis for improved varieties compared to local cultivars.

Treatment	Variable cost	Net benefit	Productivity (kg)	Marginal cost	Marginal benefit	Marginal rate of return
Melkam	18,644	97,432	3628.19	0.6	57,306	72.5
Argiti	18,664	84,824	3234.24	0.9	44,698	56.6
Tilahun	18,664	78,936	3050.3	1.1	38,810	49.14
Local	17,874	40,126	2320			

## Conclusion

Sorghum is an ideal crop for improving farmers' food security and resilience in marginal areas such as Gololcha and Shanan Kolu. As a result, scaling out the improved varieties, particularly the Melkam variety, is the most effective way to increase productivity. The Melkam variety was found to be the most productive when compared to local cultivars and improved varieties such as Argiti and Tilahun. As compared to farmer productivity, its high yield potential can increase up to 1308 kg/ha. Thus, adopting the Melkam variety along with recommended production practices will increase food availability and farmers' income. It is recommended that district agricultural offices scale up the Melkam variety with the full production package to farmers in the study area and similar agroecologies.

## Acknowledgement

The acknowledgment will go for Gololcha and Shanan Kolu agricultural office staff members and farmers in establishing the experiments and frequent follow-ups. The financial and resource support of the Melkassa agricultural research center and CultAF-II project for the implementation of the activity is well recognized.

## References

- 1. CSA (Central Statistical Agency). Reports on Area and Production of Crops. (Privater Peasant Holdings. Meher Season, Volume I. Statistical Bulletin 590. Addis Ababa, Ethiopia). 2021.
- Kinfe H, Tesfaye A. Yield performance and adoption of released sorghum varieties in Ethiopia. Edelweiss Appl Sci Technol. 2018;2(1):46-55.
- 3. Tadesse T, Seyoum A, Nega A, et al. Major achievements, challenges and prospects of sorghum and millet research and development. MERCI project TERRA project. 2019.

- Wubeneh NG, Sanders JH. Farm-level adoption of sorghum technologies in Tigray, Ethiopia. Agric Syst. 2006;91(1-2): 122-34.
- Teferi T. Adoption of improved sorghum varieties and farmers' varietal trait preference in kobo district, northwolo zone, Ethiopia. Haramaya universuty, Dire Dawa, Ethiopia. 2013.
- 6. Beshir B, Sime M. Understanding Farmers' Improved Sorghum Variety Selection Criteria. 2013.
- 7. GDOoANR. Report on agricultural production and performance evaluation. 2021.
- 8. SKOoANR. Report on agricultural production and performance evaluation. 2021.
- 9. Lobell DB, Cassman KG, Field CB. Crop yield gaps: their importance, magnitudes, and causes. Annu Rev Environ Resour. 2009;34(1):179-204.
- 10. MoAL. Crop variety registry book. Ministry of agriculture and livestock. Ethiopia. 2016.
- 11. de Rus G. Introduction to cost-benefit analysis: looking for reasonable shortcuts. Edward Elgar Publishing; 2021.
- 12. EIAR. Cereal Crop Production and Management Manual. 2016.
- 13. EIAR. Sorghum production and utilization manual. Amharic version. 2022.
- 14. Mengistu G, Shimelis H, Laing M, et al. Assessment of farmers' perceptions of production constraints, and their trait preferences of sorghum in western Ethiopia: implications for anthracnose resistance breeding. Acta Agric Scand B Soil Plant Sci. 2019;69(3):241-9.
- 15. Brown TE. Standards for MRR reference service. Arch Inform Newsletter. 1988;2(2):34-5.