



Prevalence of Escherichia Coli Among Urinary Tract Infections Patients (aged 0-70 years) Attending Hadejia General Hospital, Jigawa State-Nigeria

Hassan Muhammad

Department of Science Laboratory Technology, School of Science, Binyaminu Usman Polytechnic, Hadejia, Jigawa State-Nigeria.

Abstract:

Urinary Tract Infections (UTIs) are one of the most common nosocomial infections after Respiratory Tract Infections (RTIs) worldwide. Nowadays it represents one of the most common diseases encountered in medical practice affecting people of all ages from the neonate to the geriatric age group. The aim of this research work was to find the prevalence of Escherichia Coli among Urinary Tract Infections patients (0-70 years) attending Hadejia General Hospital, Jigawa State-Nigeria between August and September, 2019. A total of One hundred (100) urine samples comprising of 42(42%) in female and 58(58%) in male were obtained from the patients with age ranged from 0-70 years and analyzed. Samples collected were cultured and incubated at 37C for 24 hours. Isolation and Identification of bacteria was done by standard laboratory procedures. Twenty-six (26) samples yielded significance growth for E. Coli out of which 15 were from females and 11 were from males, based on cultural, microscopic and biochemical characteristics. The highest prevalence occurred within the age ranged 31-40 with females taking the lead. This means that it's necessary to practice good personnel hygiene through hand washing and used of hand sanitizer to reduce the incidence of microbial transmission.

Biography:

Hassan Muhammad Department of Science Laboratory Technology, School of Science, Binyaminu Usman Polytechnic, Hadejia, Jigawa State-Nigeria.

Publication of speakers:

1. Geng, Xiang & Xu, Jian & Ma, Xin & Wang, Xu & Huang, Jiazhang & Zhang, Chao & Wang, Chen & Muhammad, Hassan. (2014). Anatomy of the Sural Nerve With an Emphasis on the Incision for Medial Displacement Calcaneal Osteotomy. The Journal



of foot and ankle surgery : official publication of the American College of Foot and Ankle Surgeons. 54. 10.1053/j.jfas.2014.07.008.

2. Xu, Jian & Geng, Xiang & Muhammad, Hassan & Wang, Xu & Huang, Jia-Zhang & Zhang, Chao & Ma, Xin. (2014). Comparison of the incisions for the open surgical treatment of gluteal muscle contracture. Journal of pediatric orthopedics. Part B. 23.
3. Xu, Jian & Muhammad, Hassan & Wang, Xu & Ma, Xin. (2014). Botulinum Toxin Type A Injection Combined With Cast Immobilization for Treating Recurrent Peroneal Spastic Flatfoot Without Bone Coalitions: A Case Report and Review of the Literature. The Journal of foot and ankle surgery : official publication of the American College of Foot and Ankle Surgeons. 54. 10.1053/j.jfas.2014.03.008.
4. Xu, Jian & Zhang, Yijun & Muhammad, Hassan & Wang, Xu & Huang, Jiazhang & Zhang, Chao & Geng, Xiang & Ma, Xin. (2015). In vivo three-dimensional analysis of hindfoot kinematics in stage II PTTD flatfoot. Journal of orthopaedic science : official journal of the Japanese Orthopaedic Association. 20. 10.1007/s00776-015-0698-4.

11th International Conference on Clinical Microbiology and Infectious Diseases

Citation: Hassan Muhammad, Prevalence of Escherichia Coli Among Urinary Tract Infections Patients (aged 0-70 years) Attending Hadejia General Hospital, Jigawa State-Nigeria.; Euro Clinical Microbiology 2020; June 26, 2020; London, UK