

## Radiological assessment of difficult airway in thyroid malignancy: A retrospective case study

Shivaam Kesarwaani<sup>1\*</sup>, Vedula Padmini Saha<sup>2</sup>, Animesh Ghosh<sup>3</sup>, M K Nandhini<sup>4</sup>, Somnath Saha<sup>5</sup>

<sup>1</sup>Department of ENT and Head-Neck Surgery, Calcutta National Medical College, Kolkata, West Bengal, India

<sup>2</sup>Department of Plastic Surgery, R.G. Kar Medical College, Kolkata, West Bengal, India

<sup>3</sup>Department of Medical Officer & Clinical Tutor, Calcutta National Medical College, Kolkata, West Bengal, India

<sup>4</sup>Department of Junior Resident, Calcutta National Medical College, Kolkata, West Bengal, India

<sup>5</sup>Department of ENT and Head-Neck Surgery, Calcutta National Medical College, Kolkata, West Bengal, India

### Abstract

**Aims and Objective:** To propose the importance of doing a radiological assessment for any thyroid malignancy before opting for intervention, in-order to assess the severity of the disease and also to proceed with further better management.

**Materials and Methods:** Study includes a total number of 4 advanced thyroid malignancy, who presented to emergency department of ENT Head-Neck Surgery, Calcutta National Medical College. Patients were stabilized in emergency department & underwent ENT & systemic examinations. All cases were subjected to radiological investigations like plain radiography and Contrast enhanced CT scan of neck to better understand & study the disease extension. The parameters assessed were:

1. The extent of the disease.
2. Tracheal involvement.
3. Tracheal compression.
4. Other viscera involved.
5. Pretracheal space involvement.

**Result:** Tracheal compression was noted in 100% cases and tracheal involvement in 50% of total cases. Tracheal narrowing of about 70% was noticed in all cases, although the voice of the patient was absolutely normal in all the cases. Vascular involvement was seen in 75% cases and in the form of 360 degree was seen in 25%. Retrosternal extension was noted in 100% cases and pretracheal space involvement in 75% cases.

**Conclusion:** Thyroid malignancy especially anaplastic carcinoma can be very aggressive, where CT assessment is of utmost importance in view of management. It helps in assessing the risk for any intervention. It gives a cue if patient can be salvaged.

**Keywords:** Thyroid malignancy, Radiological assessment, Difficult airway, Tracheal compression, Retrosternal extension, CT imaging.

*Accepted on 16 April, 2021*

### Introduction

Thyroid malignancy with limited disease can be managed primarily by surgery but diseases with loco regional and distant metastasis with complications is difficult to manage. Airway management in large and retro-sternal thyroid malignancy with tracheal compression is often met with challenges, where radiological assessment plays a vital role in emergent airway control. Computed tomography (CT) and magnetic resonance imaging (MRI) can play an important role in preoperative and post-treatment assessment of thyroid malignancy. The radiologist as well as the treating surgeon should be aware of the pathological behavior of thyroid carcinoma, extent of disease involvement, airway patency and the characteristic imaging appearance of the primary tumor and metastases.

Adverse prognostic factors have been well-established and include age, extrathyroidal extension, tumor histology, primary tumor size, and distant metastasis [1]. Extrathyroidal extension has the greatest negative impact on prognosis, with 10-year overall survival rates dropping to 45% in patients who have extrathyroidal extension compared with 91% for those who have encapsulated tumors [2]. CT imaging at our institutions involves multidetector acquisition from the skull base to the tracheal bifurcation with or without contrast. Multiplanar 2-mm axial, coronal and sagittal images are provided for interpretation. Our MRI protocol has a similar coverage from the skull base to the tracheal bifurcation and includes the following sequences: axial and coronal T1-weighted and fat-saturated T2-weighted images, followed by post-contrast axial and coronal T1-weighted images.

## Results and Discussion

### Case 1

A 45 years old female came to the emergency department with stridor and respiratory distress, with a large swelling of the thyroid region more on the right, extending from the chin to the suprasternal notch (Figure 1). The swelling has been progressive and the patient was on alternative medicine. An urgent CECT revealed involvement of both the lobes with tracheal deviation to the left and near total occlusion. There was retrosternal and pretracheal extension of the disease with compression and 360 encompassment of carotid artery. The features shows the inoperability of the disease and also the patient was anesthetically unfit due to difficult airway, where a trial of intubation was failed (Figure 2).



**Figure 1.** CP of the patient showing a large thyroid mass R>L.




**Figure 2.** Axial section of the CECT shows involvement of both the lobes, tracheal compression and shift to the left with 360 encompassment of right carotid artery and pretracheal extension of the tumor.

### Case 2

A 32 years old female came to the emergency with stridor and respiratory distress and a large swelling of the thyroid region. This patient was also on alternative medicine (Figure 3). Urgent reveals enlargement of the right lobe of the thyroid with mediastinal and pretracheal extension (Figure 4). Airway shifts to left and compression of the vessels was seen on the right. The features precluded surgery and indicated difficult airway. Intubation was failed (Figure 5).



**Figure 3.** CP of the patient with large thyroid mass.



**Figure 4.** Sagittal section shows near total occlusion of the airway.



**Figure 5.** Axial section at the level of clavicle shows the retrosternal involvement of the disease.

### Case 3

A 48 year old female with severe respiratory distress came to the emergency department with a large neck swelling extending from the level of hyoid to the suprasternal notch. The disease has been progressive in nature and the patient was on alternative medicine (Figure 6). Urgent CECT showed gross enlargement of the right lobe of thyroid with tracheal shift and near total occlusion of the airway. The lumen was seen infiltrated by the tumor and there was retrosternal and pretracheal extension. As the patient was inoperable and as per the patient attendant's informed consent no intervention was done (Figure 7).



Figure 6. CP of the patient.




Figure 7. Sagittal section shows the near total occlusion of the tracheal lumen.

**Case 4**

A 43 year old female came to the emergency department with severe respiratory distress, where an emergency tracheostomy was performed. CECT revealed involvement of both the lobes of thyroid with retrosternal extension and 360 encompassment of the trachea with compression of airway. Later debulking surgery was performed but the patient died a week post-op due to tracheomalacia (Figure 8).



Figure 8. Axial section shows 360 encompassment of the tracheal by the tumor.

Primary thyroid carcinoma is arising from epithelial follicular cells are classified into Papillary and Follicular thyroid carcinoma which is differentiated and Anaplastic thyroid carcinoma which is undifferentiated. Medullary thyroid carcinoma arises from thyroid parafollicular cells. Papillary carcinoma is the most common and least aggressive whereas Anaplastic carcinoma is the least common and most

aggressive. In their study has revealed that the incidence of thyroid cancer in India has increased over a decade and is 3 times more common in women [3].

Though thyroid cancers can be treated with prompt intervention the delayed presentation with complications precludes intervention, the reasons of which are negligence, alternative treatment modalities, fear of surgery and financial constraints. Patient can develop various distressing symptoms such as pain, hoarseness of voice, dyspnea, dysphonia, hemoptysis, and dysphagia, due to local structure invasion.

Ultrasound is the gold standard at most institutions for the initial evaluation of thyroid disease and can be combined with fine-needle aspiration biopsy for diagnostic purposes. Though USG is most commonly used, the high frequency ultrasound used for thyroid examination has less depth penetration and so the undersurface and extracapsular extension cannot be clearly assessed [4-6]. Hence a higher imaging tool such as CT or MRI is required, which gives details of thyroid malignancy, extrathyroid extension, retrosternal extension, lymph node status especially the central compartment (LN level VI) and invasion of trachea, oesophagus and vascular components [7]. Computed tomography, a modality especially useful because it is directly interpretable by the managing surgeon, allows for a direct visual assessment of the thyroid tumor with respect to adjacent neck structures and is optimal for the assessment of tracheal or cricoid cartilage involvement while also providing excellent resolution of cervical and paratracheal lymph nodes [8] (Table 1).

Radiological features of 4 case scenarios with varied thyroid enlargements extending from the base of the tongue (BOT) .

**Table 1.** Involvement of viscera, vascular components and pretracheal space has been studied.

Characteristics	Result
Gender	
Male	0
Female	4
Age of presentation	30 to 50 years
Symptom at the time of presentation	-
Respiratory distress	4
Stridor	3
Radiological investigation done	-
CXR	4
USG	0
NCCT NECK	0
CECT NECK	4
Parameters assessed	-
a. Extent of the disease	
B/L lobe involvement	2

U/L lobe involvement	2
Right	2
Left	-
Retrosternal Extension	4
b. Tracheal involvement	2
c. Tracheal compression	4
d. Other viscera involved	3
e. Pretracheal space involvement	3
Intervention done	-
Emergency Tracheostomy	1
Surgical debulking	1

Tracheal compression was noted in 100% cases and tracheal involvement in 50% of total cases. Tracheal narrowing of about 70% was noticed in all cases, although the voice of the patient was absolutely normal in all the cases. Vascular involvement was seen in 75% cases and in the form of 360 degree was seen in 25%. Retrosternal extension was noted in 100% cases and pretracheal space involvement in 75% cases. Although intubation was tried in 2 cases and surgical debulking was done in one, all the patients succumbed to the disease due to the lack of airway control.

## Conclusion

CT scan is wiser than ultrasound in extensive disease. CT helps diagnostically and therapeutically in terms of management and decision making and it is ought to be done even in case of emergencies, before any kind of intervention, in-order to avoid any undue circumstances such as on-table death, difficult intubation and deferring-OT. Especially anaplastic carcinoma can be very aggressive, where CT assessment is of utmost importance in view of management. It helps in assessing the risk for any intervention. It gives a cue if patient can be

salvaged. Lastly it poses a great medico-legal importance for the treating surgeon.

## References

1. Shah JP, Loree TR, Dharker D, et al. Prognostic factors in differentiated carcinoma of the thyroid gland. *Am J Surg.* 1992;164:658-61.
2. Andersen PE, Kinsella J, Loree TR, et al. Differentiated carcinoma of the thyroid with extrathyroidal extension. *Am J Surg.* 1995;170:467-70.
3. Kevin Wang, Feitong Lei, Quan Chen, et al. Trends in thyroid cancer incidence in India. *J Clin Oncology.* 2020;105:2507-14.
4. Goyal A, Gupta R, Mehmood S, et al. Palliative and end of life care issues of carcinoma thyroid patient. *Indian J Palliat Care.* 2012;18:134-37.
5. Blum M, Feingold KR, Anawalt B, et al. *Ultrasonography of the Thyroid.* South Dartmouth (MA): MDText.com, Inc. 2000.
6. Shimamoto K, Satake H, Sawaki A, et al. Preoperative staging of thyroid papillary carcinoma with ultrasonography. *Eur J Radiol.* 1998;29:4-10.
7. Ishigaki S, Shimamoto K, Satake H, et al. Multi-slice CT of thyroid nodules: comparison with ultrasonography. *Radiat Med.* 2004;22:346-53.
8. Price DL, Wong RJ, Randolph GW, et al. Invasive thyroid cancer: management of the trachea and esophagus. *Otolaryngol Clin North Am.* 2008;41:1155-68.

## \*Correspondence to

Dr. Shivaam Kesarwaani

Department of ENT and Head-Neck Surgery

Calcutta National Medical College

Kolkata, India

E-mail: shivaam.kesarwaani@gmail.com