











14. Silva N, Eremeeva ME, Rozental T, et al. Eschar-associated spotted fever rickettsiosis, Bahia, Brazil. *Emerg Infect Dis*. 2011;17:275.
15. Kumar M, Krishnamurthy S, Delhikumar CG, et al. Scrub typhus in children at a tertiary hospital in southern India: clinical profile and complications. *J Infect Public Health*. 2012 ;5:82-88.
16. Huang CT, Chi H, Lee HC, et al. Scrub typhus in children in a teaching hospital in eastern Taiwan, 2000-2005. *City*. 2009;4:14-3.
17. Dolado M, JC PN. Meningoencephalitis due to *Rickettsia conorii*. Etiopathological, clinical and diagnostic aspects. *Neurologia* 1994;9:72-5.
18. Tzavella K, Hatzizisis IS, Vakali A, et al. Severe case of Mediterranean spotted fever in Greece with predominantly neurological features. *J Med Microbiol*. 2006;55:341-343.
19. Aliaga L, Sanchez-Blazquez P, Rodríguez-Granger J, et al. Mediterranean spotted fever with encephalitis. *J Med Microbiol*. 2009;58:521-525.
20. Hullatti C, Latha GS, Babu VB. Hyponatremia: a diagnostic marker for the diagnosis of Rickettsial diseases. *Int J ContempPediatr*. 2017;4:696-699.
21. Mittal V, Gupta N, Bhattacharya D, et al. Serological evidence of rickettsial infections in Delhi. *Indian J Med Res*. 2012;135:538.
22. Purvis JJ, Edwards MS. Doxycycline use for rickettsial disease in pediatric patients. *Pediatr Infect Dis J*. 2000;19:871-874.
23. Raoult D, Weiller PJ, Chagnon A, et al. Mediterranean spotted fever: clinical, laboratory and epidemiological features of 199 cases. *Am J Trop Med Hyg*. 1986;35:845-850.
24. Dolado M, JC PN. Meningoencephalitis due to *Rickettsia conorii*. Etiopathological, clinical and diagnostic aspects. *Neurologia*. 1994;9:72-75.

**\*Correspondence to**

Dr. Prakash VJ

Department of Paediatrics

Shri BM Patil Medical College and research centre,

Vijayapura

India

E-mail: drjprakashvs@gmail.com