

Role of Tumor necrosis factor alpha, Malondialdehyde & serum Iron in Anemic Tuberculosis Patients

Rashmi kulkarni¹, Ajit Deshpande², Kiran Saxena³, Akhouri RS Sinha⁴, Meena Verma⁵, Ravi Saxena⁶.

¹Department of Biochemistry Shri Aurobindo Institute of Medical Sciences, India

²Department of Community Medicine Shri Aurobindo Institute of Medical Sciences, India

³Department of Biochemistry Chirayu Medical College Hospital, Bhopal, India

⁴Department of Biochemistry College of Medical sciences, Amargadh. India

⁵Department of Biochemistry MGM Medical College, Indore, India

⁶Department of Physiology. College of medical sciences, Amargadh, India

Abstract

The role of Tumor Necrosis Factor alpha, Malondialdehyde and serum Iron in inducing anemia in Tuberculosis patients was studied. The present study was conducted in the Department of Biochemistry, Shri Aurobindo Institute of Medical Sciences, Indore. The study comprised of sixty-eight anemic pulmonary tuberculosis patients. Levels of Hemoglobin, Iron, TIBC, Ferritin, MDA & TNF α were analyzed in all of them. The inflammatory parameters TNF α & Ferritin along with oxidative marker malondialdehyde were found to be increased in anemic tuberculosis patient and levels of Hb, iron & TIBC was decreased. Overall correlation shows that the levels of Hb, Iron & TIBC were negatively correlated to the levels of Ferritin, MDA & TNF α . Ferritin was negatively correlated with Iron ($r = -0.261$, $p < 0.05$) & showed a positive correlation with MDA. ($r = 0.304$, $p < 0.05$). For Ferritin & TNF α correlation was positive, it was not significant ($r = 0.114$, NS). There was a positive significant correlation between Iron & TIBC. ($r = 0.666$, $p < 0.01$) & a negative significant correlation between the TNF α levels and the hemoglobin concentration ($r = -.273$, $p < 0.05$). Hypoferremia was induced as a consequence of inflammatory response and action of free radicals in anemic tuberculosis patients

Key word: Anemia, Hypoferremia, Ferritin, TNF α , Malondialdehyde.

Accepted September 19 2010

Introduction

Active tuberculosis infection is generally associated with major abnormalities in hemopoiesis [1]. Anemia is a common complication of pulmonary tuberculosis, the reported prevalence ranging from 16 to 76% in different studies [2]. The precise mechanism of anemia in pulmonary tuberculosis patients (PTB) is not clearly known however, anemias due to inflammation & iron deficiency have been implicated [3,4]. The pathogenesis of anemia of chronic inflammation may evolve due to cytokine-mediated defense against microbial pathogens, effectively withholding iron from microbes, which incidentally also deprive erythroid precursors of their iron supply [5,6,7]. The development of hypoferremia thus results in the development of iron-restricted erythropoiesis and anemia. Cytokine also induced alterations in cellular iron homeostasis are mediated in part by the increased production of nitric oxide (NO) and reactive oxygen species (ROS) [8].

The present study was taken to find out the role of Tumor Necrosis Factor alpha, Malondialdehyde & serum Iron in inducing Anemia in Tuberculosis Patients.

Material and Method

Present study was conducted in the Department of Biochemistry, Shri Aurobindo Institute of Medical sciences, Indore. Sixty-eight patients of anemic pulmonary tuberculosis were selected. The patients of pulmonary tuberculosis were taken from Shri Aurobindo Institute of Medical Sciences and Manorama Raje Tuberculosis Hospital, Indore. Pulmonary tuberculosis patient falling in DOTS category CAT I & aged between 18-55 yrs & hemoglobin (Hb) less than <13 g/dL for Men & HB <12 g/dL for non-pregnant Women was included. Tuberculosis patient with other disease or disorder and patients with incomplete data records and moderate to severe injury or surgery during the previous month were excluded. Those patients

who have taken antibiotics for cough like Amoxicillin, Septran, Ciprofloxacin etc were included but those patients with history of prior anti-TB treatment were not included. Patient with extra pulmonary involvement, blood donors, subjects with history of smoking and who had taken iron supplementation were excluded. The study was approved by the Institutional & Human Ethical Committee of the Sri Aurobindo Institute of Medical sciences, Indore. Informed consent was taken from each subject and then they were enrolled in the study.

5ml of fasting venous blood samples were drawn from subjects, each sample was divided into two parts: the first part was collected in EDTA while the second sample was collected in a plain tube and left to clot. Serum was obtained after centrifuge. The serum samples were aliquoted and stored according to the instructions in the kits, until the various assays described below were performed

Haemogram was estimated by automated cell counter. (Sysmex KX) Serum iron and total iron-binding capacity (TIBC) were estimated on an autoanalyser from Rosh diagnostics using commercial test kits from Randox Laboratories, UK. Ferritin was estimated by direct immuno-enzymatic determination by using test kits from Diametra, Italy. Serum TNF α was estimated by using a commercially available immunoassay Kit from DIA-CLONE, France. Serum Malondialdehyde was estimated spectrophotometrically by thiobarbituric acid method of Wilbur K M et al (1949) [9].

Correlations were calculated by Pearson's correlation coefficients (two-tailed) using SPSS Version 10. P value \leq 0.05 was used as a threshold of significance.

Result

The levels of Hb, iron & TIBC was decrease than normal ranges & ferritin, MDA & TNF values found to be in

creased than normal ranges in anemic tuberculosis patients (Table 1)

Pearson correlations (Table no 2) were carried out on the different variables, which showed that the decreasing levels Hb, Iron & TIBC was negatively correlated to the increasing levels Ferritin, MDA & TNF α .

The statistically significant correlation was found between Hb & TNF α ($r = -0.273$, $p < 0.05$), Iron & TIBC ($r = 0.666$, $p < 0.01$), Iron & ferritin ($r = -0.261$, $p < 0.05$) & Ferritin & MDA ($r = 0.304$, $p < 0.05$).

Table 1. Mean \pm SD values for hemoglobin, iron related parameters and inflammation & oxidative marker in anemic tuberculosis patient group

Parameters	Normal ranges	Anemic TB Patients n=68
		Mean \pm SD
Hb (g/dl)	Males > 13 Females > 12	9.08 \pm 1.34
Iron (μ mol/l)	Males = 10.6 – 8.3 Females = 6.6- 26.0	7.38 \pm 1.79
TIBC (μ g/dl)	250- 450	178.01 \pm 44.0
Ferritin (ng/mL)	Males = 6 – 350 Females = 20 – 400	321.25 \pm 77.8
MDA (nmol/ml)	>3	5.82 \pm 0.92
TNF (pg/ml)	> 8	57.55 \pm 14.7

Table 2. Pearson's Correlations between the Variables

	IRON r value	TIBC r value	Ferritin r value	MDA r value	TNF r value
Hb (g/dl)	.179	.172	-.033	-.120	-.273 *
IRON (μ mol/l)	-	.666**	-.261*	-.040	-.128
TIBC (μ g/dl)	-	-	-.114	-.139	-.147
Ferritin (ng/mL)	-	-	-	.304*	.114
MDA (pg/ml)	-	-	-	-	.131

r = Correlation coefficient. r^{**} = Correlation significant at $p < 0.01$ & r^* = Correlation significant at $p < 0.05$

Discussion

Anemia is one of the abnormalities found in TB Patients due to poor nutritional causes and increased in cytokines

.....

level¹. Therefore, both anemia of inflammation as well as iron deficiency anemia may coexist in pulmonary tuberculosis. Our findings were consistent with the previous reports that there was decrease in Hb, Iron & TIBC levels and increased levels of ferritin, TNF α & MDA in tuberculosis patients [10, 11, 12, 13, 14, 15].

Hypoferremia in disease associated anemia appears to be the result of a complex network of cytokines, immune cells, and acute-phase proteins [16]. TIBC values are usually high in iron-deficiency anemia and low in anemia of inflammation [17,18]. The low TIBC concentrations seen in our present patients might have been the result of an acute-phase response. Ferritin concentration in blood is considered a specific indicator of body iron stores [19], however, the concentrations can rise following an inflammatory response, irrespective of iron status [20,21,23,].

Endogenous TNF α response during inflammation is a sufficient proximal stimulus to mediate an associated anemia with decreased RBC survival & impaired erythropoiesis [24].

Recombinant TNF α and recombinant interleukin 1 (IL-1) cause a significant decrease in serum iron levels accompanied by an increase in circulating ferritin concentration and a decrease in soluble transferrin receptor levels [25,26]. Increased levels of TNF α , ferritin & decreased iron levels in present study & their correlation with each other was in accordance with the finding that tumour necrosis factor alpha induce hypoferraemia and ferritin production [16]. Similar findings was stated by Bayes et al [11] for C-reactive protein (CRP), iron & ferritin.

The increased levels of TNF α & MDA & decreased levels of Hb, Iron, TIBC reveals that the MDA along with TNF α was involve in redistribution of iron. These findings are in concordance with the literature [26]. Cytokines affect iron homeostasis in monocytes by inducing the formation of short-lived radicals such as nitric oxide (NO) and reactive oxygen species (ROS) which affect the post-transcriptional regulation of iron homeostasis via iron regulatory proteins (IRPs) [27,28].

In anemic tuberculosis patients disturbances in iron homeostasis may be one of the cause to the shift of iron from a transferrin- bound available state to a ferritin-incorporated storage state. These changes are the components of a systemic inflammatory response and action of free radical, which may be one of the major factors in the pathogenesis of anemia.

References

1. Sei Won Lee, Young Ae Kang, Young Soon Yoon, et al. The Prevalence and Evolution of Anemia Associated with Tuberculosis. *J Korean Med Sci* 2006; 21: 1028-32.
2. Goldenberg AS. Haematological abnormalities and mycobacterial infections. In *Tuberculosis 1996*; pp. 645–652 [WN Rome and S Garay, editors]. Boston, MA: Little Brownand Company.
3. Douglas SW & Adamson JW. The anaemia of chronic disorders: Studies of marrow regulation and iron metabolism. *Blood* 1975; 45: 55–65
4. Das BS, Nanda NK, Rath PK, et al. Anaemia in acute, *Plasmodium falciparum* malaria in children from Orissa state, India. *Ann Trop Med Parasitol* 1999;93: 109–118.
5. Bullen JJ, Rogers HJ & Griffiths E. Role of iron in bacterial infection. *Curr Top Microbiol Immunol* 1978; 80: 1-35.
6. Weinberg E. Iron and infection. *Microbiol Rev* 1978 ; 42: 45-66.
7. Jurado RI. Iron, infection, and anaemia of inflammation. *Clin Infect Dis* 1997; 25: 888-895
8. Igor Theurl, Verena Mattle, Markus Seifert, et al. Dysregulated monocyte iron homeostasis and erythropoietin formation in patients with anemia of chronic disease. *Blood may* 2006; 107(10): 41 4142-4148
9. Wilbur KM, Bernheim F, Shapiro OW. The thiobarbituric acid reagent as a test for the oxidation of unsaturated fatty acid by various agents. *Arch Biochem Biophys* 1949;24:305-313
10. Uma Devi, C. Mohan Rao, Vinod K. Srivastava, et al. Effect of iron supplementation on mild to moderate anaemia in pulmonary tuberculosis. *British Journal of Nutrition* 2003; 90: 541-550
11. Baynes RD, Flax H, Bothwell TH, et al. Hematological and iron-related measurements in active pulmonary tuberculosis. *Scand J Haematol* 1986; 36: 280-7
12. Kaminskaia GO, Abdullaev RIu. Iron metabolism in patients with different variants of pulmonary tuberculosis *Probl Tuberk.* 2002; 12: 49-51
13. Kaminskaia GO, Abdullaev RIu, Baturova GA, et al. The specific features of iron intake in patients treated for pulmonary tuberculosis *Probl Tuberk Bolezn Legk.* 2009; 7: 46-55
14. Nakaya M, Yoneda T, Yoshikawa M, et al. The evaluation of interleukin-8 (IL-8) and tumor necrosis factor-alpha (TNF-alpha) level in peripheral blood of patients with active pulmonary tuberculosis. *Kekkaku.* 1995; 70 8: 461-466.
15. Kharakter ZhZ, Skoraia RI, Platonova IL. The ceruloplasmin and lipid peroxidation indices in patients with pulmonary tuberculosis. *Vrach Delo.*1989; 10: 55-7
16. Feelders RA, Vreugdenhil G, Eggermont AM, et al. Regulation of iron metabolism in the acute-phase response: interferon gamma and tumour necrosis factor alpha induce hypoferraemia, ferritin production and a decrease in circulating transferrin receptors in cancer patients. *Eur J Clin Invest.* 1998; 28: 520-527

17. Fleck A & Myers MA. Diagnostic and prognostic significance of acute phase proteins. In *The Acute Phase Response to Injury and Infection* 1985:249–271 [AH Gordon and A Koj, editors]. Amsterdam: Elsevier Science Publishers.
18. Punnonen K, Irjala K, Rajamaki A. Iron deficiency anaemia is associated with high concentrations of transferrin receptor in serum. *Clin Chem* 1994; 40: 774 - 776
19. Lipschitz DA, Cook JD, Finch CA. A clinical evaluation of serum ferritin as an index of iron status. *New Eng J Med* 1974; 290: 1213-1216.
20. Dallman PR, Reeves JD, Driggers DA, et al. Diagnosis of iron deficiency: the limitations of laboratory tests in predicting response to iron treatment in 1-year-old infants. *J Pediatr* 1981;98: 376- 381.
21. Harju E, Pakarinen A & Larimi T. A comparison between serum ferritin concentration and the amount of bone marrow stainable iron. *Scand J Clin Lab Invest* 1984; 44: 555-556.
22. Henderson A Ferritin levels in patients with microcytic anaemia complicating pulmonary tuberculosis. *Tubercle* 1984;65 : 185-189.
23. Taylor PG, Martinez-Torres C ,Mendez-Castellano H, et al. The relationship between iron deficiency and anaemia in Venezuelan children. *Am J Clin Nutr* 1993; 58: 215-218.
24. Lyle L, Moldawer, Michael A. Marano, He Wei, Yuman Fong, et al. Cachectin/tumor necrosis factor- α alters red blood cell kinetics and induces anemia in vivo. *The FASEB Journal* March 1989; 3: 1637- 1643
25. Alvarez-Hernandez X, Liceaga J, McKay IC, et al. Induction of hypoferrremia and modulation of macrophage iron metabolism by tumor necrosis factor. *Lab Invest.* 1989; 61: 319-322
26. Maciejewski, JP, Selleri C, Sato T et al. Nitric oxide suppression of human hematopoiesis in vitro. Contribution to inhibitory action of interferon- γ and tumor necrosis factor- α . *J Clin Invest* 1995; 96: 1085-1092
27. Drapier JC, Hirling H, Wietzerbin J, et al. Biosynthesis of nitric oxide activates iron regulatory factor in macrophages. *EMBO J* 1993; 12: 3643-3649
28. Cairo G, Recalcati S, Pietrangelo A, et al. The iron regulatory proteins: targets and modulators of free radical reactions and oxidative damage. *Free Radic Biol Med.* 2002; 32: 1237-1243

Correspondence to:

Rashmi A Kulkarni
Department of Biochemistry,
Shri Aurobindo Institute of Medical Sciences
Indore, Madhya Pradesh
India

.....