

## Stab wound causing uncontrollable bleeding in the abdomen.

AJ Feinstein\*

Trauma, General Surgery and Critical Care, Banner University Medical Center, Phoenix, USA

Accepted on 25 October, 2021

### Introduction

Introduction: A 32-year-old man presented with a single left lateral lower quadrant stab wound and pulsatile hemorrhage. The patient was taken immediately to the operating room for laparotomy and was found to have an expanding retroperitoneal hematoma. Aortic occlusion did not slow progression. The left iliac artery could not be located. A femoral cut down and proximal control of the femoral artery was also not helpful. The hematoma was entered directly and the hemorrhaging area in the lateral psoas muscle was ligated. Results: The patient maintained palpable dorsal is pedis pulses throughout the procedure. Postoperatively, a CT (computed tomography) showed congenital absence of the left iliac artery with large retroperitoneal collateral vessels in the retro peritoneum that reconstituted in a normal femoral artery. The scan demonstrated the injury was to one of these large vessels. The patient was discharged home after an uneventful course. Discussion: Congenital malformations of the iliac artery are rare, and most are discovered incidentally on imaging. We report a case of injury to a collateral vessel resulting from a left iliac atresia. This had implications for controlling hemorrhage in this case. Knowledge of vascular anomalies is important for management of traumatic injury.

A 32-year-old man presented to our Level I trauma center after being stabbed in the abdomen. Paramedics reported finding a significant amount of blood at the scene and he was transiently hypotensive during transport. On arrival, he was complaining of lightheadedness and he was diaphoretic and tachycardic. Examination revealed a 3 cm stab wound just medial to the left anterior superior iliac spine. When direct pressure was released from the wound, copious pulsatile hemorrhage was apparent. No other traumatic injuries were noted.

Large bore intravenous lines were placed, blood transfusion was initiated and the patient was taken emergently to the operating room. In the operating room, the patient was placed in a supine position, intubated and widely prepped from the neck to the knees. A midline laparotomy was performed while maintaining direct pressure on the wound. There was no free fluid in the abdomen. A large hematoma was noted in zone 2 on the left and zone 3 of the retroperitoneum. In order to achieve proximal control, a left medial visceral rotation was performed. The distal aorta and right iliac artery were localized, isolated and looped. The left ureter was visualized and preserved. The left iliac artery could not be located. Several attempts were made to follow the iliac artery distally from the aorta without success. The hematoma continued to expand laterally in zone 2 and into the pelvis.

When the distal aorta was occluded, the hematoma appeared to expand more rapidly and did not diminish the pulsatile hemorrhage from the wound when pressure was released. During this time, another incision was made inferior to the inguinal ligament. The left proximal femoral artery was dissected and vessel loops were placed around it. Occlusion of the femoral vessel also did not diminish the hemorrhage from the wound when pressure was released. The patient was becoming more hemodynamically unstable despite resuscitation and the decision was made to open the hematoma in the absence of proximal control. The hematoma was entered adjacent to the stab wound in the lateral inferior portion of zone 2. On exposure of the psoas muscle, a large amount of arterial hemorrhage was encountered from a defect in the muscle. Several large sutures were placed using absorbable suture around the defect until the arterial hemorrhage stopped. Pressure was released from the wound and there was minimal bleeding. There was a diminished but palpable pulse in the left foot. Vessel loops were removed and the retro peritoneum and groin were packed with laparotomy pads. A negative pressure dressing was placed in the abdomen and the patient was taken directly to the CT scanner.

A contrast CT of the chest, abdomen and pelvis revealed congenital absence of the left iliac artery. Large collaterals, some arising from the thoracic aorta, reconstituted in the left femoral artery, which had good filling and runoff. The area of abdominal packing indicated the disruption of one or more of these collateral arterial vessels. The patient was taken back to the operating room the following day for removal of the abdominal packing and closure. His pulses remained intact and his hospital course was uneventful. He was discharged home on post-operative day seven.

### \*Correspondence to

AJ Feinstein

General Surgery and Critical Care

Banner University Medical Center

Phoenix

USA

E-mail: feinstein@gmail.com

**Citation:** Feinstein AJ. Stab wound causing uncontrollable bleeding in the abdomen. *J Trauma Crit Care* 2021;5(5):13.