

Status of Vaping in India and Its Efficacy as a Smoking Harm Reduction Approach

VipinThamp

Indian Institute of Public Health-Delhi, India

Abstract

An electronic cigarette is a handheld electronic device that tries to create the feeling of tobacco smoking it is a form of an ENDS (Electronic nicotine delivery system). They are battery-powered devices that delivery nicotine in an aerosol to the user. E-cigarettes contain nicotine cartridges with airflow sensors, but do not burn tobacco¹. E-cigarette was introduced in the market as an alternative approach /device for quitting smoking.² Vaping is the act of inhaling and exhaling the water vapor produced by an electric device called vaporizer³. E-Cigarettes can be categorized into three groups disposable, rechargeable and modular can be refilled by the user. E-cigarette devices were manufactured mainly in China initially when it was launched into market in 2007⁴. There are over 466 brands and 7764 unique flavors with about 242 new flavors added per month⁵. E- Cigarettes are mostly available online. Dampf, vape stop, greenvapo and develop are the fundamental online stores accessible in India⁶. A single dispensable E-Cigarettes ranges from \$ six to \$ twelve (INR 390-782) and cartridge models. are rechargeable and contain pre-filled cartridges starter kits usually range around \$ forty to \$ sixty (INR 2600 -3900)⁷.

The usage of the E-Cigarettes is more in France and United Kingdom which represents 6 percent and 3.5 percent of the adult population¹. In United states e-cigarette usage has been highest among never smokers aged 18-24 yrs contributing about 9.7 percent of the total population as compared to other age groups⁸. In India, there are almost 75 companies which are supplying e - cigarettes through online in India according to association of vapers India and use is more common in adolescents because consumption of e- cigarettes occurs through online and there is no authority website protection for e- cigarette advertisement and it is available in various flavors. States like Karnataka, Kerala, Punjab, J&K, Mizoram & Maharashtra have banned e- cigarettes under the Drugs and Cosmetics Act, 1940, and Food Safety & Standards (Prohibition and Restriction on Sales) Regulation, 2011⁹.

Conventional cigarettes have been used for more than a century and smokers are preferring conventional cigarettes

as compared to e-cigarettes worldwide, more information about the long term health hazards of the conventional cigarettes is available, whereas e-cigarettes are newer products which has been implemented 10 years before because of this the data available on the long term health consequences is limited .Some studies are mentioning the benefits of e-cigarettes over conventional cigarette and in contrast other studies are mentioning its health hazards equal to conventional cigarette so it is difficult to come to a conclusion that whether the product reduces morbidity and mortality. The present study has been conducted to understand the status of vaping and its efficacy as an alternative product for tobacco cessation.

Methods

Data sources and searches

Literature search strategies was developed using medical subject headings (MESH) based on the terms “electronic cigarette(s)” OR “e-cig” OR “electronic nicotine delivery system(s)” OR “alternative -nicotine delivery devices” OR “vaping” What's more, "status AND "quit" OR "smoking suspension" OR "smoking decrease". This electronic based search strategy was used for identification of published papers in PubMed. Search results were not limited by language, but all identified studies were in English. This strategy for the date bases was run or conducted from January 2007 to March 2018.

A total of 11 unique articles met the inclusion criteria. Articles titles and abstracts were screened based on the status of vaping and its role on tobacco cessation in addition relevant reports, articles, journals were referred from electronic cigarette research website.

Criteria for considering studies for the review

Inclusion criteria Eligible

Study designs included systematic reviews and meta-analysis, analytical study designs (case-control studies, cohort studies, analytical cross-sectional studies), descriptive studies (case report, case series, cross-sectional studies), qualitative research and outcome studies on the

status and the efficacy of e-cigarettes as a smoking cessation or reduction device were included. We considered all study populations that were defined as “adult” by the study authors. The studies that defined adult smokers, current smokers, former smokers, and dual users were used for further analysis.

Exclusion Criteria

Editorials, conference papers, studies which were unrelated to topic concerned and studies were excluded. At the end articles relevant will be reviewed.

Data extraction

A standardized data extraction form was developed to collect information from the selected studies done on e-cigarettes and vaping. Standard data describing each study was extracted, this includes general information of the study (author, year of publication), study characteristics (study design, study location), primary and secondary outcome measures and end points, study strengths and limitations.

Selection of Studies

Studies were reviewed based on the exclusion and inclusion criteria, in three stages.

- During the first stage of title screening, titles of the studies identified from the search were assessed for inclusion.
- If the study was related to the objectives of the current study, then it was moved to abstract screening. In the next stage of abstract screening, abstracts of these selected titles that were approved and included for the final stage of full text screening.
- In the third stage of full text screening, full texts of abstracts selected in the previous stage were screened for eligibility.

Discussion

Initial searches conducted via PubMed using the key words electronic cigarette, e-Cigarette, vaping, and electronic nicotine delivery systems, yielded a total of 537 records were retrieved through the Search Strategy, of which 188 articles were screened for eligibility after full text

screening. Finally, 11 articles were included for the review.in which one article could be extracted from the first objective (status of vaping in India) and nine articles were extracted related to the second objective (harm reduction approach-smoking reduction) and Figure 1, shows the flow chart of the search strategy.

1.To know the presence and usage of E-cigarettes in India

The data on e-cigarettes in India both in offline outlets, as well as online retail websites, is still not available. E-cigarettes are available in wide variety of flavors like chocolate, strawberry, blueberry etc. flavors similar to cigarette flavors such as Tobacco, Classic Original, Cool Menthol etc. to more exotic flavors such as Cappuccino, Double Apple, Virginia Leaf etc. One can also find dessert flavors such as Caramel Mocha etc. and the possibilities are truly unimaginable and endless¹⁰. Nicotine strength will depend on your current smoking habits or choice of cigarette you smoke. If you are a heavy smoker then 1.8%, 1.6%, or a similar %, is a good starting point. A lighter smoker could start with 1.2%, 0.8% or similar and someone who prefers no nicotine can choose 0. A study conducted in India to check for the online ENDS availability reported that findings out of 65 models (34 brands) which include 45 models of electronic cigarettes,12 e-shisha, 2 e-hookah and 6 e-cigar. In these products information regarding the ingredients, flavors, nicotine strength and claims such as “health related-no tar no harm environmental friendly” were mentioned¹¹.But studies on e-cigarettes are claiming it as not an eco-friendly product as it contains substances like formaldehyde, acetaldehyde and acrolein that are released into the environment which are classified as carcinogens by International Agency for Cancer Research (IARC).Apart from that multiple studies have reported both acute and long-term effects on health so long-term studies on humans must be done on this topic in order to generate an evidence. Few states in India such as Punjab, Karnataka, and Maharashtra have taken steps to cease the sale of ENDS in areas under their jurisdictions¹².

2. To assess whether use of E-Cigarettes can be helpful in reduction of behavior related to smoking cigarettes

E-cigarette manufacturers and users suggest that e-cigarettes are used as smoking-cessation aids as a large proportion of users manage to quit smoking or substantially reduce smoking consumption^{13–16},but these studies used

convenience samples and suffer from selection bias and do not generalize the general population. In an investigation led on the clients drawing nearer vape shops evaluated the smoking status of members by estimating breathed out carbon monoxide and distinguished that 66 percent of members had quit smoking.¹⁷ This study cannot access the random sample of users and could over-represent the dedicated and advanced consumers. Randomized controlled trials represent the gold-standard in assessing the efficacy of any medical intervention. Studies have shown better results for newer generation and modest results for first generation.¹⁸ Many cohort studies have shown mixed results with some showing the odds of quitting while others showing the opposite effect ^{19–21}. Meta-analysis has also shown mixed results and systematic review have reported that the e-cigarettes help in quitting smoking.^{22,23} Another systematic review found that e-cigarette use was associated with 28 percent reduced chances of quitting,²⁴ while the most recent one found limited evidence for a positive or negative effect of e-cigarettes on smoking cessation, again rating the evidence as low or very low certainty²⁵. The systematic reviews were of good quality, there were major problems in the studies that were included to the analyses and were rated 'low' by GRADE standards due to the small number of trials, low event rates and wide confidence intervals around the estimated means. There is an evidence based on the meta-analysis done that usage of e-cigarettes leads to use of conventional cigarettes. There is a reverse temporal association also that usage of combustible tobacco cigarettes at the baseline was positively associated with the use of e-cigarettes. Follow up must be done for months' period. The two randomized controlled trials used outdated and poor-quality products that were already withdrawn and replaced by more advanced products at the time of the studies' publication.^{26,27} The third study provided nicotine-containing e-cigarette liquids for a limited period of time, while for the rest of the study follow up participants could not easily obtain such liquids because they were banned in that country at the time of the study.²⁸ Cohort studies suffered from very strong bias as many of them included subjects who had already failed to quit smoking with the use of e-cigarettes at baseline, resulting in bias of the outcome being present at the start of the studies.²¹ Additionally, they usually assessed ever or current (past 30-day) use, definitions that include a lot of experimenters rather than regular users ²⁹. Population studies have shown that e-cigarettes have helped people

quit smoking, with increasing proportions of e-cigarette users being former smokers. There has been an increase in prevalence of e-cigarette use from 700,000 in 2013 to 2.9 million in 2017.³⁰ The use of the e-cigarettes have risen from 33% in 2012 to 52% in 2017.

In the European Union, an estimated 6.1 million smokers have managed to quit smoking with the help of e-cigarettes, while an additional 9.2 million smokers have reduced their smoking consumption.³¹ When current and daily e-cigarette use was assessed separately from ever-use reported smoking cessation and reduction rates were by far higher^{31,32}. Cross sectional studies showed 60% higher odds of quitting smoking as compared to other nicotine replacement therapy.³³ Cross-sectional studies also have serious limitations, such as the lack of temporal association and causality, self-report bias and no objective assessment of the smoking status or the span of smoking end. Likewise, these examinations neglect to investigate what number of the individuals who guarantee that they have halted with the guide of e-cigarettes would have halted in any case and what number of the individuals who utilized an e-cigarette yet neglected to stop would have halted had they utilized another method³⁴.

Conclusion:

The present study concluded that there is a lack of data regarding the use of e-cigarettes and its safety and efficacy. In order to cover this lacunae, we need to give emphasis on the reporting of e-cigarette in the GATS (Global Adult Tobacco Survey). To examine whether e-cigarettes can be used as a source of harm reduction in the population and to find out the effects of E-cigarettes on them, extensive research should be carried out to explore the complex interactions between smoking and E-Cigarette. Strength This study performs the comprehensive review of literature to know the status and presence of e-cigarettes in India and the efficacy of the e-cigarettes as a smoking cessation device globally.

Limitations

1. Articles related to the topic were scarce and access to those data sources were limited
2. Review studies were considered mainly for data analysis as the systematic review and meta-analysis were limited in number.

Conflict of interest

There was no conflict of interest

Funding

No Funding source available

Acknowledgement

I would like to thank faculties of my institution Indian Institute of Public Health, Delhi (IIPH) for giving me support and guiding me in completing my research, I would like to thank my supervisor Dr. Rajesh Nair without whose support this research would not have been possible.

References

1. SEATCA_E-Cig. SEATCA_E-Cig.2014.
2. Heydari G, Ahmady AE, Chamyani F, Masjedi M, Fadaizadeh L. Electronic cigarette, effective or harmful for quitting smoking and respiratory health: A quantitative review papers. Lung India.2017;34(1):25.
3. Ernst et al E cigarette. EY-E-Cigarette-Report.Pdf; 2016.
4. Grana R, Benowitz N, Glantz SA. E-cigarettes: a scientific review. Circulation. 2014;129(19):1972-1986.
5. Huey S. Escape the vape: Health hazards of the latest nicotine craze : Nursing2018.2018.
6. Online Shopping India. Which are the best online stores for buying electronic cigarettes& liquids in India? - Quora.2017.
7. IEC Vaping Authority. E-Cigarettes vs. Cigarettes: A Cost Analysis – IEC Vaping Authority. IEC Best E-Cig & Vape Guides.2018.
8. NHIS US E-Cigarette. NHIS US E-Cigarette. 2015.
9. Times of India. Govt likely to stub out 'toxic' e-cigarettes - Times of India. The Times of India. <http://timesofindia.indiatimes.com/india/govt-likely-to-stub-out-toxic-e-cigarettes/articleshow/59855386.cms>. Published 2017. Accessed September 12,2017.
10. Lite joy Ukindia 2018. How to choose your eliquid | LitejoyIndia.
11. Mohanty VR, Chahar P, Balappanavar AY, Yadav V. Electronic Nicotine Delivery Systems (ENDS): Mapping the Indian Online Retail Market. Nicotine Tob Res. 2017;19(11):1386- 1389.
12. Zee Media Bureau. Karnataka becomes third state to ban e-cigarettes in the country. Zee News. 2016.
13. Dawkins L, Turner J, Roberts A, Soar K. "Vaping" profiles and preferences: an online survey of electronic cigarette users. Addiction.2013;108(6):1115-1125.
14. Farsalinos KE, Romagna G, Tsiapras D, Kyrzopoulos S, Voudris V. Characteristics, Perceived Side Effects and Benefits of Electronic Cigarette Use: A Worldwide Survey of More than 19,000 Consumers. Int J Environ Res Public Health.2014;11(4):4356-4373
15. Farsalinos KE, Romagna G, Tsiapras D, Kyrzopoulos S, Spyrou A, Voudris V. Impact of Flavour Variability on Electronic Cigarette Use Experience: An Internet Survey. Int J Environ Res Public Health.2013;10(12):7272-7282.