

The potential of kebar grass (*Biophytum petersianum klotzsch*) on mice (*Mus musculus*) during lactation period on liver histopathological were exposed by carbofuran.

Epy Muhammad Luqman^{1*}, Muhammad Andika Aulia Rakhman¹, Sri Hidanah¹, Widjiati¹, Viski Fitri Hendrawan²

¹Departement of Veterinary Science, Universitas Airlangga Kampus C Unair Jalan Mulyorejo, Surabaya, Indonesia

²Department of Animal Reproduction, Universitas Brawijaya, Malang, Jawa Timur, Indonesia

Abstract

This study aims to determine the potential of kebar grass extract in reducing the impact of liver damage in mice (*Mus musculus*) during lactation exposed to carbofuran. The 42 lactation mice (*Mus musculus*) used in the study were divided into seven groups, each group consisting of six mice. Carbofuran, chewed grass, and vitamin C are exposed orally on days 1 to 14 after birth. This group consisted of C (aquadest control), P1 (carbofuran 1/4 LD₅₀ 0.0125 mg/day), P2 (carbofuran 1/8 LD₅₀ 0.00625 mg/day), P3 (carbofuran 1/4 LD₅₀ 0.0125 mg/day+extract of kebar grass 3.375 mg 0.2 ml), P4 (carbofuran 1/8 LD₅₀ 0.00625 mg/day+extract of kebar grass 3.375 mg 0.2 ml), P5 (carbofuran 1/4 LD₅₀ 0.0125 mg/day+vitamin C 5 mg 0.2 cc), and P6 (carbofuran 1/8 LD₅₀ 0.00625 mg/day+vitamin C 5 mg 0.2 cc). On the 15th day after birth, mice were sacrificed and their liver organs taken for microscopic examination with hematoxylin-eosin staining. The results of this study indicate that the liver histopathology of mice suffered damage characterized by an increase in the number of inflammation cell, degeneration, and necrosis in line with the carbofuran dose given during the lactation period (P1 and P2). The administration of kebar grass extract (P3 and P4) is more effective than vitamin C (P5 and P6) in reducing the amount of inflammation cell, degeneration, and hepatocyte necrosis caused by carbofuran administration in lactation mice. This can be seen in the amount of liver damage in group P3 and P4 mice less than the amount of liver damage in group P5 and P6.

Key words:

Pesticide stress, Carbofuran, Histopathology, Kebar grass, Lactation, liver.

Accepted on 29th September, 2021.

Introduction

Pesticides are widely used to control plant-disturbing organisms such as cabbage, spinach and carrots [1]. Excessive exposure to insecticides can result in disturbed soil fertility. In addition, agricultural product residues will have an impact on human and animal health [2].

Carbofuran is a type of carbamate insecticide that often causes poisoning in non-target organisms such as plants, animals and humans [3]. Excessive use of carbofuran can result in a reduction in the number of individuals, barriers to metabolic activity, behavioral barriers, and reproduction [4].

The target of organ damage due to the effects of carbofuran are the brain, liver, muscles and heart [5]. The liver is the largest glandular organ that has a role in detoxifying the metabolic waste materials of food, drugs, and toxic substances that enter the body [6].

Toxic substances carried in the bloodstream can cause damage to hepatocytes around the central vein which will first experience necrosis. The formation of reactive oxygen species

(ROS) and lipid peroxidation due to ROS can cause hepatocyte death [7].

Carbofuran given orally has been shown to stimulate ROS which causes oxidative stress in the body if the doses are given too much [8]. This imbalance between prooxidants and antioxidants is called oxidative stress. This usually occurs due to lack of antioxidants and excess production of free radicals [9].

Hepatocytes are the main tissue that is targeted for an increase in the concentration of free radicals because the liver is the site of the metabolic process of xenobiotic compounds that will induce hepatocyte death. Carbofuran causes ROS, causing degeneration and necrosis damage and inflammation cell infiltration in the liver [7].

Lactation period is the period that occurs when a mother has given birth to a child. Giving carbofuran to the parent of the lactation period is considered to show more significant results compared to the parent who is not in the lactation period. This is due to the parent in the lactation period more susceptible to certain chemical compounds than the parent not in the lactation period [10].

Citation: Luqman EM, Aulia Rakhman MA, Hidanah S, et al.. The potential of kebar grass (*Biophytum petersianum klotzsch*) on mice (*Mus musculus*) during lactation period on liver histopathological were exposed by carbofuran. *J RNA Genomics* 2021;17(6):1-7.

Kebar grass (*Biophytum petersianum Klotzsch*) is a plant belonging to the family group Oxalidaceae found in Kebar District, West Papua. Kebar grass usually grows naturally and is spread almost throughout the Kebar District. Charged grass contains flavonoids, retinol, and tocopherol.

According to Hudson [11]. Chewed grass contains flavonoids which have the function of primary antioxidants because they are free radical acceptors so they can inhibit free radical chain reactions in lipid oxidation that can prevent membrane damage. Polyphenol compounds such as flavonoids work by converting H_2O_2 to H_2O and O_2 .

The content of retinol or vitamin A in chilled grass functions as an antioxidant. Beta carotene works by reacting with free radicals and causing free radicals to become stable. Beta carotene or retinol in collaboration with vitamin E (tocopherol) contained in kebar grass and vitamin C (ascorbic acid).

Vitamin C which is transformed into radicals is stabilized by the natural antioxidant glutathione [12]. The content of vitamin E in the resistant grass prevents free radicals by giving H atoms from hydroxyl groups to peroxy lipid radicals. Radicals formed from alpha-tocopherol will be stabilized through electron delocation in the aromatic ring [13].

Based on this background it is necessary to conduct research on the potential of kebar grass extract (*Biophytum petersianum Klotzsch*) in reducing the impact of damage to the liver of mice (*Mus musculus*) during lactation exposed to carbofuran.

The data obtained is expected to provide additional reference results and proof about the potential of kebar grass extract in reducing liver damage in mice during lactation exposed to carbofuran orally. In addition, the data obtained can also be used in providing information to the public about the dangers posed by excessive use of carbofuran, especially in the parent during the lactation period.

Materials and Methods

Materials

This study used a Completely Randomized Design with seven treatment groups and six repetitions. The experimental animal used was lactation mice (*Mus musculus*). This research was conducted at the Animal Cage in the Faculty of Veterinary Medicine, Airlangga University.

Mice are placed in plastic cages covered with wire and given husks as a base. Mice were fed pellets and drinking distilled water ad libitum every day during the study. Materials used in this study were mice (*Mus musculus*) lactation period, kebar grass (*Biophytum petersianum Klotzsch*), CMC Na, Ethanol 70%, carbofuran (2,3-Dihydro-2,2-dimethyl-7-benzofuranol N-methylcarbamate 98%) from Aldrich Chemistry with Bellstain Registry number 1428746, Product of USA.

Pellet feed for mice, aquadest as a carbofuran solvent, vitamin C, drinking water, husk as a base of the cage, ether, 10% formalin, alcohol. The tools used in this study were plastic

cages and wire mesh for experimental animal cages, drinking containers, sonde needles, test tubes, and 3 ml syringe.

Equipment used for the termination of mice is anesthetic jars, surgical scissors, scalpels, tweezers, and small pot bottles. Equipment for making liver histology preparations include glass objects, glass cover, automatic tissue processor, water bath, hot plate, microtome, and blade. Liver histology examination using an Olympus CX-41 camera and microscope.

Methods

The sample used in this study was the liver of the lactation period which was exposed to carbofuran. The number of samples used was 42 mice during lactation period. The mice used were divided into seven groups namely C, P1, P2, P3, P4, P5, and P6. Each group consisted of six replications.

Mice were given carbofuran, kebar grass, vitamin C orally using 1 ml tuberculin on the first day after giving birth until the 14th day. C: aquadest control group (0.5 ml/head/day), P1: carbofuran 1/4 LD₅₀ (0.0125 mg/day), P2: carbofuran 1/8 LD₅₀ (0.00625 mg/day), P3: carbofuran 1/4 LD₅₀ (0.0125 mg/day) +extract of kebar grass 3.375 mg in 0.2 ml, P4: carbofuran 1/8 LD₅₀ (0.00625 mg/day)+extract of kebar grass 3.375 mg in 0.2 ml, P5: 1/4 LD₅₀ carbofuran (0.0125 mg/day)+vitamin C 5 mg in 0.2 cc, and P6: carbofuran 1/8 LD₅₀ (0.00625 mg/day) +vitamin C 5 mg in 0.2 cc. All groups were given treatment for 14 days.

On the 15th day, mice were sacrificed for liver organ harvesting and histopathological preparation.

Mice are sacrificed by euthanasia. Mice were put into a jar that was given ether. Furthermore, abdominal cavity surgery is performed to remove the liver. Organs are placed into small pots containing 10% formalin buffer solution, then making histopathological preparations using HE staining.

Each mouse was made one liver preparation and each preparation was observed in five microscopic fields of view, namely in the four corners and the middle part of the preparation with a X 400 magnification.

Data analysis

Histopathological preparations of the liver were scored with degrees of liver damage with a score range of 0-5, depending on the extent of change and the degree of liver damage. The method used in granting this score uses the Knodell score method [14].

The results of scoring changes that occur in liver preparations are then analyzed using the Kruskal Wallis test followed by the Mann Whitney test with the Statistical Product and Service Solution (SPSS) application.

Results and Discussion

Microscopic observation of the liver histology preparations of mice (*Mus musculus*) lactation exposed to carbofuran then given kebar grass which is then compared with lactation during

mice given carbofuran and ascorbic acid or vitamin C. The results of scoring on mice's liver tissue showed degeneration, necrosis, and inflammation cell.

Degeneration

Degeneration observations were carried out microscopically using histopathological preparations with HE staining from the liver of mice (*Mus musculus*) during lactation.

The observation was carried out using a microscope with a X 400 magnification. The presence of degeneration is marked by the size of the cytoplasm of cells that are enlarged and cytoplasm that is cloudy because there are vacuoles (Table 2).

Table 1: Liver Damage Scoring.

Rate of Change	Score	Information
Inflammation Cells	0	No inflammation cell
	1	Inflammation cells occurs <25% of all fields view
	3	Inflammation cells occurs >25%-50% of all fields view
	5	Inflammation of inflammatory cells occurs >50% of all fields view
Degeneration	0	No degeneration cell
	1	Degeneration cells occurs <25% of all fields view
	3	Degeneration cells occurs >25%-50% of all field of view
	5	Degeneration cells occurs >50% of all fields of view
	0	No necrosis cells
Necrosis	1	Necrosis cells occurs < 25% of all fields view
	3	Necrosis cells occurs > 25% -50% of all fields view
	5	Necrosis cells occurs > 50% of all fields view

The results of the statistical analysis in Table 2 show that there are significant differences ($p < 0.05$) between the C group and the P1, P2, P3, P4, P5, and P6 groups. There were significant differences ($p < 0.05$) between groups P1 and groups P2, P3, P4, P5, and P6.

There were significant differences ($p < 0.05$) between groups P2 with groups P3, P4, P5, and P6. There were significant differences ($p < 0.05$) between groups P3 and groups P4, P5, and P6. There was a significant difference ($p < 0.05$) between groups P4 and groups P5 and P6.

Table 2: The scoring results of liver damage in mice exposed to lactation carbofuran.

Treatment	Degeneration Score (Mean \pm SD)	Necrosis Score (Mean \pm SD)	Inflammatory Cell Score (Mean \pm SD)
C	0.00 \pm 0.00 ^a	0.00 \pm 0.00 ^a	0.00 \pm 0.00 ^a
P1	4.73 \pm 0.23 ^f	4.87 \pm 0.23 ^f	4.20 \pm 0.00 ^f
P2	3.93 \pm 0.23 ^e	3.93 \pm 0.23 ^e	3.80 \pm 0.00 ^e
P3	1.27 \pm 0.23 ^c	2.06 \pm 0.46 ^c	0.80 \pm 0.20 ^c
P4	0.80 \pm 0.00 ^b	1.27 \pm 0.12 ^b	0.40 \pm 0.00 ^b
P5	2.73 \pm 0.61 ^d	3.40 \pm 0.80 ^{de}	2.73 \pm 0.92 ^{de}
P6	2.06 \pm 1.23 ^d	3.13 \pm 0.23 ^d	2.20 \pm 0.40 ^d

Note: C (aquadest control group), P1 (Carbofuran 1/4 LD₅₀ 0.0125 mg/day), P2 (carbofuran 1/8 LD₅₀ 0.00625 mg/day), P3 (extract of kebar grass 3.375 mg+carbofuran 1/4 LD₅₀ 0.0125 mg/day), P4 (extract of kebar grass 3.375 mg+carbofuran 1/8 LD₅₀ 0.00625 mg/day), P5 (vitamin C 5 mg+carbofuran 1/4 LD₅₀ 0.0125 mg/day), P6 (vitamin C 5 mg+carbofuran 1/8 LD₅₀ 0.00625 mg/day).

In this study administration of carbofuran with doses of 1/4 LD₅₀ and 1/8 LD₅₀ in lactation mice gave a significant difference ($p < 0.05$) and could increase hepatocyte degeneration in proportion to the increase in carbofuran dose given to lactation mice (Table 2).

Carbofuran can cause toxic effects that cause ROS. Excessive amounts of ROS can result in the formation of hydroxyl radicals that are very reactive and harmful to the body. Hydroxyl radicals disrupt the normal function of cells by causing reversible lesions in the form of degeneration.

This is consistent with research conducted by [15]. Which states an increase in ROS in excessive amounts is very dangerous to the body and can cause the formation of hydroxyl radicals that are very reactive.

In this study the administration of kebar grass to the lactation mice that were previously given carbofuran with doses of 1/4 LD₅₀ and 1/8 LD₅₀ can provide a significant difference ($p < 0.05$) and can reduce the degeneration of mouse hepatocytes (Table 2).

The rate of decrease in hepatocyte degeneration of mice produced by flavonoids is better for mice exposed using carbofuran at a dose of 1/8 LD₅₀ compared to mice exposed using carbofuran at a dose of 1/4 LD₅₀ due to lower toxicity. Charged grass contains flavonoids and is classified as a non-enzymatic antioxidant.

Citation: Luqman EM, Aulia Rakhman MA, Hidanah S, et al.. The potential of kebar grass (*Biophytum petersianum klotzsch*) on mice (*Mus musculus*) during lactation period on liver histopathological were exposed by carbofuran. *J RNA Genomics* 2021;17(6):1-7.

The antioxidant effect of flavonoids enhances the regeneration process by destroying free radicals, providing competitive substrates for unsaturated lipids in the membrane and accelerating the repair mechanism of damaged cell membranes so that the amount of hepatocyte degeneration decreases [16].

In this study the administration of ascorbic acid or vitamin C to the lactation mice that were previously given carbofuran at a dose of 1/4 LD50 and 1/8 LD50 can reduce degeneration of mouse hepatocytes even though there were no significant differences (Table 2).

The rate of decrease in hepatocyte degeneration of mice produced by ascorbic acid is better for mice exposed using carbofuran at a dose of 1/8 LD50 compared to mice exposed using carbofuran at a dose of 1/4 LD50 due to lower toxicity.

Ascorbic acid or vitamin C acts as an antioxidant causing a decrease in the amount of hepatocyte degeneration. Ascorbic acid can stabilize free radicals by completing the lack of electrons possessed by free radicals.

Ascorbic acid protects the substrate by substitution in the oxidation process and inhibits a chain reaction from the formation of free radicals that can cause oxidative stress [17].

Necrosis

Observation of necrosis was carried out microscopically using histopathological preparations with HE staining of the liver of mice (*Mus musculus*) during lactation.

The observation was carried out using a microscope with a X 400 magnification [18]. observation of necrosis is seen by the presence of nuclei that undergo picnosis, karyorrhexis and karyolysis (Table 2).

The results of the statistical analysis in Table 2 show that there are significant differences ($p < 0.05$) between the C group and the P1, P2, P3, P4, P5, and P6 groups. There were significant differences ($p < 0.05$) between groups P1 and groups P2, P3, P4, P5, and P6 [19].

In this study the administration of carbofuran with a dose of 1/4 LD50 and 1/8 LD50 in the lactation mice gave a significant difference ($p < 0.05$) and can increase hepatocyte necrosis in proportion to the increase in carbofuran dose given to the lactation mice (Table 2).

Necrosis is an advanced stage of degeneration that has passed the point of no return.

Necrosis is cell damage that causes tissue cell death while the individual is still alive [18].

Pathogenically the occurrence of necrosis is a continuation of degeneration which has reached the point of no return [18].

This is consistent with research conducted by [19].

Which states the level of toxicity of a compound in the body is caused by the period of exposure, method and dosage, and the level of immunity between individuals [20].




Figure 1. Comparison of microscopic features of lactating hepatocyte degeneration of mice (*Mus musculus*) during lactation with H.E staining, X 400. Arrows indicate cell degeneration.




Figure 2. Comparison of microscopic images of mice hepatocyte necrosis (*Mus musculus*) during lactation with H.E staining, X 400. Arrows indicate cell necrosis.




Figure 3. Comparison of microscopic images of inflammation cell in the liver of lactation mice (*Mus musculus*) with H.E staining, X 400. Arrows indicate infiltration of inflammatory cells.

In this study the administration of kebar grass to lactation mice that were previously given carbofuran at doses of 1/4 LD₅₀ and 1/8 LD₅₀ can provide a significant difference ($p < 0.05$) and can reduce mice hepatocyte necrosis (Table 2) [21].

The reduced level of hepatocyte necrosis of mice produced by flavonoids is better for mice exposed using carbofuran at a dose of 1/8 LD₅₀ compared to mice exposed using carbofuran at a dose of 1/4 LD₅₀ due to lower toxicity [22].

It is suspected that administration of kebar grass extract can restore hepatocyte function so that an increase in the number of cell necrosis can be prevented [23-25]. This is in accordance with research conducted by [20].

Which states that there is a decrease in the number of hepatocyte necrosis due to the presence of flavonoid content in kebar grass that works by suppressing the formation of free radicals or ROS through inhibition of enzymes, metal ion chelating involved in the production of free radicals and reducing free radicals (free radical scavengers) [26].

In this study the administration of ascorbic acid or vitamin C to the lactation mice that were previously given carbofuran at a dose of 1/4 LD₅₀ and 1/8 LD₅₀ can reduce mice hepatocyte necrosis even though there were no significant differences (Table 2).

Decreased levels of mice hepatocyte necrosis produced by ascorbic acid are better for mice exposed using carbofuran at a dose of 1/8 LD₅₀ compared to mice exposed using carbofuran at a dose of 1/4 LD₅₀ due to lower toxicity [27].

Ascorbic acid or vitamin C acts as an antioxidant that causes a decrease in the number of hepatocyte necrosis.

Ascorbic acid or vitamin C works by donating electrons so that it can prevent the formation of other compounds from the oxidation process by releasing a carbon chain [21].

Inflammation cells

Observation of inflammation cell was carried out microscopically using histopathological preparations with HE staining from the liver of mice (*Mus musculus*) during lactation [28].

The observation was carried out using a microscope with a X 400 magnification. Observation of inflammation cell was seen in the porta or Kiernan triangles (Table 2).

The results of the statistical analysis in Table 2 show that there are significant differences ($p < 0.05$) between the C group and the P1, P2, P3, P4, P5, and P6 groups. There were significant differences ($p < 0.05$) between groups P1 and groups P2, P3, P4, P5, and P6 [29]. There was a significant difference ($p < 0.05$) between P2 groups and P3, P4, and P6 groups.

Citation: Luqman EM, Aulia Rakhman MA, Hidanah S, et al.. The potential of kebar grass (*Biophytum petersianum klotzsch*) on mice (*Mus musculus*) during lactation period on liver histopathological were exposed by carbofuran. *J RNA Genomics* 2021;17(6):1-7.

There were significant differences ($p < 0.05$) between groups P3 and groups P4, P5 and P6. There was a significant difference ($p < 0.05$) between groups P4 and groups P5 and P6.

In this study the administration of carbofuran with a dose of $1/4 LD_{50}$ and $1/8 LD_{50}$ in the lactation mice gave a significant difference ($p < 0.05$) and can increase inflammation cell in the liver in proportion to the increase in carbofuran dose given to the lactation mice (Table 2) [30].

The presence of inflammatory cells in the tissue is also a response to injury by carbofuran in the form of membrane damage and necrosis.

These events due to oxidative stress can cause cellular protein damage and cause necrosis [22]. This is consistent with research conducted by [23].

Which states that inflammation cell is the body's defensive reaction to all forms that attack the body.

In this study the administration of kebar grass to lactation mice that were previously given carbofuran at doses of $1/4 LD_{50}$ and $1/8 LD_{50}$ can provide a significant difference ($p < 0.05$) and can reduce inflammation cell infiltration in the liver of mice (Table 2) [31].

The level of reduction in inflammation cell in the liver of mice produced by retinol and tocopherol was better for mice exposed using carbofuran at a dose of $1/8 LD_{50}$ compared to mice exposed using carbofuran with a dose of $1/4 LD_{50}$ due to lower toxicity.

There is a decrease in the amount of inflammation cell in the liver due to the presence of retinol and tocopherol content in the kebar grass.

Retinol or vitamin A as an antioxidant works by reacting with free radicals and free radicals become stable.

Tocopherol or vitamin E as a non-enzymatic antioxidant that collaborates with oxygen to destroy Reactive Oxygen Species (Sayuti, 2015; Widayati, 2012) [12].

In this study the administration of ascorbic acid or vitamin C to lactation mice that were previously given carbofuran at doses of $1/4 LD_{50}$ and $1/8 LD_{50}$ can reduce inflammation cell infiltration in the liver of mice, although there were no significant differences (Table 2).

The level of reduction in inflammation cell in the liver of mice produced by ascorbic acid is better for mice exposed using carbofuran at a dose of $1/8 LD_{50}$ compared to mice exposed using carbofuran at a dose of $1/4 LD_{50}$ due to lower toxicity. Ascorbic acid or vitamin C acts as an antioxidant that causes a reduction in the amount of inflammation cell in the liver.

Ascorbic acid or vitamin C functions as an antioxidant because it effectively captures free radicals, especially ROS or reactive oxygen compounds [24]. The administration of kebar grass extract is considered better than the administration of ascorbic acid or vitamins in treating degeneration, necrosis, and infiltration of inflammatory cells in mice during the lactation period caused by carbofuran with a dose of $1/4 LD_{50}$ or $1/8 LD_{50}$. This is due to the presence of flavonoids, retinol and tocopherol in kebar grass.

Polyphenol compounds such as flavonoids can inhibit oxidation reactions through radical scavenging mechanisms by donating an unpaired electron in free radicals so that the number of free radicals is reduced [13]. Retinol or vitamin A as an antioxidant works by reacting with free radicals and free radicals become stable. Tocopherol or vitamin E has properties that are easily oxidized so that they can protect other compounds from oxidation. As an antioxidant tocopherol can fight destructive oxygen, lipid peroxide, and free radicals and stop free radical chain reactions [25].

Conclusion

The administration of kebar grass extract is better in dealing with degeneration, necrosis, and infiltration of inflammatory cells in mice during lactation caused by administering carbofuran at a dose of $1/4 LD_{50}$ or $1/8 LD_{50}$ compared with the provision of vitamin C. The content of flavonoids, retinol, tocopherol, vitamins A and vitamin E in kebar grass can inhibit oxidation reactions.

Acknowledgements

The authors express sincere thanks to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia for funding the research, and the Dean of Faculty of Veterinary Medicine Universitas Airlangga for providing all necessary facilities and funds in conducting this research work.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Sayuti K, Rina Y. Natural and synthetic antioxidants. Andalas University Press. Padang. Sudarmo, S. 1991. Pesticide. Kanisius. Yogyakarta. 2015.
2. Jaiswal SK, Siddiqi NJ, Sharma B. Carbofuran induced oxidative stress in rat heart: Ameliorative effect of vitamin C. *Hindawi Publish Corp J*. 2013:1-10.
3. Indraningsih. Effect of carbamate insecticide use on animal health and its products. *Wartazoa J*. 2008;8:101-14.
4. Tannock J, Wessel CL. Determination of carbofuran residue and metabolites in plant material. *Pesticide Sci J*. 1981;12:228-34.
5. Kaur M, Shandir R. Comparative effect of acute and chronic karbofuran exposure on oxidative stress and drug

- metabolizing enzymes in liver. *Drug Chem Toxicol J*. 2006;29:415-21.
6. Gibson J. *Modern physiology and anatomy for nurses*. Edisi ke 2. EGC. Jakarta. 2003;2:534-6.
 7. Prayitno J. Effects of giving effervescent tablets combination of ethanol extracts of dewandaru leaves (*Eugenia uniflora L*) and sambiloto (*Andrographis paniculata* [Burm.f.] Ness) on liver function in glucose-ridden mice. [Thesis]. Fakultas Farmasi Universitas Muhammadiyah Surakarta. Surakarta. 2009.
 8. Chew B P, Park JS. Carotenoid action on the immune response. *Nutr J*. 2004;134:257-61.
 9. Brambilla D, Mancuso C, Scuderi MR, et al. The role of antioxidant supplement in immune system, neoplastic, and neurodegenerative disorders: A point of view for an assessment of the risk/benefit profile. *Nutr J*. 2008;7:29-38.
 10. Mansour T, Haikal M, Mossa AH, et al. Toxic effect of five insecticides on their mixture on male albino rats. *Egypt Soc Toxicol J*. 2008;39:85-94.
 11. Hudson BJ. *Food antioxidant* (Elsevier Applied Science Series). London. 1990.
 12. Santoso B, Kilmakossu A, Sambodo P. Effects of saponin from *biophytum petersianum klotzsch* on ruminal fermentation, microbial protein synthesis and nitrogen utilization in goats. *Anim Feed Sci Technol J*. 2007;137:58-68.
 13. Pokorny J, Yanishlieva N, Gordon M. *Antioxidant in food: Practical applications*. CRC Press. New York. 2001.
 14. Brunt ME. Grading and staging the histopathological lesions chronic hepatitis: The knodell histology activity index and beyond. *Hepato J*. 2000;31:241-6.
 15. Lobo V, Patil A, Phatak A, et al. Free radical, antioxidants, and functional foods: impact on human health. *Pharmacogn Rev*. 2010;4:118-26.
 16. Potential of vitamin E (α -tocopherol) against spermatogenic cells, diameter of seminiferous tubules, and levels of malondialdehyde (MDA) testes of mice (*Mus musculus*) induced 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). Tesis. Program studi ilmu biologi reproduksi fakultas sains dan Teknologi Universitas Airlangga. Surabaya. 2013.
 17. Christijanti W, Nur RU, Iswara A. Effect of provision of antioxidants vitamins c and e on spermatozoa quality of white rat exposed to allethrin. *Biol Educ J*. 2010;2:18-26.
 18. Pringgoutomo S, Sutisna H, Achmad T. *Pathology textbook I (General)*. Sagung Seto. Jakarta. 2002.
 19. Ndaong NA, Wijayanti AD, Widyarini S. Efek deltamethrin exposure to broilers on alanine aminotransferase enzyme activity, aspartate aminotransferase and overview of histopathology. *J Kajian Veter*. 2014;2:79-87.
 20. Halliwell B, Gutteridge JMC. *Free Radical in Biology and Medicine*. Oxford University Press. New York.
 21. Muhammad I. Antioxidant effects of vitamin c on male rats (*rattus norvegicus l*) due to cigarette smoke exposure. Thesis. Sekolah Pascasarjana Institut Pertanian Bogor. Bogor. 2009.
 22. Wahjuni S. *Malondialdehyde monograph precursor oxidative stress*. Udayana University Press. Denpasar. 2012:22-4.
 23. Yang B, Weiyang Z, Zhenzhen H, et al. Involvement of oxidative stress and inflammation in liver injury caused by perfluorooctanoic acid exposure in Mice. *Biomed Res Int*. 2014;2014:409837.
 24. Frei B. Reactive oxygen species and antioxidant vitamins: Mechanisms of action. *Am J Med*. 1994;97:5S-13S.
 25. Lamid A. Vitamin E as an antioxidant. *J Media Penelitian dan Pengembangan Kesehatan*. 1995;5:14-6.
 26. Arimbi, Azmijah A, Darsono R, et al. *General veterinary pathology textbook*. Edisi ke 2. Airlangga University press. Surabaya. 2015.
 27. Gupta RC, Goad JT, Kadel WL. Carbofuran-induced alterations (in vivo) in high-energy phosphates, creatine kinase (CK) and CK isoenzymes. *Arch Toxicol*. 1991;65:304-10.
 28. Kamboj SS, Kumar V, Kamboj A. Mitochondrial oxidative stress and dysfunction in rat brain induced by carbofuran exposure. *Cell Mole Neurobiol J*. 2008;28:961-9.
 29. Knodell RG, Ishak KG, Black WC, et al. Formulation and application of a numerical scoring system for assessing histological activity in asymptomatic chronic active hepatitis. *Hepato J*. 1981;1:431-5.
 30. Rai DK, Rizvi PK, Watal SI, et al. Carbofuran induced toxicity in rats protective role of vitamin C. *Exp Toxicol Pathol J*. 2009;61:531-5.
 31. Yanti EZ, Hermadi HA, Rejeki PS, et al. The increasing inflammatory cells, degeneration and duodenal necrosis on infant mice from their carbofuran exposed mothers. *J Int Dental Med Res*. 2020;13:785-90.

*Correspondence to

Epy Muhammad Luqman
 Departement of Veterinary Science,
 Universitas Airlangga Kampus C Unair Jalan Mulyorejo
 Surabaya
 Indonesia
 E-mail: epy-m-l@fkh.unair.ac.id