



Treatment of COVID-19 by Controlling the Activity of the Nuclear Factor-Kappa B

Mahmoud Saad Mohamed Elkhodary

Suez Canal University, Egypt

Abstract:

Heavy infection of the virus leads to overproduction of cytokines. The overproduction of cytokine (cytokines storms) is responsible for the critical cases and deaths of COVID-19. The nuclear factor kappa-B stimulates the expression of the genes, which is responsible for cytokines storm and RNA transcription. The COVID-19 virus can be controlled by inhibition of nuclear factor kappa-B. Nuclear factor kappa-B is controlled by inhibition of hydrogen peroxide and inhibitor kappa-B kinase enzyme.

Biography:

Mahmoud Saad Mohamed Elkhodary, Suez Canal University, Egypt

Recent Publications:

 How to Return the Death Programs of Cancer Cells to Work again and Cure Cancer within a Short Time?. Mahmoud Saad Mohamed El-Khodary, Sa-



har Ezeldien Hasal,, Wael A. Hassan, Maather M. El-Lamie, Ismail A. M. Eissa, Waleed F. Khalil, Salah M. Aly. DOI: 10.4236/cellbio.2019.82002 Jun. 30, 2019

Webinar on Novel COVID-19 and its impact on the Healthcare System | July 15, 2020 | Dubai, UAE

Citation: Mahmoud Saad Mohamed Elkhodary, Treatment of COVID-19 by Controlling the Activity of the Nuclear Factor-Kappa B, Novel Covid-19, 2020, July 15, Dubai, UAE