

## **Trichomoniasis in Nigeria: a review.**

**Omorodion Oriri Asemota\***

Industrial Safety and Environmental Technology Department, Petroleum Training Institute, Effurun, Nigeria

### **Abstract**

**Sexually Transmitted Parasitic Diseases (STPDs) are infectious diseases that are transmitted via unprotected sex which could potentially predispose men and women to different reproductive health issues. The data presented in this report shows that, trichomoniasis caused by *Trichomonas vaginalis* (TV) remain the only reported sexually-transmitted parasitic disease in Nigeria. Also, of the 36 states in the country, only 26 states including the Federal Capital Territory (FCT), Abuja have partly evaluated the prevalence of the disease. Furthermore, the prevalence data on the burden of TV infections in Nigeria in form of Years Lived with Disability (YLD), the distribution of the infection in the six geopolitical zones and the prevalence over the years are presented. Notably, YLD for TV infection was highest in the South-South; while the factors possibly impacting on the burden of TV in Nigeria are highlighted.**

**Keywords:** Sexually transmitted parasitic diseases (STPDs), Trichomoniasis, Nigeria.

*Accepted on April 18, 2018*

### **Introduction**

Sexually Transmitted Diseases (STDs) being transmitted by sex through different routes are amongst the commonest communicable diseases globally with more than 1 million new infection daily [1]. The burden of STDs largely occurs in developing countries [2]; with the most vulnerable, disproportionately affected by other health and social-economic issues. In developing countries, STDs are among the leading causes of Disability-Adjusted Life Years (DALYs) lost for women of reproductive age [3].

Parasitic STDs include trichomoniasis, amoebiasis and giardiasis [4,5]. Infection with Sexually Transmitted Parasitic Diseases (STPDs) has resulted in debilitation or anatomic deformities that make sex impossible as a result of direct damage to the male and female reproductive organs including impairing fertility via the inhibition of gamete production [6].

*Trichomonas vaginalis* is the most common STPDs in Nigeria; and studies on STPDs remain relatively scanty [7-10]. In addition, knowledge about *T. vaginalis* and trichomoniasis is hugely lacking; and this identified gap may have impacted on the burden of infection.

This review article thus present prevalence data for Nigeria by states; and highlight the gaps in information regarding various aspects of the disease. Further, the implication of the relatively lack of awareness about *T. vaginalis* infection were noted and discussed.

### **Prevalence and Distribution of STPDs**

In Nigeria, amongst the vast array of endemic parasitic diseases, only trichomoniasis is sexually transmitted. There are other parasites that could be transmitted via oral-anal and oral-genital routes and as such can elicit other forms of sexually transmitted diseases. Of the 36 states and Abuja, *T. vaginalis* studies could only be accessed for twenty-six (26) states (Table 1). These studies spanned through the different geo-political zones (South-East, South-South, South-West, North-West, North-East and North-North) of the country. In spite of this, there is still limited/lack of literature on the prevalence of *T. vaginalis* infection in some other states. From available data, the prevalence across the country ranged between 0-51.8%. Majority of these studies were carried out on pregnant women in the different geo-political zones, and samples were mostly vaginal swab (Table 1).

**Table 1.** Prevalence of trichomoniasis across zones and states.

Zone	State	Participants	Number tested	Prevalence (%)	Sample (s) used	References
South-East	Ebonyi	HIV/AIDS patients	970	36.6	Urine	[11]
	Anambra/Enugu	Male adolescents	361	25.2	Urine, urethral smears, semen	[12]

	Ebonyi	HIV-seropositive women	250	24.4	Urine, HVS	[13]
	Abia	Women	600	18.67	Urine, HVS	[7]
	Anambra	Women between 20 years and above	200	17.5	HVS	[14]
	Abia	Pregnant women	200	13	HVS	[15]
	Anambra	Rural women	100	15	HVS	[16]
	Anambra	Women	1440	21.53	Urine, HVS	[17]
	Anambra	Females	100	2.3	Blood, HVS	[18]
	Ebonyi	Pregnant women	688	12.9	Vaginal swab	[19]
	Ebonyi	Antenatal patients	200	27.5	HVS	[20]
	Ebonyi	Single and married pregnant women	1025	12.3	Vaginal swab	[21]
	Imo	Female/adult out patients	1241	10.2	HVS	[22]
	Imo	Males	178	3.4	US	[22]
	Imo	Males and females	5883	1.3	US	[22]
South-South	Edo	Pregnant women	150	0	Urine	[23]
	Cross-River	Pregnant women	562	5.2	HVS	[24]
	Bayelsa	Females patients with vaginal complaints	1240	6.5	HVS, ECS	[25]
	Edo	Pregnant/non-pregnant women	300	2.8	HVS	[26]
	Akwa-Ibom	Pregnant women	260	17.7	HVS	[10]
	Rivers	Women attending antenatal, gynaecology, family planning and sexually transmitted disease clinics	220	4.5	HVS	[27]
	Rivers	Women from urban and sub-urban areas	1431	0.63	HVS	[28]
	Edo	Adolescent girls in secondary schools	272	9.2	HVS	[29]
	Akwa-Ibom	HIV-seropositive people	150	16.2	HVS, urine	[30]
	Akwa-Ibom	Pregnant women	560	44.5	HVS, ECS	[31]
	Delta	Women at Effurun	100	6	HVS	[32]
	Delta	Secondary school girls, teachers, undergraduates and petty traders	240	22.5	Urine, HVS	[33]
South-West	Ogun	HIV-infected and non-infected women	100	5	Urine, HVS	[34]
	Osun	Female patients	310	8.1	HVS, ECS, US	[35]
	Oyo	Pregnant women	100	6	HVS	[36]
	Osun	Pregnant women	100	2	HVS	[37]
	Ogun	Pregnant women	300	10.3	HVS	[38]
	Ogun	Pregnant women	200	20	HVS, Urine	[39]

	Lagos	Pregnant women	198	18.7	HVS	[40]
	Ibadan	ARFH attendees	200	1.5	Blood, HVS, ECS	[41]
	Oyo	Pregnancy ante-natal patients	150	25.3	HVS	[42]
	Lagos	Women (outpatients)	544	3.3	HVS	[43]
	Ondo	Seropositive and sero negative pregnant women	240	50 and 22	HVS	[44]
	Ondo	Seropositive and sero negative males and females	100	0 and 6	Urine and HVS	[45]
	Ibadan	Sexual partners of women with confirmed diagnosis of TV	44	45.5	Urethral swab	[46]
	Ogun	Female students Babcock University	200	12.5	HVS	[47]
	Lagos	Patients with virginal discharge	200	74.5	HVS	[48]
	Ogun	University male female students in redemption camp	60	1.67 and 3.3	urine	[43]
North-West	Kaduna	Pregnant women	120	19.2	HVS, Blood	[49]
	Kebbi	Female patients	220	4.1	Urine, HVS	[50]
	Jigawa	Female patients	150	1.3	HVS	[51]
	Kano	Females with abnormal vaginal discharges	2000	1.5	ECS, HVS	[52]
	Kaduna	Females living in Palladan	150	13.67	HVS	[53]
	Kaduna	HIV-seropositive antenatal clients	102	5.9	HVS	[54]
North-East	Borno	Female IDPs	200	20.5	HVS	[8]
	Borno	Pregnant women	919	10.99	HVS	[55]
	Adamawa	Health males, females, antenatal women, and sick males and females without any history of Trichomoniasis	1520	2.6	Blood and genital swab	[56]
North-Central	Plateau	HIV-1 infected women	168	51.8	HVS	[57]
	Nasarawa	Pregnant women	960	7.5	Urine, HVS	[58]
	Nasarawa	Female patients	200	8.5	Vaginal discharge	[59]
	Kwara	Pregnant women	250	20	HVS, Urine	[60]
	Plateau	Female	150	4	HVS	[61]
	FCT, Abuja	Primary school pupils	402	3.73	Urine, stool	[62]
	Kwara	Pregnant women	230	4.7		[63]
	Kwara	HIV-seropositive antenatal clients	160	5.6	ECS, HVS	[64]
	Benue	Females patients	200	10.5	HVS, Urine	[65]
	Benue	Male and female patients	400	12	Urine	[66]

Nasarawa	Female patients	100	12	HVS	[67]
Nasarawa	Male and female patients	1050	15.71	Urine, HVS	[68]
FCT, Abuja	Male and female patients	286	15	Blood, ECS, HVS, Urine, Urethral swabs, Urethral discharge	[69]
Nasarawa	Female students of Bingham University	200	4.5	Low vaginal swabs	[70]
Plateau	HIV positive and negative patients	700	17	HVS	[71]

HVS: High Vaginal Swab; ECS: Endocervical Swap; US: Urethral Sample; HIV: Human Immunodeficiency Virus; AIDS: Acquired Immune Deficiency Syndrome.

The prevalence of *T. vaginalis* infection is relatively higher in sexually active women [9]. In males, *T. vaginalis* infection is generally trivial or asymptomatic. Asymptomatic carriers can serve as vectors for the disease, making it important to treat male partners. The parasite resides in the female's lower genital tract and the male urethra and prostate.

Reports from Nigeria suggest that, trichomoniasis could be higher in urban areas than in the rural communities [72,73]. For example, in a study carried out by [74], a higher prevalence of *T. vaginalis* was documented for subjects that resided in urban areas (57.70%) than those that lived in rural communities (39.16%). Also, prevalence was highest amongst the sexually active group of age 11-45 years old [7].

Amongst pregnant women, *T. vaginalis* is generally high [9,39]. In Zaria, pregnant women between ages 16 to 25 were reported to be *T. vaginalis* positive with prevalence of 53.57% [43]. In Lagos, South-West, Nigeria, *T. vaginalis* prevalence of 1.8% was recorded for pregnant women between 21 to 30 years old. Similarly, in Abeokuta, South-West Nigeria, pregnant women between the age 20 and 30 had prevalence of 21.3%. Meanwhile in Anambra, South-East Nigeria, non-pregnant women had higher rate of infection (17.8%) than pregnant ones (16.7%) [14]. Furthermore, a study in Maiduguri showed a high rate of 20.8% infection among non-pregnant women [8].

In relation to marital status, a prevalence of 2.9% was recorded among Lagos married women compared to 0.4% prevalence in unmarried women [43]. Amadi and Nwagbo [7] reported 19.72% for single women in Abia, South-East, Nigeria with 21.6% and 11% among Abeokuta and Maiduguri married women respectively [8,38].

Data have shown that HIV could escalate infection with *T. vaginalis* [61,75,76]. Isiaka-Lawal et al. [64] observed that the prevalence of trichomoniasis was higher in HIV infected women than their uninfected counterparts in north-central Nigeria, while in Lagos, a prevalence of 35.8% were noted in HIV positives [77].

Available data had revealed an increased risk of *T. vaginalis* infection in persons with poor personal hygiene and low socio-economic status [78]. Levels of education have been linked to prevalence rate. For instance, non-formal educated women had

an infection rate of 22.3% while tertiary educated women had only 1.0% [8]. Also, Usanga et al. [24] reported that women with primary school education had 6.4% prevalence rate. In addition, poverty, unemployment, violence against women and children [17,24], and young age could raise the risk associated with acquiring *T. vaginalis* infection [63,79].

## Diagnosis

Diagnostic methods available for trichomoniasis are wet mount preparation, staining methods, culture in laboratory medium, and molecular methods. Historically, detection of the parasite is made possible by examination of urine and High Vaginal Swab (HVS) in a drop of saline or trichomonas diluents for the characteristic wobbling and rotating motion. Amadi and Nwagbo [7] reported that either urine sample or vaginal swab is insufficient for proper diagnosis of *T. vaginalis* infection and have suggested that for better results both urine and vaginal swab should be used. Diagnosis of trichomoniasis has relied mostly on wet mount demonstration and staining of the parasite in the laboratory with success rate of between 20 and 80% [80]. A combination of cultural method with microscopic wet mount demonstration is now the acceptable procedure for effective diagnosis [81].

New molecular diagnostic tests with improved sensitivity have been developed in response to the increasing recognition by stakeholders of the importance of this wide-spread STI. Thus the detection of *T. vaginalis*, including rapid antigen detection and nucleic acid amplification tests, has significantly improved the quality of diagnostics for trichomoniasis, particularly in women [82]. In America, Guillermo et al. [83] reported 97% sensitivity for molecular amplification diagnostic methods, as against 70 and 36% sensitivities for culture and wet preparations respectively. Also, studies of genetic typing of the parasites have been reported to be more sensitive than other methods [84,85]. However, in Nigeria there is no report of the application molecular diagnosis tools on *T. vaginalis* infection, which may influence the prevalence data as presently reported from less reliable tests.

## DALYs and Infection Burden

Evaluation of the disease burden is essential in determining the cost-effectiveness of control so as to ensure that control

programmes are focused appropriately. Disability Adjusted Life Year (DALYs) is the trending metric measurement to assess disease burden [86]. DALYs include years of life lost due to mortality (YLLs) and years lived with disability (YLDs). Most *T. vaginalis* survey is mainly on women, but reports of infection burden measured in DALYs are unavailable. Here we have estimated value of YLD based on available prevalence data and morbidity of the infection [87]. YLD due to TV in this review was estimated using the formula below [88].

$$YLD = P \times DW$$

Where, *P*=number of prevalent cases, *DW*=Disability Weight.

Here, prevalence was used rather than incidence [88], because in Nigeria, surveys are widely reported in prevalence (Table 2). *DW* (0.027) of infection which is represented on a scale of 0-1 (0=perfect health; 1=death), was adopted [89]. YLDs due to TV, according to the geopolitical zones are shown in Table 2. This review shows that the YLD from *T. vaginalis* prevalence data in Nigeria ranged from 21 to 63 with the highest and least disease burden recorded in South-South and North-West respectively.

**Table 2.** *T. vaginalis* mean prevalence and estimated YLD according to geopolitical zones (1984-2017).

Region	Mean ± SD (%)	YLD /10 000
South-East	16.12 ± 9.9	44
South-South	23.3 ± 16.3	63
South-West	17.3 ± 20.2	47
North-West	7.6 ± 7.3	21
North-East	11.4 ± 9.0	31
North-Central	12.5 ± 12.4	34

## Control

Control and elimination may depend largely on extensive proper sex education, especially for the adolescent and youths [7,8]. Also suggested are adequate treatment of both spouses, implementation of effective screening programmes and treatment at no cost [8]. For improved health, adequate personal hygiene, avoidance of promiscuity, improved education of women on safe sex and the need to know partners' STI status are advocated [14,77]. It is recommended that routine STIs screening in sexually active patients especially among the young and singles should be incorporated into hospital care [41]. This is needed to prevent transmission of the parasite, because some infected women and most infected men show no signs of the disease like liquid discharge from the vagina or penis, irritation while urinating and genital itching. So this YLD information is vital to focusing efforts on getting women tested and treated Based on the results from this study, trichomoniasis control in Nigeria should especially be of high priority in the South-South.

## Conclusion

Data available in this review have clearly outlined the prevalence of trichomoniasis in Nigeria, with the South-South having the highest disease burden. The dearth of information and very poor awareness of this STDs among the populace may have impacted on the burden of infection. Thus, increased awareness on the parasite and the disease at all levels is highly recommended. Also, ensuring that health workers include *T. vaginalis* as part of a routine procedure in the course of screening for STDs is thus advocated. Furthermore, providing better sensitive methods for diagnosis is the way to go as this would ensure near-accurate information on the prevalence of the disease in any given area in Nigeria.

## References

1. World Health Organization Report. Fighting disease, Fostering development. World Health Organization, Geneva, Switzerland 1996; 133.
2. WHO. Prevention and control of sexually transmitted infections: Draft global strategy. Ref Type: Generic 2006.
3. World Health Organization. Global prevalence and incidence of selected curable sexually transmitted infections: overview and estimates. Geneva, Switzerland 2001.
4. Okwute LO. A review of sexually transmitted diseases (STDs) of parasitic origin: The case of giardiasis. Afr J Biotechnol 2008; 7: 4979-4981.
5. Levine GI. Sexually transmitted parasitic diseases. Primary Care 1991; 18: 101-128.
6. Okafor CF, Omudu AE. Parasitic diseases and sexual disability: A critical review of some parasitic diseases with serious repercussions. Anim Res Int 2005; 2: 255-260.
7. Amadi ANC, Nwagbo AK. Trichomonas vaginalis infection among women in Ikwuano, Abia State, Nigeria. J Appl Sci Env Manag 2013; 17: 389-393.
8. Hamafyelto HS, Ikeh IE. Prevalence of Trichomonas vaginalis infection among female internally displaced persons in Maiduguri, Nigeria. Int J Trop Dis Health 2017; 27: 1-7.
9. Jatau ED, Olonitola OS, Olayinka AT. Prevalence of Trichomonas infection among women attending antenatal clinics in Zaria, Nigeria. Ann Afr Med 2006; 5: 178-181.
10. Opara K, Udoidiung N, Atting I, Basse E, Okon O, Nwabueze A. Risk factors for vaginal trichomoniasis among women in Uyo, Nigeria. Intern J Health 2008; 9: 1-5.
11. Nweze EI, Mouneke GN. Trichomonas vaginalis in HIV/AIDS subjects in Nigeria. Asian Pac J Trop Dis 2011; 4: 282-286.
12. Onyemelukwe NF, Nwokocha ARC. Trichomonas vaginalis infection in male adolescents in parts of South Eastern Nigeria. IOSR J Dent Med Sci 2013; 12: 52-53.
13. Uneke CJ, Alo MN, Ogbu O, Ugwuoru DC. Trichomonas vaginalis infection in human immunodeficiency virus-

- seropositive Nigerian women: The public health significance. *Onl J Health Allied Sci* 2007; 6: 1-7.
14. Iwueze MO, Ezeanyanwu LN, Okafor FC, Nwaorgu OC, Ukibe SC. Prevalence of *Trichomonas vaginalis* infection among women attending hospitals/health centres in Onitsha community, Onitsha North Local Government area of Anambra State. *Bioscientist* 2014; 2: 54-64.
  15. Kanu AM, Ihekumere I, Kalu EJ. Frequency distribution of Trichomoniasis in pregnant women in Aba, South East, Nigeria. *Univ J Microbiol Res* 2015; 3: 53-55.
  16. Onyido AE, Umeanaeto PU, Irikannu KC, Ekwunife CA, Ezeanya LC, Nwangwu UC, Ugha CN, Obiechina IO. Prevalence of *Trichomonas vaginalis* among the rural women of Ekwulumili community Anambra state, Southeastern Nigeria. *Nat Sci* 2014; 12: 129-134.
  17. Ulogu IO, Obiajuru IO, Ekejindu IM. Prevalence of trichomoniasis amongst women in Nnewi, Anambra state, Nigeria. *Niger J Parasitol* 2007; 28: 6-10.
  18. Orji NM. Reproductive tract infections among females in Ihiala Local Government area, Anambra State, Nigeria. *Int J Sci Eng Appl Sci* 2015; 1: 22-28.
  19. Okonkwo EC, Iroha IR, Onwa NC, Nworie O, Oji Egwu A. *Trichomonas vaginalis* associated with adverse pregnancy outcomes: implications for maternal health care delivery system in South Eastern Nigeria. *British J Med Med Res* 2012; 2: 568-574.
  20. Sunday-Adeoye IM, Adeoye JOK, Odidika UJU, Okonta P. The prevalence of *Trichomonas vaginalis* and *Candida albicans* infection in the lower genital tracts of antenatal patients in Abakaliki, Southeastern Nigeria. *Nepal J Obstetr Gynaecol* 2009; 4: 11-14.
  21. Okonkwo EC, Amadi ES, Idioha JC, Nworie O, Nwuzo AC, Onwe JC. Prevalence of *Trichomonas vaginalis* among pregnant women in Abakaliki, Ebonyi State. *Int J Curr Res* 2010; 11: 11-15.
  22. Acholonu ADW. Trichomoniasis: A little recognized sexually transmitted disease but with grave consequences. *Publ Lect Nigeria Acad Sci Lagos* 1998.
  23. Ochei KC, Obeagu EI, Ugwu GU, George CN. Prevalence of *Trichomonas vaginalis* among pregnant women attending hospital in Irrua Specialist Teaching Hospital in Edo State, Nigeria. *IOSR J Dent Med Sci* 2014; 13: 79-82.
  24. Usanga V, Abia-Bassey L, Inyang-Etoh P, Udoh S, Ani F, Archibong E. *Trichomonas vaginalis* infection among pregnant women in Calabar, Cross River State, Nigeria. *Intern J Gynecol Obstetr* 2009; 14: 1-7.
  25. Pondie K, Jeremiah I, Lawani E, Nsikak E. The prevalence of symptomatic vulvo-vaginal candidiasis and *Trichomonas vaginalis* infection and associated risk factors among women in the Niger Delta Region of Nigeria. *Int STD Res Rev* 2017; 5: 1-10.
  26. Akinbo FO, Mokobia CN, Ande AB. Prevalence of trichomoniasis among pregnant women in Benin City. *Sahel Med J* 2017; 20: 67-71.
  27. Olorode OA, Mark OO, Ezenobi NO. Urinogenital trichomoniasis in women in relation to candidiasis and gonorrhoea in University of Port-Harcourt Teaching Hospital. *Afr J Microbiol Res* 2014; 8: 2482-2485.
  28. Abah AE. *Trichomonas vaginalis* infection in a typical urban and a sub-urban area of Rivers state, Nigeria. *Asian J Med Health* 2017; 6: 1-6.
  29. Akinbo FO, Oronsaye IS. *Trichomonas vaginalis* infection among adolescent girls in some secondary schools in Benin City, Edo State, Nigeria. *Afr J Clin Exp Microbiol* 2017; 18: 223-229.
  30. Akpan UI. The prevalence of *Trichomonas vaginalis* in Uyo Local Government area of Akwa Ibom State, Nigeria. *Int J Mod Biol Med* 2013; 4: 134-139.
  31. Ekanem EI, Ekott M, Udo AE, Efiok EE, Inyang-Out A. Prevalence of sexually transmitted diseases in pregnant women in Ikot Epene, a rural community in Akwa Ibom State, Nigeria. *Op J Obstetr Gynaecol* 2012; 2: 49-55.
  32. Okoko FJ. Prevalence of Trichomoniasis among women at Effurun metropolis, Delta State, Nigeria. *Continent J Biol Sci* 2011; 4: 45-48.
  33. Wokem GN, Ndukwu CB. Re-evaluation of vulvovaginal trichomoniasis among women in Niger Delta region, Nigeria. *Glob J Pure Appl Sci* 2015; 21: 13-19.
  34. Oyewole IO, Anyasor GN, Michael-Chikezie EC. Prevalence of STI pathogens in HIV-infected and Non-infected women: implications for acquisition and transmission of HIV in Nigeria. *Asian J Med Sci* 2010; 2: 163-166.
  35. Olaniran O, Osevwe AJ, Balogun IA, Hassan-Olajokun RE, Oyetoke OO. The prevalence of *Trichomonas vaginalis* in Wesley Guide Hospital, Ilesha, Nigeria. *Int J Microbiol Res Rev* 2017; 6: 345-350.
  36. Donbraye E, Donbraye-Emmanuel OOB, Okonko IO, Okedeji IO, Alli JA, Nwanze JC. Detection and prevalence of *Trichomonas vaginalis* among pregnant women in Ibadan, South-Western Nigeria. *W Appl Sci J* 2010; 11: 1512-1517.
  37. Olowe OA, Makanjuola OB, Olowe R, Adekanle DA. Prevalence of vulvovaginal candidiasis, trichomoniasis and bacterial vaginosis among pregnant women receiving antenatal care in South Western Nigeria. *Eur J Microbiol Immunol* 2014; 4: 193-197.
  38. Etuketu MI, Mogaji OH, Alabi MO, Adeniran AA, Oluwole SA, Ekpo FU. Prevalence and risk factors of *Trichomonas vaginalis* infection among pregnant women receiving antenatal care in Abeokuta, Nigeria. *Afr J Infect Dis* 2015; 9: 51-56.
  39. Ojurongbe O, Taiwo BO, Dina BO, Sina-Agbaje OR, Bolaji OS, Adeyeba AO. Prevalence of *Trichomonas vaginalis* infection among pregnant women in Abeokuta, Nigeria. *Sierra Leone J Biomed Res* 2010; 2: 82-86.
  40. Oyetunde TO, Olamide F, Ifeoluwa TO. *Trichomonas vaginalis* infection in Nigerian pregnant women and risk factors associated with sexually transmitted infections *Int J STD AIDS* 2015; 27: 1187-1193.

41. Okonko IO, Okerentugba PO, Adejuwon AO, Onoh CC. Prevalence of sexually transmitted infections (STIs) among attendees of Lead City University Medical Centre in Ibadan, South-Western, Nigeria. *Arc Appl Sci Res* 2012; 4: 980-987.
42. Bolaji OS, Adejare FO, Adeyeba OA, Ojurongbe O. Comparison of methods of diagnosis of Trichomoniasis in pregnancy among ante-natal patients in some parts of Oyo state. *Int J Pharm Med Biol Sci* 2013; 2: 5-12.
43. Adeoye GO, Akande AH. Epidemiology of Trichomonas vaginalis among women in Lagos metropolis, Nigeria. *Pak J Biol Sci* 2007; 10: 2198-2201.
44. Ebhodaghe BI, Ako-Nai KA, Aderoba AK, Anderson WA, Kassim OO. Co-infections of human immunodeficiency virus and sexually transmitted infections among hiv seropositive pregnant women at healthcare centres in Akure, Southwestern Nigeria. *Int Arc Med* 2016; 9: 1-8.
45. Olorunfemi OB, Ijabadeniyi AO. Prevalence of Trichomonas vaginalis in urogenital tract infection in Akure South Local Government area, Ondo state. *Ife J Sci* 2005; 7: 159-161.
46. Bakare RA. Prevalence of Trichomonas vaginalis among the sexual partners of women with Trichomoniasis in Ibadan, Nigeria. *Afr J Clin Exp Microbiol* 2003; 4: 107-115.
47. Senchi JH, Samson ES, Olumuyiwa A, Elejo IG, Oluchi OG, Olusola OA. The prevalence of Trichomonas vaginalis infection and associated risk factors among undergraduate female students of Babcock University, Ilishan-Remo, Ogun state, Nigeria. *Int STD Res Rev* 2017; 6: 1-13.
48. Anorlu RI, Fagbenro-Beyioku AF, Fagorala T, Abudu OO, Galadanci HS. Prevalence of Trichomonas vaginalis in patients with vaginal discharge in Lagos, Nigeria. *Niger Postgrad Med J* 2001; 8: 183-186.
49. Avidime S, Sulayman HU, Adesiyun AG. Prevalence of Trichomonas vaginalis and HIV co-infection among asymptomatic pregnant women in Zaria, Northern Nigeria. *J Health Res Rev* 2017; 1: 49-53.
50. Abubakar A, Arzika S, Aliyu F, Umar H, Abubakar HM. Detection and prevalence of Trichomonas vaginalis among women in Birnin Kebbi, North West, Nigeria. *Int J Res Find Eng Sci Technol* 2013; 1: 23-27.
51. Ahmad MM, Yahaya G, Shuaibu I, Abdullahi II. Urinary tract infections among female patients attending Barnin-Kudu General Hospital, Jigawa, Nigeria. *Dutse J Pure Appl Sci* 2016; 2: 166-170.
52. Nwadioha S, Egesie JO, Emejuo H, Iheanacho E. Prevalence of pathogens of abnormal vaginal discharges in a Nigerian Tertiary Hospital. *Asian Pac JTrop Med* 2010; 3: 483-485.
53. Akafyi DE, Oko JO, Umar M, Obafemi A, Michael R. Prevalence of bacterial, Trichomonas and Candidal Vaginosis among females in Angwan-Fulani, Palladan in Zaria, Nigeria. *J Appl Life Sci Int* 2016; 5: 1-6.
54. Isaac N, Onwuhafua PI, Oguntayo AO, Opaluwa AS. Routine screening for Trichomonas vaginalis among human immunodeficiency virus-seropositive antenatal clients in Zaria: A necessity or option? *Trop J Obstetr Gynaecol* 2016; 33: 322-326.
55. Mairiga AG, Balla HJ, Ahmad MI. Prevalence of Trichomonas vaginalis infections among antenatal clients in Maiduguri, Nigeria. *Int J Biol Med Res* 2011; 2: 998-1002.
56. Abdulazeez A, Alo E, Naphthali R. Urino-genital trichomoniasis and human immune-deficiency virus concurrent infection in Adamawa state, Nigeria. *Intern J Paras Dis* 2007; 3: 1-6.
57. Anejo-Okopi AJ, Simji G, Okon EB, Ejeliogu UE, Okojokwu OJ, Okechalu J, Mafuyai S, Efekamaraye E, Isa SE. Prevalence and factors associated with Trichomonas vaginalis among HIV-1 positive women attending tertiary hospital in Jos, Nigeria. *Saudi J Pathol Microbiol* 2017; 2: 269-275.
58. Abioye JOK, Ihuma JO, Abdullahi DK, Kamoh N, Awari YS, Gemma K, Babatunde SK. Trichomonas vaginalis among pregnant women attending New Nyanya General Hospital, Nasarawa State, Nigeria. *Int J Sci Nat* 2014; 5: 437-439.
59. Yako AB, Maria FI, Hassan SC, Chassed G, Katuka G. Trichomonas vaginalis: the most prevalent sexually transmitted infection (STI) among female patients, a cause for medical care at Nagari Hospital (NH), Keffi, Nasarawa State, Nigeria. *W Sci Res J* 2014; 2: 67-70.
60. Bolaji OS, Ajadi TS, Bello SA, Ojurongbe O, Adeyeba OA. Prevalence of Trichomonas vaginalis among pregnant women from selected hospitals in Ilorin metropolis. *J Pharm Sci Innov* 2015; 4: 108-111.
61. Okojokwu OJ, Akpakpan EE, Kolawole OT, Ndubuisi JC, Okopi JA. Epidemiology of Trichomonas vaginalis infection among women in Jos metropolis, Nigeria. *Biomed Nurs* 2015; 1: 7-11.
62. Olarenwaju CA. Incidence of trichomoniasis among primary school pupils in Gwagwalada, FCT, Nigeria. *J Life Sci Technol* 2016; 4: 40-43.
63. Aboyeji AP, Nwabuisi C. Prevalence of sexually transmitted diseases among pregnant women in Ilorin, Nigeria. *J Obstetr Gynaecol* 2003; 23: 637-639.
64. Isiaka-Lawal SA, Nwabuisi C, Fakeye O, Saidu R, Adesina KT, Ijaiya MA, Jimoh AA, Omokanye LO. Pattern of sexually transmitted infections in human immunodeficiency virus positive women attending antenatal clinics in North-Central Nigeria. *Sahel Med J* 2014; 17: 145-150.
65. Okoh ME, Igbaaka IM. Prevalence of Trichomonas vaginalis among females visiting some selected hospitals in Makurdi, Benue State Nigeria. *Int J Microbiol Biotechnol* 2017; 2: 43-47.
66. Gberikon GM, Aguoru CU, Yandev D. Prevalence of Schistosoma haematobium and Trichomonas vaginalis in relation to age distribution and marital status among

- patients attending four selected hospitals in Gboko, Benue state of Nigeria. *Int J Pure Appl Sci Technol* 2015; 29: 26-30.
67. Hussaini Y, Adegda JA, Abdulkadir A, Suleiman A, Murtala SE, Jibrin Y, Yakubu Y. Prevalence of *Trichomonas vaginalis* among female patients attending the General Hospital Nasarawa, Nasarawa State. *Cell Biol* 2017; 5: 8-11.
  68. Alaku IA, Abdullahi AG, Attah DD. Prevalence of human *Trichomonas vaginalis* infections in some parts of Nasarawa State, Nigeria. *J Med Biol Sci* 2014; 4: 8-16.
  69. Raimi MO, Ochayi EO. Assessment of the rate of sexually transmitted diseases in Kubwa F.C.T, Abuja, Nigeria. *Sci J Publ Health* 2017; 5: 365-376.
  70. Adogo LY, Oyewole EA, Anyanwu NCJ, Omebije PE. Prevalence and correlates of *Gardnerella vaginalis* and *Trichomonas vaginalis* among female students in Bingham University. *Int J Trop Dis Health* 2016; 20: 1-8.
  71. Nwadioha SI, Bako IA, Onwuezobe I, Egah DZ. Vaginal trichomonas among HIV patients attending primary health care centres, Jos, Nigeria. *Asian Pac J Trop Dis* 2012; 2: 337-341.
  72. Njoku AJ, Obiajuru IOC, Njoku CJ, Nwokoro EA, Uwaezuoke IC, Anosike IC. Prevalence of *Trichomonas vaginalis* infection among students of tertiary institutions in Imo state, Nigeria. *Niger J Parasitol* 2000; 21: 83-94.
  73. Obiajuru IOC. Prevalence of *Trichomonas vaginalis* in Owerri, Imo state. M.Sc. Thesis, Imo State University, Owerri 2000.
  74. Obiajuru IOC, Ogbulie JN. Prevalence of sexually transmitted diseases in three zones of Imo state, Nigeria. *Niger J Microbiol* 2007; 21: 1491-1498.
  75. Sorvillo F, Kerndt P. *Trichomonas vaginalis* and amplification of HIV-1 transmission. *Lancet* 1998; 351: 213-214.
  76. Laga M, Manoka A, Kivuvu M. Non-ulcerative sexually transmitted diseases as risk factors for HIV-1 transmission in women: results from a cohort study. *AIDS* 1993; 7: 95-102.
  77. Oyetunde TS, Chelsea NE. Frequency of Human Immunodeficiency Virus (HIV) in *Trichomonas vaginalis* infected women in Badagry, Lagos, Nigeria. *J Reprod Infertil* 2016; 17: 61-63.
  78. Thomas CQ, King KH. *Trichomoniasis*. Tropical and geographical medicine. McGraw Hill Book Company, New York 1985; 340.
  79. Obiajuru OC, Noku AJ, Ogbulie JN. Studies on the prevalence of sexually transmitted diseases amongst teenage hawkers in Lmo state Nigeria. *Int J Env Health Human Develop* 2004; 4: 104-119.
  80. Fouts AC, Kraus SJ. *Trichomoniasis vaginalis*: reevaluation of its clinical presentation and laboratory diagnosis. *J Infect Dis* 1980; 141: 137-143.
  81. Arora DR, Arora B. *Medical Parasitology* (2nd Edn.). CBS. Publisher, New Delhi India 2005; 41-43.
  82. Hobbs MM, Sena AC. Modern diagnosis of *Trichomonas vaginalis* infection. *Sex Transm Infect* 2013; 89: 434-438.
  83. Guillermo M, Thomas CQ, Anne R, Kelly TM, Charlotte AG. Diagnosis of *Trichomonas vaginalis* infection by PCR using vaginal swab samples. *J Clin Microbiol* 1998; 36: 3205-3210.
  84. van Der Schee C, van Belkum A, Zwiijgers L, van Der Brugge E, Oneill EL, Luijendijk A, van Rijsoort-Vos T, van Der Meijden WI, Verbrugh H, Sluiters HJ. Improved diagnosis of *Trichomonas vaginalis* infection by PCR using vaginal swabs and urine specimens compared to diagnosis by wet mount microscopy, culture, and fluorescent staining. *J Clin Microbiol* 1999; 37: 4127-4130.
  85. Vanacova S, Tachezy J, Kulda J, Flegr J. Characterization of trichomonad species and strains by PCR fingerprinting. *J Eukaryot Microbiol* 1997; 44: 545-552.
  86. Brooker S. Estimating the global distribution and disease burden of intestinal nematode infections: adding up the numbers-a review. *Int J Parasitol* 2010; 40: 1137-1144.
  87. Bartsch SM, Hotez PJ, Asti L, Zapf KM, Bottazzi ME, Diemert DJ. The global economic and health burden of human hookworm infection. *PLoS Negl Trop Dis* 2016; 10: 0004922.
  88. Murray CJL, Ezzati M, Flaxman AD, Lim S, Lozano R, Michaud C. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; 380: 2197-223.
  89. Salomon JA, Haagsma JA, Davis A, de Noordhout CM, Polinder S, Havelaar AH. Disability weights for the Global Burden of Disease 2013 study. *Lancet Glob Health* 2015; 3: 712-723.

#### \*Correspondence to

Omorodion Oriri Asemota

Industrial Safety and Environmental Technology Department

Petroleum Training Institute

Nigeria