

Vascular endothelial growth factor (*VEGF*) rs3025039 polymorphism is associated with increased risk of osteosarcoma.

Jian Li^{1,2}, Tao Zhou², Hao Lin², Xiaoqiang Chen², Shaoqian Wang², Zongsheng Yin^{3*}

¹First Affiliated Hospital of Anhui Medical University, Hefei, Anhui, 230032, PR China

²Department of Orthopedics, Ma'an Shan People's Hospital, Ma'an Shan 243100, Anhui, PR China

³Department of Orthopedics, First Affiliated Hospital of Anhui Medical University, Hefei 230032, Anhui, PR China

Abstract

Background: The role of vascular endothelial growth factor (*VEGF*) rs3025039 polymorphism on osteosarcoma risk was not fully clear. Thus, we did a case-control study to evaluate the association between *VEGF* rs3025039 polymorphism and osteosarcoma risk.

Method: This study included 242 patients with osteosarcoma and 253 controls. The genotyping was conducted using the matrix-assisted laser desorption ionization time of flight mass spectrometry (MALDI-TOF MS).

Results: The *VEGF* rs3025039 TT genotype was significant higher in the osteosarcoma group than in the control group (OR=2.42, 95% CI 1.00-5.85, P=0.04). The TT genotype and CT genotype was also significantly associated with osteosarcoma risk (OR=1.43, 95% CI 1.00-2.05, P=0.04). Under the allelic model, T allele of rs3025039 was significantly associated with higher osteosarcoma risk (OR=1.41, 95% CI 1.05-1.90, P=0.02). In addition, *VEGF* rs3025039 TT genotype was not associated with tumor location (OR=1.02, 95% CI 0.59-1.77, P=0.94) and metastasis (OR=1.76, 95% CI 0.90-3.43, P=0.10).

Conclusion: In conclusion, this study confirmed that *VEGF* rs3025039 polymorphism was significantly associated with higher risk of osteosarcoma.

Keywords: Vascular endothelial growth factor, Osteosarcoma, Genetic.

Accepted on June 15, 2017

Introduction

Osteosarcoma originates from primitive bone-forming mesenchymal cells and has been identified as an aggressive sarcoma of the bone [1]. Osteosarcoma has a wide range of histological appearances. Conventional osteosarcoma may be classified as osteoblastic, chondroblastic or fibroblastic, depending on the predominant type of extracellular matrix present [2]. Despite combined therapy, more than 30% of the patients showed the recurrence or metastatic disease during the first five years after diagnosis [3].

Vascular endothelial growth factor (*VEGF*) plays an important role in the maintenance of endothelial integrity, endothelial survival and the physiological function of endothelium [4]. Niu et al. suggested that knockdown of *VEGFA* by siRNA inhibited proliferation, migration, and invasion of U2OS cells [5]. Peng et al. demonstrated that *VEGF* silencing could suppress cells proliferation, promote cells apoptosis and reduce osteosarcoma angiogenesis through inactivation of *VEGF/PI3K/AKT* signaling pathway [6]. Han found that *VEGF* is related to the grade and metastasis of osteosarcoma [7]. The human *VEGF* gene is located on chromosome 6p21.3 with 7 introns and 8 exons and shows polymorphism [8]. The role of

VEGF rs3025039 polymorphism on osteosarcoma risk was not fully clear. Thus, we did a case-control study to evaluate the association between *VEGF* rs3025039 polymorphism and osteosarcoma risk.

Methods

Study population

This study consisted of 242 patients with osteosarcoma. They were treated in the First Affiliated Hospital of Anhui Medical University, Ma'asshan hospital and Drum Tower Hospital between 2010 and 2017. The control group contained 253 healthy subjects who came from our hospital's physical checkup center during the same period of time. All control subjects had no history of cancer. Written permission was obtained from all the participants and the study was approved by the Research Ethics Committee of First Affiliated Hospital of Anhui Medical University.

Sample collection

Peripheral venous blood samples (10 mL) were collected from all patients using vacutainer tubes in the morning. Blood

Vascular endothelial growth factor (VEGF) rs3025039 polymorphism is associated with increased risk of osteosarcoma

samples (5 mL) for genetic analyses were transferred into tubes which contained ethylenediamine tetra-acetic acid (EDTA). Genomic DNAs were isolated using genomic DNA extraction kit (QIA amp DNA Blood Mini Kit, Qiagen, Berlin, Germany) under the manufacturer's instructions.

Genotyping method

The SNP genotyping was conducted using the matrix-assisted laser desorption ionization time of flight mass spectrometry (MALDI-TOF MS) (Sequenom Inc., San Diego, CA, USA). The PCR reaction was performed in a 20 µl reaction mixture containing 40-160 ng DNA template, 0.25 units Taq DNA polymerase (Sangon Biotech), 250 µM each deoxyribonucleotide triphosphate (dNTP) (Sangon Biotech), 0.5 µM forward primer, 0.5 µM reverse primer, and 1X PCR buffer with 1.2 mM MgCl₂. The SNaPshot assay (Applied Biosystems) was performed to confirm genotypes of 6 DNA samples. All genotyping procedures were carried out in a double-blind manner and the whole assays were proved to be reliable.

Statistical analysis

Data are presented as mean ± standard deviation (SD) or number (percentage). The chi-square test and Student's t-test were used to compare case and control groups, as appropriate. The statistical comparison between 2 groups was conducted using the t-test or the analysis of variance (ANOVA). Genotype distribution in the control group was tested by Hardy-Weinberg equilibrium (HWE). The differences in genotype and allele distribution between the patients and the controls are represented as odds ratio (OR) and 95% confidence interval (CI). P values for all tests are 2-tailed, and <0.05 was considered as statistically significant. Statistical analysis was conducted using SPSS 18 software (SPSS Inc., Chicago, IL, USA).

Results

Table 1 shows the demographic and clinical characteristics of 242 patients with osteosarcoma and 253 controls. Comparisons between the osteosarcoma group and the control group demonstrated that family history of cancer was significantly higher in patients with osteosarcoma than in controls (P<0.05). No statistical differences were seen in age and gender (all P>0.05).

Table 1. Characteristics of the cases and controls.

Characteristics	Case (n=242)	Control (n=253)	P value
Age at diagnosis	26.1 ± 16.6	28.1 ± 12.2	0.17
Sex			0.87
Male	118	122	
Female	124	131	
Family history of cancer			<0.01

Yes	198	142
No	44	111
Tumor location		
Extremities	210	
Non-extremities	32	
Metastasis		
Yes	68	
No	174	

The genotype and allele frequencies of the VEGF rs3025039 polymorphism in patients and controls are displayed in Table 2. The VEGF rs3025039 TT genotype was significant higher in the osteosarcoma group than in the control group (OR=2.42, 95% CI 1.00-5.85, P=0.04). The TT genotype and CT genotype was also significantly associated with osteosarcoma risk (OR=1.43, 95% CI 1.00-2.05, P=0.04). Under the allelic model, T allele of rs3025039 was significantly associated with higher osteosarcoma risk (OR=1.41, 95% CI 1.05-1.90, P=0.02). In addition, VEGF rs3025039 TT genotype was not associated with tumor location (OR=1.02, 95% CI 0.59-1.77, P=0.94; Table 3) and metastasis (OR=1.76, 95% CI 0.90-3.43, P=0.10; Table 3).

Table 2. The genotype and allele frequencies of VEGF polymorphism in the cases and controls.

Genotype	Case	Control	OR	95% CI	P value
CC	131	159	1.00	Reference	
CT	95	86	1.34	0.92-1.94	0.12
TT	16	8	2.42	1.00-5.85	0.04
TT+CT	111	94	1.43	1.00-2.05	0.04
C	357	404	1.00	Reference	
T	127	102	1.41	1.05-1.90	0.02

Table 3. Relation of VEGF polymorphism and characteristics.

Characteristics	TT+CT (n=111)	CC (n=94)	OR	95% CI	P value
Tumor location					
Extremities	49	42	1.00	Reference	
Non-extremities	62	52	1.02	0.59-1.77	0.94
Metastasis					
Yes	80	77	1.00	Reference	
No	31	17	1.76	0.90-3.43	0.10

Discussion

In the current study, we investigated the association between the VEGF rs3025039 polymorphism and osteosarcoma risk. The VEGF rs3025039 TT genotype was significant higher in

the osteosarcoma group than in the control group. The TT genotype and CT genotype was also significantly associated with osteosarcoma risk. Under the allelic model, T allele of rs3025039 was significantly associated with higher osteosarcoma risk. In addition, *VEGF* rs3025039 TT genotype was not associated with tumor location and metastasis.

Zhang et al. found that vascular endothelial growth factor overexpression predicted a worse prognosis for laryngeal cancer patients [9]. Zhao et al. indicated that high plasma/intratumoral *VEGF*-A level at baseline could predict poor treatment effect (depressed PFS and OS) of BEV-based chemotherapy in mCRC [10]. Cai et al. found that high level of *VEGF* is associated with poor outcomes in HCC patients treated with sorafenib [11]. Hui et al. indicated that the OS of the *VEGF*-positive group with ovarian cancer was significantly poorer than the *VEGF*-negative group [12]. Ma et al. suggested that both the *VEGF* +936C/T and -634G/C polymorphisms influence breast cancer susceptibility and tumor growth, instead of metastasis [13].

This study has some limitations. The selective bias was mostly controlled by the design of a hospital-based case-control study. As other case-control studies, this study raises concern about recall bias and residual confounding. The major difficulty is still the inability to separate exposures to factors prior to clinical onset from exposures to factors after clinical onset.

In conclusion, this study confirmed that *VEGF* rs3025039 polymorphism was significantly associated with higher risk of osteosarcoma risk.

References

- Maire G, Martin JW, Yoshimoto M, Chilton-MacNeill S, Zielenska M, Squire JA. Analysis of miRNA-gene expression-genomic profiles reveals complex mechanisms of microRNA deregulation in osteosarcoma. *Cancer Genet* 2011; 204: 138-146.
- Marina N, Gebhardt M, Teot L, Gorlick R. Biology and therapeutic advances for pediatric osteosarcoma. *Oncologist* 2004; 9: 422-441.
- Ottaviani G, Jaffe N. The epidemiology of osteosarcoma. *Cancer Treat Res* 2009; 152: 3-13.
- Ferrara N. Molecular and biological properties of vascular endothelial growth factor. *J Mol Med (Berl)* 1999; 77: 527-543.
- Niu J, Sun Y, Guo Q, Niu D, Liu B. miR-1 Inhibits Cell Growth, Migration, and Invasion by Targeting VEGFA in Osteosarcoma Cells. *Dis Markers* 2016; 2016: 7068986.
- Peng N, Gao S, Guo X, Wang G, Cheng C, Li M, Liu K. Silencing of VEGF inhibits human osteosarcoma angiogenesis and promotes cell apoptosis via VEGF/PI3K/AKT signaling pathway. *Am J Transl Res* 2016; 8: 1005-1015.
- Han G, Wang Y, Bi W, Jia J, Wang W, Xu M. Effects of vascular endothelial growth factor expression on pathological characteristics and prognosis of osteosarcoma. *Clin Exp Med* 2016; 16: 577-584.
- Arcondéguy T, Lacazette E, Millevoi S, Prats H, Touriol C. VEGF-A mRNA processing, stability and translation: a paradigm for intricate regulation of gene expression at the post-transcriptional level. *Nucleic Acids Res* 2013; 41: 7997-8010.
- Zhang LP, Chen HL. Increased vascular endothelial growth factor expression predicts a worse prognosis for laryngeal cancer patients: a meta-analysis. *J Laryngol Otol* 2017; 131: 44-50.
- Zhao L, Zhang D, Ma H, Jin M, Huang F, Zhang T. High VEGF-A level at baseline predicts poor treatment effect of bevacizumab-based chemotherapy in metastatic colorectal cancer: a meta-analysis. *Panminerva Med* 2016; 58: 48-58.
- Cao G, Li X, Qin C, Li J. Prognostic Value of VEGF in Hepatocellular Carcinoma Patients Treated with Sorafenib: A Meta-Analysis. *Med Sci Monit* 2015; 21: 3144-3151.
- Hui G, Meng M. Prognostic value of vascular endothelial growth factor expression in women with ovarian cancer: A meta-analysis. *J BUON* 2015; 20: 870-878.
- Ma J, Hu W, Zhang P, Sun Y, Wang N, Teng X, Qiao Q. The Association Between VEGF +936C/T and -634G/C Polymorphisms and Breast Cancer Susceptibility, Tumor Growth, and Metastases: Evidence From 20,728 Subjects. *Cancer Invest* 2015; 33: 312-317.

*Correspondence to

Zongsheng Yin

Department of Orthopedics

First Affiliated Hospital of Anhui Medical University

PR China