

Water soluble carbon nanotubes increase agricultural yields.

Nadeem Kizilbash^{1*}, Jamal Alruwaili¹, Abdul Hai¹, Hassan M. Khachfe² and Jaweria Ambreen³

¹Department of Medical Laboratory Technology, Faculty of Applied Medical Sciences, Northern Border University, Arar-914131, Saudi Arabia

²Lebanese Institute for Biomedical Research & Application (LIBRA), Lebanese International University, Beirut, Lebanon

³Department of Chemistry, COMSATS University Islamabad, Islamabad-45550, Pakistan

Abstract

The introduction of nanotechnology-based applications in agriculture is important to tackle the difficulties faced by agriculture and food sector across the globe. Nanomaterials facilitate controlled delivery of nutrients with increased crop protection. Due to their direct and indirect applications, nanomaterials can support the development of high-tech agricultural farms. This is exactly what is needed to maximize agricultural output by improvement of the ability of plants to absorb nutrients from the soil.

Keywords: Nanotechnology, Nanomaterials, Agriculture farms.

Accepted on April 10, 2020

Introduction

Carbon Nanotubes (CNTs) were first discovered independently by Iijima & Ichihashi and Bethune et al. [1,2]. They are cylindrical in shape and can be either single-walled (SWCNTs) or multi-walled (MWCNTs) [3]. As their name, 'Carbon Nanotubes,' implies, CNTs have diameters in the range of nanometers. Single-walled carbon nanotubes (SWCNTs) are less than 1 nm in diameter while multi-walled carbon nanotubes (MWCNTs) have diameters in excess of 100 nm. Recent studies have shown the beneficial effects of CNTs on plants, both in vitro and in vivo [4]. These studies have shown that both MWCNTs [5] and SWCNTs can cross the cell walls of plant cells and can be transported to specific cellular organelles [6, 7].

Production of Water-soluble CNTs

One of the drawbacks pertaining to the use of CNTs in agriculture is their poor water solubility. Numerous approaches have been used to improve the solubility of CNTs in water [8]. In recent years, chemical modifications have improved the solubility of CNTs in water [9]. CNTs can be modified both covalently and non-covalently by introducing various functional groups on their outer surfaces [10,11]. One convenient method is the ozone-induced oxidation of CNTs in the gas phase [12]. In addition, the UV-ozone oxidative approach has also been used [13]. Another method for surface modification of CNTs uses air atmospheric pressure dielectric barrier discharge (APDBD) [14,16].

Advantages for the Use of Carbon Nanotubes in Agriculture

A number of different CNTs with differing sizes and properties have been used in plant systems [17]. They have been shown to influence both growth and development. The presence of CNTs in the soil can induce changes in metabolic functions of plants

resulting in an increase in biomass and yield. Some studies have shown the effect of MWCNTs on seed germination and growth in radish (*Raphanus sativus*), rapeseed (*Brassica napus*), rye, lettuce, maize, and cucumber [18,19]. MWCNTs have been shown to have a positive effect on seed germination of *Brassica juncea L.* and *Phaseolus mungo L.* [20]. Similarly, Ma et al. investigated the effect that SWCNTs have on root growth of carrot (*Daucus carota*), cucumber, lettuce, onion (*Allium cepa*), and tomato [21,22]. Srinivasan found an increase in seed germination and growth rate in tomato seeds when exposed to CNTs [22]. A higher rate of germination was observed in hybrid Bt cotton (*Gossypium hirsutum*) [23], Indian mustard (*Brassica juncea*), urad bean (*Vigna mungo*), and rice (*Oryza sativa*) after treatment with MWCNTs [18,22,24]. In short, CNTs have been shown to be beneficial in agriculture as well as having possible use in other disciplines (Figure 1).




Figure 1: Various applications of Carbon Nanotubes.

Many studies have investigated the accumulation of CNTs in plants (Table 1). In 2007, Lin and Xing found that ryegrass (*Lolium perenne*) increases in root length on exposure to MWCNTs. Canas et al. investigated the effect of uncoated-SWCNTs on cabbage (*Brassica oleracea*), carrot (*Daucus carota*), cucumber (*Cucumis sativus*), lettuce (*Lactuca sativa*),

onion (*Allium cepa*) and tomato (*Lycopersicon esculentum*) [25]. CNTs have also been shown to more than double the growth rate of tobacco cell cultures [3]. A summary of the effect of CNTs on seed germination and plant development is shown in (Table 1).

Table 1: Effect of CNTs on seed germination and plant development.

Crops	Type of nanoparticles	Concentration (mg/L)	Effect
Wheat, maize, peanut, garlic, bulb	Water soluble MWCNTs	20,50	Enhanced germination
Barley, maize, soybean	MWCNTs	50,100,200	Enhanced germination
Barley, maize, soybean	MWCNTs	25,50,100 (spray)	At 25 mgL no effect; at 100 mgL increased germination rate
Tomato	MWCNTs	10,20,40	Increased seed moisture content; enhanced germination
Rice	CNTs	50,100,150	Enhanced germination
Rice	CNTs	n.a.	Increased seed moisture content; enhanced germination
Tomato, onion, turnip, radish	CNTs	10,20,40	Enhanced germination of tomato and onion

Production of Water Soluble Carbon Nanotubes-added Biofertilizer

Biofertilizers provide nutrients to the crops through nitrogen fixation and solubilizing of phosphates by the use of microbial inoculants. The microbes that are utilized in biofertilizer are first characterized in the laboratory. The steps involved in the production of biofertilizer include selection of a suitable bacterial strain, isolation of the bacteria and preparation of a carrier system followed by mixing, curing, packaging, storage and dispatching of product (Figure 2).




Figure 2: Flow chart showing the different steps involved in the commercial production of CNT-added Biofertilizer.

Mechanism of Action of Water Soluble Carbon Nanotubes

Most CNTs can be absorbed by plants [3]. However, the extent of their absorption depends on the plant species and the properties of CNTs being used [26]. The influence of CNTs can be neutral, positive or negative depending upon the concentration, chemical properties, structure and size of CNTs [27]. In some plants, biological pathways are affected by the

interaction of CNTs at the gene level. The production of fruit is significantly accelerated by MWCNTs [28] but the absorption of MWCNTs can also lead to formation of vacuoles, shrinkage of cytoplasm and non-appearance of nuclei in some plants [29,30]. It has been hypothesized that water soluble CNTs enhance the capillary action of water absorption [31]. It is already known that there is water present in xylem channels. The molecular transport takes place faster via xylem when water soluble CNT are present in water which results in very rapid flow of water, as predicted by molecular dynamics simulations [32].

Drawbacks for the use of Carbon Nanotubes

Some studies have provided information about toxicity possessed by CNTs [33]. However, very few studies have investigated the toxicity of CNTs in environmental conditions. One of the reasons why the toxicity displayed by CNTs is less understood is the low solubility and reactivity of CNTs with soil and background radiation from residual carbon present in the soil. Also, little information is available about the nutritional effects and genetic modifications induced by CNTs [34]. At high concentrations, CNTs have been found to have a detrimental effect on plant growth, while at lower levels they exhibit a beneficial effect or no effect at all [34]. A common mechanism explaining the toxicity caused by CNTs is the production of oxidative stress by formation of reactive oxygen species (ROS). The toxicity caused by MWCNTs includes histone modifications and DNA methylation [35]. On exposure to CNTs, plant cells experience damage to cell membrane, loss of mitochondrial dehydrogenase activity and production of ROS [35].

Conclusions

CNTs can be readily absorbed by plants which result in their internal translocation. The CNT-added biofertilizer provides large surface area, high reactivity, compatible pore size and

particle morphology which helps in providing nutrients to plants for improving their growth and yields. A positive physiological response is displayed by plants when CNTs are administered at low concentrations. At higher concentrations, usually detrimental effects on plant growth are observed. The effect of CNTs on the surrounding environment also needs to be further investigated before their widespread use in agriculture and plant protection can be recommended.

References

- Bethune DS, Kiang CH, De Vries G, et al. Cobalt-catalysed growth of carbon nanotubes with single-atomic-layer walls. *Nature* 1993;363(6430):605-607.
- Khodakovskaya MV, Kanishka de Silva, Alexandru SB, et al. Carbon nanotubes induce growth enhancement of tobacco cells. *ACS Nano*. 2012;6 (3):2128-2135.
- Yang W, Ratinac KR, Ringer SP, et al. Carbon nanomaterials in biosensors: should you use nanotubes or graphene? *Angew. Chem. Int. Ed.* 2010;49:2114-2138.
- Serag MF, Kaji N, Tokeshi M, et al. Carbon nanotubes and modern nanoagriculture. *Nanotechnology and Plant Sciences*. 2015;183-201.
- Lin S, Reppert J, Hu Q, et al. Uptake, translocation, and transmission of carbon nanomaterials in rice plants. *Small* 2009;5:1128-1132.
- Samaj J, Baluska F, Voigt B, et al. Endocytosis, actin cytoskeleton, and signaling. *Plant Physiol*. 2004;135:1150-1161.
- Liu Q, Chen B, Wang Q, et al. Carbon nanotubes as molecular transporters for walled plant cells. *Nano Lett*. 2009;9:1007-1010.
- Adenuga AA, Truong L, Tanguay RL et al. Preparation of water soluble carbon nanotubes and assessment of their biological activity in embryonic zebrafish. *Int J Biomed NanosciNanotechnol*. 2013;3(1-2):38-51.
- Li X, Fan Y, Watari F. Current investigations into carbon nanotubes for biomedical application. *Biomed Mater*. 2010;5(2):022001-12.
- Wu HC, Chang X, Liu L, et al. Chemistry of carbon nanotubes in biomedical applications. *J Mater Chem*. 2010;20:1036-1052.
- Parisi C, Vigani M, Rodríguez Cerezo E. Proceedings of a workshop on Nanotechnology for the agricultural sector: from research to the field Luxembourg. 2014.
- Khot LR, Sankaran S, Maja JM, et al. Applications of nanomaterials in agricultural production and crop protection: a review. *Crop Prot*. 2012;35:64-70.
- Hong, J, Peralta Videa JR, GardeaTorresdey JL. Nanomaterials in agricultural production: benefits and possible threats? In: Shamim N, Sharma VK, editors. *Sustainable nanotechnology and the environment: advances and achievements*. Washington, D.C: American Chemical Society. 2013;73-90.
- Gogos A, Knauer K, Bucheli TD. Nanomaterials in plant protection and fertilization: current state, foreseen applications, and research priorities. *J Agric Food Chem*. 2012;60:9781-9792.
- Sarlak N, Taherifar A, Salehi F. Synthesis of nanopesticides by encapsulating pesticide nanoparticles using functionalized carbon nanotubes and application of new nanocomposite for plant disease treatment. *J Agric Food Chem*. 2014;62:4833-4838.
- Suresh Kumar RS, Shiny PJ, Anjali CH, et al. Distinctive effects of nanosizedpermethrin in the environment. *Environ SciPollut Res Int*. 2013;20:2593-602.
- Lin D, Xing B. Phytotoxicity of nanoparticles: inhibition of seed germination and root growth. *Environ Pollut*. 2007;150(2):243-250.
- Pereira A, Grillo R, Mello NF, et al. Application of poly(epsilon-caprolactone) nanoparticles containing atrazine herbicide as an alternative technique to control weeds and reduce damage to the environment. *J Hazard Mater*. 2014;268:207-215.
- Ghodake G, Seo YD, Park D, et al. Phytotoxicity of carbon nanotubes assessed by *Brassica juncea* and *Phaseolusmungo*. *J NanoelectronOptoelectron*. 2010;5(2): 157-160.
- Ma X, Geiser Lee J, Deng Y, et al. Interactions between engineered nanoparticles (ENPs) and plants: phytotoxicity, uptake and accumulation. *Sci Total Environ*. 2010;408(16): 3053-3061.
- Srinivasan C, Saraswathi R. Nano-agriculture-carbon nanotubes enhance tomato seed germination and plant growth. *Curr Sci*. 2010;99:273-275.
- Nalwade AR, Neharkar SB. Carbon nanotubes enhance the growth and yield of hybrid Bt cotton Var. ACH-177-2. *Int J AdvSciTechnol Res*. 2013;3:840-846.
- Morla S, RamachandraRao CSV, Chakrapani R. Factors affecting seed germination and seedling growth of tomato plants cultured in vitro conditions. *J ChemBiol Phys Sci B*. 2011;1:328-334.
- Canas, JE, Long M, Nations S, et al. Effects of functionalized and nonfunctionalized single-walled carbon nanotubes on root elongation of select crop species. *Environ Toxicol Chem*. 2008;27(9):1922-1931.
- Pandey K, Mohamed HL, Victoria KH et al. Effects of carbon-based nanomaterials on seed germination, biomass accumulation and salt stress response of bioenergy crops. *PLoS One*. 2018;13(8):e0202274.
- Georgakilas V, Perman JA, Tucek J et al. Broad of carbon nanoallotropes: classification, chemistry, and applications of fullerenes, carbon dots, nanotubes, graphene, nanodiamonds, and combined superstructures. *Chem. Rev*. 2015;115(10):4744-4822.
- Line C, Larue C, Flahaut E. Carbon nanotubes: Impacts and behaviour in the terrestrial ecosystem - A review. *Carbon* 2017;123:767-785.
- McGehee DL, Lahiani MH, Irin F et al. Multiwalled Carbon Nanotubes Dramatically Affect the Fruit Metabolome of Exposed Tomato Plants. *ACS Appl Mater Interfaces*. 2017;9(38):32430-32435.

Citation: Nadeem Kizilbash, Jamal Alruwaili, Abdul Hai, Hassan M. Khachfe and Jaweria Ambreen. Water soluble carbon nanotubes increase agricultural yields. *J Biochem Biotech* 2020;3(1):*{pages}*.

29. Ghosh M, Chakraborty A, Bandyopadhyay M, et al. Multi-walled carbon nanotubes (MWCNT): induction of DNA damage in plant and mammalian cells. *J. Hazard. Mater.* 2011;197:327-336.
30. Kalra A, Garde S, Hummer G. Osmotic water transport through carbon nanotube membrane. 2003;PNAS100(18): 10175-10180.
31. Hummer G, Rasaiah JC, Noworyta JP. Water conduction through the hydrophobic channel of a carbon nanotubes. *Nature.* 2001;414:188-190.
32. Ghosh M, Bhadra S, Adegoke A, et al. MWCNT uptake in *Allium cepa* root cells induces cytotoxic and genotoxic responses and results in DNA hyper-methylation. *Mutat. Res.* 2015;774:49-58.
33. Verma SK, Ashok KD, Saikat G, et al. Applications of carbon nanomaterials in the plant system: A perspective view on the pros and cons. *Sci Total Environ.* 2019;667:485-499.
34. Viswanathan C, Jian-Kang Z. Epigenetic regulation of stress responses in plants. *Curr Opin Plant Biol.* 2009;12(2): 133-139.

***Correspondence to:**

Nadeem Kizilbash
Department of Medical Laboratory Technology
Faculty of Applied Medical Sciences
Northern Border University
Arar-914131, Saudi Arabia
E-mail: fsd707@gmail.com