

7th International Conference and Exhibition on
PHARMACOLOGY AND ETHNOPHARMACOLOGY
&
**5th GLOBAL PHYSIOTHERAPY, PHYSICAL
REHABILITATION AND SPORTS MEDICINE**

March 27-28, 2019 | Amsterdam, Netherlands

Antonios Trochopoulos et al., Asian J Biomed Pharmaceut Sci 2019, Volume 9 | DOI: 10.4066/2249-622X-C1-018

NEW CURCUMIN FORMULATION FOR THE TREATMENT OF CUTANEOUS T-CELL LYMPHOMA

**Antonios Trochopoulos^{1,3}, Mirela Marinova^{1,3}, Maya Zaharieva², Krassimira Yoncheva¹,
Martin R Berger³ and Spiro Konstantinov¹**

¹Medical University of Sofia, Bulgaria

²Bulgarian Academy of Sciences, Bulgaria,

³German Cancer Research Centre, Germany

Cutaneous T-cell lymphomas (CTCL) are a group of heterogeneous life-threatening extranodal T-cell lymphoproliferative neoplasms, of which Mycosis Fungoides (MF) and Sézary syndrome (Séz) are the most prominent subtypes. Taking into account the acquired resistance of malignancies in general, curcumin, a natural pigment with proven antineoplastic effect and insignificant toxicity, could serve as a therapeutic agent in combination regimes. Moreover, with inflammation playing a major role in the pathogenesis of CTCL, it is clear that compounds with antineoplastic/anti-inflammatory activity like curcumin, are favorable. In this study, we tested the antitumor efficacy of curcumin in ethanol solution, as well as incorporated into nanoparticles (mixed micelles based on Pluronic®123 and Pluronic® 127). Both forms of curcumin were tested on 3 CTCL cell lines, namely HuT-78, HH and MJ. MTT-dye reduction assay showed cytotoxic effects in all 3 CTCL cell lines for both curcumin formulations with the IC₅₀ values varying from 29.01 µM to 31.17 µM (ethanol solution) and 4.134 µM to 29.76 µM (nano-curcumin). The nano-formulation exerted faster cytotoxic effects (MTT-dye assay), which can be explained by its faster internalization into the cells as measured by fluorescent microscopy and HPLC analysis of the curcumin content in cell culture medium. Western blot analysis showed downregulation of important protein molecules regarding proliferation and survival such as: WT-1, ALK, p-JAK2, p-JAK3 for both formulations. In conclusion, in this study we compared two different curcumin formulations and, as the active ingredient is the same, we saw similar down-regulating effects on signal transduction proteins responsible for proliferation and apoptosis escape. It can be assumed that the inclusion of curcumin into polymeric nanoparticles will ameliorate its low bioavailability and support its faster internalization into the tumor cells.

BIOGRAPHY

Antonios Trochopoulos did master's in pharmacy at Medical University of Sofia, Bulgaria. In 2016 he did his PhD in department of pharmacology, toxicology, pharmacotherapy in Medical University of Sofia. In 2017-2018 he worked in German Cancer Research Center (DKFZ), Germany.

trochopoulos.antonios@outlook.com