

GLOBAL PHARMA SUMMIT

&

2nd International Conference on

GASTROENTEROLOGY AND HEPATOLOGY

November 23-24, 2018 | Bangkok, Thailand

Asian J Biomed Pharmaceut Sci 2018, Volume 8 | DOI: 10.4066/2249-622X-C5-015

ENDOSCOPIC ULTRASOUND GUIDED LEFT LOBE LIVER LESIONS BIOPSY (EUS-LLB) WITH ATYPICAL MALIGNANCIES, AN ALTERNATIVE APPROACH TO RADIOLOGICAL IMAGE GUIDED BIOPSY: INITIAL EXPERIENCE FROM A SINGLE TERTIARY CARE CENTRE

Adeel Urrehman and Mohammad Mubarak

Sindh institute of Urology and transplantation, Pakistan

Introduction: Endoscopic ultrasound (EUS) is gaining attraction as an alternative method of biopsy. It offers a more targeted approach for focal lesions in liver especially those areas which are accessible via EUS-guided method with its high diagnostic yield and limited adverse event profile making it more promising².

Methods: A total of 7 patients from Feb 2018 till September 2018 underwent EUS-guided Left Lobe lesions biopsies. All patients gave informed written consent and procedure related details were explained. Coagulation profile and preprocedural Nil per oral was similar as per standard Gastrosocopy and procedures were done under conscious sedation. EUS Guided LLB performed using 22G FNA needle, 2 passes were done with slow pull technique over one minute with 10-15 strokes in each pass to obtained core samples. All patients were discharged after 2 hrs of observation like post gastroscopy practice with nil complications. Duration of procedures ranges from 15-30 minutes.

Results: Case 1, 78-Years-old male with history of gastrectomy 5 years back for biopsy proven gastric cancer, presented with weight loss and vague epigastric pain, CT scan showed left lobe SOL. Biopsy showed well differentiated Adenocarcinoma from GI Tract. Case 2, 58-years-old male presented with weight loss and Liver mass on CT Scan. Hepatitis B & C screen was negative with normal AFP levels. Biopsy revealed Sarcomatoid Carcinoma. Case 3, 53-Years-old male presented with weight loss and abdominal pain. He had multiple lesions identified on CT scan in liver. Biopsy revealed Neuroendocrine Tumour. Case 4, 35-years-old male presented with abdominal pain. Laboratory tests showed anti-HCV reactive with normal AFP levels. CT Scan showed liver lesion suggestive of atypical Hepatocellular carcinoma. Biopsy revealed Smooth Muscle Tumour. As part of workup he also had gastroscopy and Colonoscopy with no evidence of luminal malignancy. Case 5, 60-years-old female presented with Weight loss, CT scan showed pancreatic malignancy with liver metastasis. Biopsy revealed Metastatic Adenocarcinoma. Case 6, 42-years-old female presented with Obstructive jaundice from Ampullary Carcinoma. She underwent EUS staging which revealed left lobe lesion. Biopsy revealed Metastatic Adenocarcinoma. Case 7, 32-years-old male presented with Weight loss and Obstructive jaundice, CT scan showed left lobe malignancy consistent with Cholangiocarcinoma with normal AFP and CA19-9 levels, his hepatitis B & C screen negative, biopsy revealed Lymphoproliferative disease (Lymphoma).

Conclusion: EUS-guided LLB is an alternative new technique for biopsy of liver lesions with suspected atypical malignancies. It appears to have higher level of safety and accuracy for targeted lesional biopsies. Further larger series are required for more supportive evidence.

Keywords: Endoscopic Ultrasound, Left lobe lesions, Atypical Malignancies and CT Scan.