


6th International Conference on
World Congress and Expo on

Otolaryngology: ENT Surgery

&

Cell & Stem Cell Research

September 10-11, 2018 | Paris, France



Alexander V Galazyuk

Northeast Ohio Medical University, USA

From a basic science phenomenon to a potential tinnitus treatment

Despite the ubiquity of tinnitus, its pathophysiology is poorly understood and there is no FDA approved cure or treatment. For more than 100 years, however, it has been known that a long sound stimulus can briefly eliminate or reduce tinnitus (Spaulding, 1903), a phenomenon known as residual inhibition. About 80% of patients with tinnitus describe some degree of residual inhibition, but the underlying mechanisms remain unknown. Knowledge about this natural internal ability to suppress tinnitus by a sound might shed light on brain abnormalities leading to tinnitus. More importantly, it might help to identify therapeutic strategies for tinnitus treatment. There is basic agreement among scientists that elevated spontaneous activity in the auditory system is linked to tinnitus. Our research in mice has demonstrated that neurons in the central auditory system show a suppression of their spontaneous activity immediately after presentation of a 30-second sound. The duration of this suppression is long-lasting, resembling the duration of residual inhibition in humans. There are additional striking similarities between the nature of this suppression in mice and the basic features of residual inhibition observed in tinnitus patients. These similarities strongly suggest that this suppression could be an underlying mechanism of residual inhibition.

Our most recent work is aimed to reproduce the effects of long sounds by other means. We found that a class of neurochemical


receptors-metabotropic glutamate receptors-play a key role in the sound-induced suppression of spontaneous activity. Furthermore, the drugs targeting these metabotropic glutamate receptors, administered systemically, can reversibly suppress both spontaneous activity of auditory neurons as well as tinnitus in tinnitus-positive mice, for at least two hours. We are exploring whether these drugs could provide a potential therapeutic approach for tinnitus suppression in humans.

Supported by grants R01 DC011330 and 1F31DC013498-01A1 from the National Institute on Deafness and other Communication Disorders.

Speaker Biography

Alexander V Galazyuk accepted an appointment as assistant professor at the Northeast Ohio Medical University in Rootstown, Ohio, where he is currently an associate professor of Neurobiology. He has worked in the field of auditory neuroscience throughout his career. Much of his research was focused on the brain mechanisms underlying our ability to analyse precise timing information in sounds.

e: agalaz@neomed.edu

 Notes: