

Joint Event
3rd International Conference on
Spine and Spine Disorders
&
International Conference on
Addiction Research and Therapy

November 26-27, 2018 | Dubai, UAE



Walter Bini

Waldkrankenhaus Bad Duben, Germany

Adipose derived mesenchymal stem cells in the treatment of DDD. Protocol and personal experience

Lumbar degenerative disc disease (DDD) poses an ongoing challenge as far as treatment options and alternatives, especially when considering younger patients. Over 80% of the adult population presents with one or more episodes of ongoing-progressive low-back pain (LBP). The primary cause is associated with degeneration of the intervertebral disc and which is triggered by a decrease of the nucleus pulposus cell population, as evidenced in histological studies. Definitely, in the presence of a "black disc" without profusion and neurological compromise, microsurgery or even fusion surgery should not be contemplated. Numerous percutaneous techniques have been propagated as proper way to treat this condition throughout the literature in the past years. They have been primarily focused on the treatment of the pain generated by the involved disc and the subsequent segmental insufficiency, without addressing the degeneration of the disc and for this have had limited success and remain as pain management tools. Some significant trials in the past (i.e. Chondrocyte transplantation trial) and the increasing recent research and achievements with more biological strategies as far as tissue regeneration , have motivated the development of a new treatment concept initially applicable to the lumbar spine which will be presented and discussed. Advancements have led to a significant improvement in the

understanding of the cell environment and tissue transplantation at a molecular, cellular and immunobiological level. Adipose tissue has already become a central source of clinical and research work involving adipose tissue derived progenitor cells. Endothelial and mesenchymal stem cells derived from adipose tissue are being considered and used in an array of clinical conditions and seem to have clear therapeutic benefits for many disease conditions including those affecting bone, cartilage and muscle. The use of an accessible source with abundant cells which have a high potential for regeneration clearly is superior in comparison to the chondrocyte option for the lumbar disc. Mesenchymal cells have a high self renewal capacity and a potential for multi lineage differentiation. For this, adipose tissue derived MSCs (ADMSCs) are optimal candidates for tissue regeneration and can be obtained from the patient in a one step procedure-treatment.

Speaker Biography

Walter Bini has completed his diploma at Westminster School, Simsbury Conn, USA and post-graduate degree at Universidad de Zaragoza, Facultad de Medicina, Zaragoza-Spain. In 2014, he was the Middle East Chairman of ISLASS. He was head of Neurosurgery at Sheikh Khalifa General Hospital, UAQ-UAE from 2014-2016. Currently, he is Consultant Neurosurgeon in Orthopedic Department, spine section of Lanzo Hospital COF, Lanzo d'Intelvi in Italy and also visiting consultant Neurosurgeon in Orthopedic Department at Healthpoint Hospital, UAE. Currently he is Neurosurgeon at Waldkrankenhaus Bad Duben, Germany.

e: binidr4@gmail.com



Notes: