

13th World Cancer Congress

February 25-26, 2019 | Paris, France

Coexisting Human Papilloma Virus-Type 16 & Human Herpes Virus-Type 8 in the presence of repeated electromagnetic field were found to be major contributing factor for rapid development of Cancer

Yoshiaki Omura

International College of Acupuncture and Electro-Therapeutics, USA

Our study indicated that Human Papilloma Virus-Type 16 (HPV-16) & Human Herpes Virus-Type 8 (HHV-8) in presence of repeated electromagnetic field (EMF) become major contributing factors for rapid development of cancer. Originally, the author found frequent coexistence of HPV-16 & single-cell parasite *Toxoplasma Gondii*. Eventually *Toxoplasma Gondii* are often infected with HHV-8. As potential source of this infection, author found most eggs are infected with these 2 viruses & *Toxoplasma Gondii*. Infections are limited to egg yolk while most egg whites are not infected. Many cats & dogs are also infected by these infections. Our study indicated that optimal dose of Vitamin D3 has the following 10 unique beneficial effects: 1) significant Anti-cancer effects without side effects; 2) marked decrease in DNA mutation which is proportional to 8-OH-dG; 3) marked urinary excretion of Viruses, Bacteria, Fungi, & Toxic substances, including Asbestos & toxic metals such as Hg, Pb, & Al; 4) marked increase in Acetylcholine in the brain & the rest of the body; 5) marked increase in DHEA; 6) marked decrease in β -Amyloid (1-42) in brain; 7) marked decrease in Cardiac Troponin I. 8) Marked increase in Thymosin α 1 (which has powerful, anti-cancer effects & anti-inflammatory & cancer-stimulating effects); 9) Marked increase in Thymosin β 4 (which has significant improvement in the circulatory system); 10) anti-allergic effects. Therefore,

using unique 10 beneficial effects of optimal dose of Vitamin D3 given orally every 8 hours since one optimal dose lasts average 8 hours, we can obtain significant reduction of cancer markers including Integrin α 5 β 1 (for example 1500ng or higher value in advanced cancers reduced to 0.001~0.005ng) & 8-OH-dG (60ng or higher value can be reduced to 0.1~0.15ng or less) without any side effect. Our safe, effective treatment of hopelessly advanced cancer patients with multiple metastases consist of combination of following 4 methods: 1) Individualized optimal dose of Vitamin D3 every 8 hours, 2) Selective drug uptake method by manual stimulation of organ representation area of hands, 3) Manual stimulation of newly discovered Thymus gland representation area of hands, 4) Elimination or reduction of cancer-contributing factors including viral infections of HPV-16, HHV-8, electromagnetic field, & any harmful drinks, things to eat (for example, currently available toxic white rice and toxic chemical-covered vegetables without washing with hot water several times), wear on body surface as well as optimal dose of Vitamin D3-inhibiting substances including Vitamin C over 150mg contrary to Linus Pauling's use of large dose of Vitamin C which inhibits most important 10 unique beneficial effects of optimal dose of Vitamin D3.

e: icaet@yahoo.com