

Joint Event

7th International Conference on

Otolaryngology: ENT Surgery

2

2nd International Conference on Dental Health and Oral Hygiene

September 05-06, 2019 | London, UK

Mohammad Arif Achakzai, Otolaryngology online

Volume 9 | ISSN: 2250-0359

Frequency of head & neck tumour in southern region (Balochistan) Pakistan

Mohammad Arif Achakzai

Sandeman Provincial Hospital, Pakistan

Objective: To evaluate the different types of head & Neck Tumours.

Methodology: All patients admitted in the ORL and Head & Neck Surgery Department Unit, at Sandeman provincial Hospital & Bolan Medical Complex Hospital Quetta were included in this study. The demographic and other data of the patients were recorded.

Results: The age of patients ranged from 13-70 years (mean 38.39) with 86 (55.48%) male and 69 (44.52%) Female. The metastatic squamous cell carcinoma in neck are 80 (Mean 55.17%). Some are 42 (mean 28.97%) benign, few are 13 (mean 8.97%) lymphatic origin, very few are 5 (mean 3.45%) vascular and 5 (mean 3.45%) neuronal origin.

Conclusion: The mean age of patients 38.39 and the mostly neck tumour are metastatic sqaumous cell Carcinoma.

Speaker Biography

Mohammad Arif Achakzai is the Assistant Professor in ENT Department Unit II, at the Bolan Medical University & Health Sciences, Sandeman Provincial Hospital Quetta, Pakistan. He has attended several National and International conferences across the globe. He has also published a good number of research papers in reputed journals which has a good number of citations.

e: drarifachakzai@gmail.com

