

MASS SPECTROMETRY, PROTEOMICS AND POLYMER CHEMISTRY

May 20-21, 2019 | Rome, Italy

Yuichi Negishi, J Chem Tech App 2019, Volume 3



Yuichi Negishi

Tokyo University of Science, Japan

BIOGRAPHY

Yuichi Negishi received his PhD Degree from Keio University, Japan in 2001. He is the Professor of Tokyo University of Science, Japan. He has over 150 publications that have been cited over 8,000 times. In his publications, 10 papers were categorized to Top 1% cited papers and 18 papers were selected as cover picture of the journal. His publication H-index is 45. He has been awarded several prizes, including the PCCP Prize in 2007, CSJ Award for Young Chemists in 2008, Japan Society of Molecular Science Award for Young Chemists in 2012 and Yagami Prize in 2017.

negishi@rs.kagu.tus.ac.jp

HIGH-PERFORMANCE LIQUID CHROMATOGRAPHY MASS SPECTROMETRY IN THE STUDY OF SMALL METAL CLUSTERS

Small metal clusters have attracted considerable attention as new functional nanomaterials because they have size-specific properties and functions that are not found for corresponding bulk metal. In particular, hydrophilic thiolate-protected gold clusters (hereinafter referred to as hydrophilic gold clusters) exhibit high biocompatibility and luminescence quantum yield in addition to pollution-free properties. Therefore, hydrophilic gold clusters are expected to be used in biomedical and environmental applications. Replacing some of the Au atoms in these clusters with different elements may impart them with even more useful functions. However, the synthesis of hydrophilic metal clusters has been less studied because of the complexity involved in evaluating the mass distributions of product mixtures. In this work, we found two hydrophilic interaction liquid chromatography (HILIC) columns for high-performance liquid chromatography (HPLC) suitable for the high-resolution separation of hydrophilic metal clusters. The mass distributions of the product mixtures of hydrophilic metal clusters were evaluated via HPLC mass spectrometry (LC/MS) using these HILIC columns. Consequently, we observed multiple clusters that had not been previously reported for glutathionate (SG)-protected gold clusters (AuN(SG)_M). Additionally, author demonstrated that Au_n-xM_x(SG)_M alloy clusters (M = Ag, Cu or Pd) in which part of the Au in the AuN(SG)_M cluster is replaced by a heteroelement can be synthesized, similar to the case of hydrophobic alloy clusters. It is easy to evaluate the mass distributions of hydrophilic metal clusters using this method. Thus, remarkable progress in the synthesis techniques of hydrophilic metal clusters through the use of this method is anticipated, as the situation for hydrophobic metal clusters.