

5th Global Experts Meeting on Infectious Diseases

August 08-09, 2022 | Webinar

Valentina A Divocha, J Infect Dis Med Microbiol 2022, Volume 06

Presence of cellular components in vaccines and Immunobiological the preparations.

Valentina A Divocha

Ukrainian Research Institute for Medicine Odessa, Ukraine

Now preventive maintenance of a flu by means of vaccination is conventional and is supported by experts of all world. To check presence of trypsin-like proteinase and its inhibitor in antifu and other vaccines and in immunobiological blood preparations of domestic and foreign manufacture. In work following commercial preparations have been used: "Interferon leukocytic human", "the Immunoglobulin of human placental, donor 10 %, a gonococcal vaccine a herpetic vaccine (Odessa), vaccines for preventive maintenance of a flu, a season 2002/2003 - "Influvac" which consists of hemagglutinins and a neuraminidase of a virus of a flu, strains: A/Moscow/10/99 (H3N2), A/New Caledonia/20/99 (H/N), B/Hong Kong/330/2001, "Fluarix" which consists of hemagglutinins of strains (H1N1) A/New Caledonia (H3N2), A/Panama and B/Shandong 17/97 and "Vaxigrip" which consists of three strains of a flu virus, a vaccine for preventive maintenance of a hepatitis A - "Avaxim", a blood preparation received from a heparin (the antifactor of Ha) - "Fraxiparine", a preparation from a blood of calfs for a hemodialysis - "Solcoseryl". Preparations were investigated before the termination of a period of validity.

Results: Work is devoted to study presence of components of a cell-owner and its inhibitor in vaccines and blood preparations and to define presence trypsin-like proteinase and its inhibitor in vaccines and blood preparations. It is revealed that anti influenza vaccines (influvac, vaxigrip, fluarix), herpetic and tularemic vaccines contained an inhibitor of trypsin-like proteinase in considerable quantity. Commercial preparations from a human donor blood (an immunoglobulin, interferon, fraxiparine and solcoseryl) contained as trypsin-like proteinase, and its inhibitor. The immunoglobulin contained in 4,0 times more inhibitor, than interferon.

Hence, the modern vaccines applied to prophylaxis and treatment, are insufficiently cleared. Presence of cellular components (enzymes and inhibitors) could lead to allergization and follow complication which is not very known.

Biography

Valentina Divocha in 1967 graduated from I.I. Mechnikov Odessa State University, Faculty of Biology (Department of Virology). In 1973 continued her postgraduate study at Odessa Institute of Virology and Epidemiology (specialty virology). In 1974 she was awarded her candidate degree with the thesis "Interaction of Coxsackie B viruses with sensitive cell cultures and their antigenic relationships. She has currently working as the head of the Laboratory of Experimental and Clinical Pathology for Ukrainian Research Institute of Transport Medicine, is the supervisor of the nine research programs in virology and biochemistry.

Received: July 15, 2022; **Accepted:** July 17, 2022; **Published:** July 20, 2022
