

HbA1C tracking: paving the way to better diabetes outcomes.

Suya Saini*

Department of Medicine Pt. B.D, Sharma PGIMS, Rohtak, Haryana, India

Introduction

Diabetes, a chronic metabolic condition characterized by elevated blood sugar levels, places a significant burden on those affected. Effective diabetes management requires diligent monitoring and control of blood glucose levels. Among the key markers used in diabetes care, glycated hemoglobin, commonly known as HbA1c, stands as a linchpin in understanding an individual's long-term glycemic control. In this article, we explore the significance of HbA1c, how it is measured, its clinical utility, and its crucial role in diabetes management.

The HbA1c marker

HbA1c is a term that might sound complex, but its significance is straightforward. It refers to hemoglobin that has bonded with glucose in the bloodstream. Hemoglobin, a protein found in red blood cells, is responsible for transporting oxygen throughout the body. When glucose molecules in the blood attach themselves to hemoglobin, forming glycated hemoglobin, it creates a permanent record of the average blood glucose levels over the past two to three months [1].

Measuring HbA1c

The measurement of HbA1c is a simple blood test, typically performed in a healthcare setting. It reflects the percentage of hemoglobin that has glycated, providing an estimate of average blood glucose levels over a defined period.

The American Diabetes Association recommends that individuals with diabetes should aim for an HbA1c level of less than 7% to reduce the risk of complications, although target ranges may vary depending on individual circumstances [2]. Higher HbA1c levels indicate poorer blood glucose control, while lower levels suggest better control.

Clinical utility of hba1c

HbA1c has become a cornerstone in diabetes management for several reasons

Long-term view: Unlike daily glucose monitoring, HbA1c offers a longer-term perspective, revealing how well blood sugar has been controlled over the previous few months [3]. This is invaluable for evaluating overall glycemic management.

Treatment adjustment: HbA1c levels guide healthcare providers in making adjustments to treatment plans. If levels

are consistently high, it may signal the need for changes in medication, lifestyle, or both.

Risk assessment: HbA1c serves as a predictor of the risk of diabetes-related complications. Higher levels are associated with an increased risk of heart disease, kidney disease, nerve damage, and other diabetes-related complications.

Motivation: For individuals with diabetes, HbA1c provides tangible feedback on the effectiveness of their efforts in managing the condition [4]. It can be a source of motivation to maintain better control.

The challenges of HbA1c

While HbA1c is an invaluable tool, it's important to acknowledge its limitations. It does not provide real-time data, and it may not reflect short-term fluctuations in blood glucose levels. Additionally, certain medical conditions and factors, such as anemia or kidney disease, can affect HbA1c measurements. These factors highlight the need for a comprehensive approach to diabetes management, which includes regular blood glucose monitoring, dietary control, physical activity, and medication as prescribed [5].

Conclusion

Glycated hemoglobin, or HbA1c, is a fundamental marker for diabetes management. It provides a valuable long-term perspective on glycemic control, guiding treatment decisions and assessing the risk of diabetes-related complications. While it's not without limitations, HbA1c is a critical component of diabetes care, offering individuals and healthcare providers an essential tool in the ongoing battle against this pervasive chronic condition. With diligent monitoring and a comprehensive approach, individuals with diabetes can use HbA1c to better manage their condition and reduce the risk of complications.

References

1. Aueyong OB. Decrease in HbA1c with the Use of a Continuous Glucose Monitoring Device (Doctoral dissertation, University of Massachusetts Global).
2. Lippi G, Targher G. Glycated hemoglobin (HbA1c): old dogmas, a new perspective?. *Clin Chem Lab Med.* 2010;48(5):609-14.
3. Gregg EW, Geiss L, Zhang P, et al. Implications of risk stratification for diabetes prevention: the case of hemoglobin A1c. *Am J Prev Med.* 2013;44(4):S375-80.

*Correspondence to: Suyas Saini, Department of Medicine Pt. B.D, Sharma PGIMS, Rohtak, Haryana, India, E-mail:suyasha022@gmail.com

Received: 04-Sep-2023, Manuscript No. AADY-23-118327; Editor assigned: 06-Sep-2023, PreQC No. AADY-23-118327(PQ); Reviewed: 20-Sep-2023, QC No. AADY-23-118327;

Revised: 22-Sep-2023, Manuscript No: AADY-23-118327(R); Published: 29-Sep-2023, DOI:10.35841/aady-7.5.169

4. Kropff J, DeVries JH. Continuous glucose monitoring, future products, and update on worldwide artificial pancreas projects. *Diabetes Technol The.* 2016;18(S2):S2-53.
5. Rajagopal L, Ganesan VI, Abdullah S, et al. Exploring the interrelationship between electrolytes, anemia, and glycosylated hemoglobin (Hba1c) levels in type 2 diabetics. *Asian J Pharm Clin Res.* 2018;11(1):251-6.