

Parasitic agents in the food chain: an overview of their prevalence and implications.

Lihua Ling*

Department of Microbiology, Faculty of Medicine Kuwait University Kuwait

Abstract

Parasitic agents are organisms that live on or within a host organism and obtain nutrients at the expense of the host. In the food chain, parasitic agents can have a significant impact on both the host organism and other organisms in the chain. Parasites can infect a wide range of host organisms, including humans, animals, and plants, and can cause a variety of diseases and health problems. In some cases, parasites can even lead to death. Parasites can be transmitted through various routes, including contaminated food and water, contact with infected animals or their waste, and through insect vectors such as mosquitoes and ticks. In addition to affecting the health of individual organisms, parasitic agents can also have broader ecological impacts. For example, they can disrupt food webs, alter species interactions, and impact ecosystem functioning. Understanding the ecology of parasitic agents is therefore important for managing their impact on both human health and ecological systems. This requires a multi-disciplinary approach that incorporates knowledge from fields such as biology, epidemiology, and ecology.

Keywords: Food chain.

Introduction

Parasites are organisms that live on or within another organism (host) and benefit from the host's resources. Parasites are found in every ecosystem and can impact the survival and reproduction of the host organism. In the food chain, parasites are an essential component, and their presence can impact the transmission of diseases among hosts, predators, and prey. The food chain network is a complex system that connects different organisms in a linear or non-linear way, and parasites can impact this network by changing the interactions among species. This essay discusses the role of parasitic agents in the food chain network, how they are transmitted, and the impact they have on the host organism [1].

Transmission of parasitic agents in the food chain network

Parasites can be transmitted in different ways, including through ingestion of contaminated food or water, sexual contact, or via a vector. In the food chain network, parasites can be transmitted from prey to predator, predator to prey, or between organisms at the same trophic level. For example, if a predator eats an infected prey, it can become infected with the same parasite [2,3]. Alternatively, a parasite can use a vector to infect a host organism. Vectors are organisms that transmit parasites from one host to another, such as mosquitoes, ticks, and fleas. For example, a mosquito can transmit the malaria parasite to a human when it feeds on their blood. In the food

chain network, a predator can act as a vector by transmitting parasites from one prey species to another [4].

Impact of parasitic agents on the host organism

Parasitic agents can impact the host organism in various ways. They can cause diseases, weaken the immune system, and reduce the host's ability to survive and reproduce. Some parasites can also alter the behavior of the host organism, making them more susceptible to predation or less likely to reproduce. For example, the lancet liver fluke, a parasite of sheep, can infect the brain of the host and alter its behavior, causing it to move towards water where it can be eaten by the next host in the parasite's life cycle [5].

Parasitic agents can also have a significant impact on the population dynamics of host organisms. For example, if a parasite infects a large proportion of a population, it can cause a decline in the population size. This can have a cascading effect on the food chain network, as predators that rely on the host organism for food may also experience a decline in their population size.

Impact of parasitic agents on the food chain network

The presence of parasitic agents can impact the interactions among species in the food chain network. For example, if a parasite infects a predator, it can reduce its ability to hunt, which can impact the population size of its prey. Alternatively, if a parasite infects a prey species, it can make them more

*Correspondence to: Lihua Ling, Department of Microbiology, Faculty of Medicine Kuwait University Kuwait. E-mail: Lihualing6754@cdc.gov

Received: 10-Mar-2023, Manuscript No. AAPDDT-23-91354; Editor assigned: 11-Mar-2023, Pre QC No. AAPDDT-23-91354(PQ); Reviewed: 22-Mar-2023, QC No. AAPDDT-23-91354; Revised: 23-Mar-2023, Manuscript No. AAPDDT-23-91354(R); Published: 31-Mar-2023, DOI: 10.35841/2591-7846-8.2.138

susceptible to predation, which can impact the population size of the predator. Parasitic agents can also impact the flow of energy and nutrients in the food chain network. For example, if a parasite infects a host organism, it can reduce its ability to acquire and use nutrients, which can impact the transfer of energy to higher trophic levels.

Conclusion

Parasitic agents are an essential component of the food chain network, and their presence can impact the transmission of diseases among hosts, predators, and prey. Parasites can be transmitted through ingestion of contaminated food or water, sexual contact, or via a vector. They can cause diseases, weaken the immune system, and reduce the host's ability to survive and reproduce. Parasitic agents can also impact the interactions among species in the food chain network, as well as the flow of energy and nutrients.

References

1. Poulin R. Parasite biodiversity revisited: frontiers and constraints. *Int J Parasitol.* 2014;44(9):581-9.
2. Dobson, A. (2017). Parasites and food webs. In *Food Webs* (pp. 259-269). Academic Press
3. Lafferty KD, Kuris AM. Parasitic castration: the evolution and ecology of body snatchers. *Trends Parasitol.* 2009;25(12):564-72.
4. Hudson PJ, Dobson AP, Lafferty KD. Is a healthy ecosystem one that is rich in parasites?. *Trends in ecology & evolution.* 2006;21(7):381-5.
5. Thompson RC, Lymbery AJ, Smith A. Parasites, emerging disease and wildlife conservation. *Int J Parasitol* 2010;40(10):1163-70.