High-resolution simulations of human brain cancer translational nano drugs delivery treatment process under synchrotron radiation.

Alireza Heidari*

Faculty of Chemistry, California South University, California, USA

Accepted on September 25, 2017

Image Article

In the following images, we present different high-resolution

simulations of human brain cancer translational Nano drugs delivery treatment process under synchrotron radiation (Figure 1) [1-7].



Figure 1. Images results of high-resolution simulations of human brain cancer translational Nano drugs delivery treatment process under synchrotron radiation [1-7].

Citation: Heidari A. High-resolution simulations of human brain cancer translational nano drugs delivery treatment process under synchrotron radiation. J Transl Res. 2017;1(1):1-3.

Furthermore, we simulated synchrotron radiation in this treatment process type according to the following plots (Figure 2) [1-7].



Figure 2. Simulation of human brain cancer translational Nano drugs delivery treatment process under synchrotron radiation.

References

- Heidari A, Brown C. Study of Composition and Morphology of Cadmium Oxide (CdO) Nanoparticles for Eliminating Cancer Cells. Journal of Nanomedicine Research. 2015;2(5):20.
- Heidari A, Brown C. Study of Surface Morphological, Phytochemical and Structural Characteristics of Rhodium (III) Oxide (Rh₂O₃) Nanoparticles. International Journal of Pharmacology. 2015;1:15-9.
- 3. Heidari A. An Experimental Biospectroscopic Study on Seminal Plasma in Determination of Semen Quality for Evaluation of Male Infertility. Int J Adv Technol. 2016;7:e007.
- Heidari A. Extraction and Preconcentration of N-Tolyl– Sulfonyl–Phosphoramid–Saeure–Dichlorid as an Anti– Cancer Drug from Plants: A Pharmacognosy Study. J Pharmacogn Nat Prod. 2016;2:e103.
- Heidari A. A Thermodynamic Study on Hydration and Dehydration of DNA and RNA–Amphiphile Complexes. J Bioeng Biomed Sci. 2016;S:006.
- Heidari A. Computational Studies on Molecular Structures and Carbonyl and Ketene Groups' Effects of Singlet and Triplet Energies of Azidoketene O=C=CH–NNN and Isocyanatoketene O=C=CH–N=C=O. J Appl Computat Math. 2016;5:e142.

 Heidari A. Study of Irradiations to Enhance the Induces the Dissociation of Hydrogen Bonds between Peptide Chains and Transition from Helix Structure to Random Coil Structure Using ATR–FTIR, Raman and ¹HNMR Spectroscopies. J Biomol Res Ther. 2016;5:e146.

*Correspondence to:

Alireza Heidari Faculty of Chemistry California South University California, USA Tel: +1 800-477-2254 E-mail: Alireza.Heidari@calsu.us