

## Pharmacological activities palmatine alkaloid compound isolated from *Gutteria friesiana* prospects for new drug development.

Soane Kaline Morais Chaves, Chistiane Mendes Feitosa\*, Felipe Pereira da Silva Santos and Joilane Alves Pereira Freire

<sup>1</sup>Universidade Federal do Piauí, Brazil

<sup>2</sup>Universidade Federal do Piauí, Brazil

### Abstract

Medicinal plants have been used for thousands of years for the treatment of numerous diseases becoming an important source in the search for new drugs. Among the substances, palmatine, arising from natural sources, an alkaloid has been investigated with regard to their pharmacological actions and has demonstrated promising activity for the treatment of various diseases highlighting Alzheimer's disease (AD). The aim of this study was to verify the state of the art and technique on palmatine with emphasis on their pharmacological activity *in vitro*, *in vivo* and *ex vivo* which makes it promising in developing new drugs, primarily for the treatment of AD. 164 studies were found using the keywords "palmatine", "palmatine", "palmatine activity" and "activity palmatine". Patent deposit was not observed. The palmatine shown promising pharmacological activities, including: anti-inflammatory, anti-depressive, anti-pyretic, among others. Thus, it was observed that the alkaloid demonstrated some important pharmacological actions, including related to the therapy of neurodegenerative diseases.

**Keywords:** Human foetal skin, Burn model, Wound healing, Human peripheral blood mononuclear cells (hPBMCs), Matrix metalloproteinases (MMPs).

Accepted on November 28, 2016

### Introduction


The medicinal plants have been used since the dawn of civilization for therapeutic purposes. The early civilizations have realized that some plants contain active principles that could be used for the treatment of diseases. Thus, nature has become a promising source of new substances for drug development [1-3].

Among the substances found in natural sources can highlight the palmatine an alkaloid and a major component of herbal preparations mainly used in traditional medicine Chinese, Korean and Indian. The palmatine can be found in various medicinal plants such as *Coptis chinensis*, *Rhizoma coptidis*, *Corydalis yanhusuo*, *Radix tinosporae*, among others [4,5].

The palmatine (Figure 1) was isolated of the plant species *Gutteria friesiana*. This species is a small tree known as "envireira" or "envira" found the Brazilian and Colombian Amazon basin and used in traditional medicine for various purposes. Studies of extracts, fractions, essential oils and compounds isolated of *Gutteria friesiana* showed activities as antitumor, antimicrobial activities and action larvicide against larvae of *Aedes aegypti* [6-10].

The palmatine has presented certain pharmacological activities that may be related, for example, their ability to interact with

proteins and nucleic acids, some of which are important in the treatment of neurodegenerative diseases [11,12].



**Figure 1.** Molecular structure of palmatine.

Neurodegenerative diseases currently affecting millions of people and the incidence of new cases is mainly related to the increase in the aging population [13,14]. According to studies, the number of people affected by major neurodegenerative diseases will increase by 50% by 2030 [15]. Alzheimer's disease (AD) is a neurodegenerative disorder with cognitive and neuropsychiatric symptoms that triggers a progressive

disability both in memory and other cognitive functions [16,17]

The DA is an important public health problem, especially in countries where there is an increase in life expectancy, as well as affect functional and socially family and society [18,19].

Several studies are being conducted to evaluate the pharmacological activities associated with palmatine alkaloid, from *in vitro* assays to *in vivo* and *ex vivo* tests. Thus the aim of this review was to assess the state of the art and technique on palmatine with emphasis on their pharmacological activities evaluated *in vitro*, *in vivo* and *ex vivo*.

## Materials and Methods

A literature about palmatine was performed through Science Direct, PubMed, LILACS, MEDLINE, Portal Capes, Web of Science, Scopus and Scielo databases. The key words used were: "palmatina", "palmatine", "atividade palmatina" and "activity palmatine". Articles published were selected during the period from 1993 to 2015. The selection of studies was based on analysis of titles and abstracts. A technological prospecting was performed in patents filed at the European Patent Office (EPO), Google patents, National Institute of Industrial Property (INPI), United States Patent and Trademark Office (USPTO) and World Intellectual Property Organization (WIPO). The period and the search terms were the same as for the scientific basis.

## Results and Discussion

The search on a scientific basis using the keyword "palmatine" resulted in obtaining 164 studies, whereas no studies were found using the term "palmatina". For the term "atividade palmatina" were no studies in scientific bases and regarding the keyword "activity palmatine" were observed 51 studies as shown in Table 1. After analysis of titles and abstracts and elimination of repeated articles were selected 18 studies that directly evaluated the pharmacological activities of palmatine. There were no claims of patent applications in selected bases. The studies found were selected those that evaluated directly the activity of the alkaloid *in vitro*, *in vivo* and *ex vivo*.

**Table 1.** Quantification of publishing on palmatine found in scientific database for each search descriptor.

Databases	Keywords			
	Palmatine	Palmatina	Atividade palmatina	Activity palmatine
LILACS	0	0	0	0
MEDLINE	7	0	0	3
Portal Capes	da 108	0	0	42
PubMed	0	0	0	0
SciELO	2	0	0	0
Science Direct	23	0	0	3

Scopus	7	0	0	0
Web of Science	17	0	0	3
Total publications	164	0	0	51

### Pharmacological activities in *in vitro* assays

There are several types and methodologies of *in vitro* assays to prove or disprove pharmacological actions. Certain *in vitro* assays were performed with palmatine and some activities were attributed to the substance from these tests (Table 2).

In *in vitro* assays that assess the ability of the substance to inhibit the enzyme acetylcholinesterase, one element for the treatment of Alzheimer's disease. Mak et al. [20] showed that the combination of palmatine and berberine resulted in inhibition synergistic *in vitro* of human recombinant AChE, the combination being a potential therapeutic strategy. Furthermore, it was observed sequestering capacity of ONOO-radicals, in other words, the antioxidant capacity with IC50 value of 28.70 micrometers [21]. The palmatine has also demonstrating other pharmacological actions such as the selective inhibition of prostate cancer cells and selective cytotoxic activity against cancer cell lines in the breast (MCF-7) and glioma cells (U251) [10,22]. Li et al. [23] suggested that some alkaloids, including the palmatine, would be responsible for the remarkable anti-inflammatory activity of *Rhizoma coptidis*.

### Pharmacological activities in *in vivo* and *ex vivo* assays

Tests *in vivo* and *ex vivo* are widely used for the elucidation of pharmacological actions of numerous compounds. As well as *in vitro* tests, certain activities *in vivo* and *ex vivo* of palmatine have been proven in the literature (Table 3).

Ning et al. [24] evaluated the effect of palmatine isolated of *Coptis chinensis* in hamsters fed high fat diet. It was observed that the compound reduced the level of serum total cholesterol (TC), triglycerides (TG) and low density lipoprotein (LDL), as well as increased excretion of TC and total bile acids in animals.

Another study showed that the alkaloid had effect on motor activity and the concentration of monoamines in brain regions of rats. The substance increased the hypomotility induced by  $\alpha$ -methyl-p-tyrosine, reserpine and 5-hydroxytryptophan and reduced hypermotility caused by L-DOPA with benserazide and p-chlorophenylalanine. The concentration of dopamine and homovanillic acid was significantly reduced in the cortex. In contrast the 5-HT levels in the cortex and 5-hydroxyindole acetic acid amounted [25].

In relation to Alzheimer's disease, certain behavioral tests and *ex vivo* tests are also suitable for evaluating the potential use of these compounds in the treatment of disease. The palmatine in doses 0.5 and 1 mg/kg, intraperitoneally, was assessed by the Morris maze test and it was found that the substance

## Hepatoprotective and Antioxidant Effect of Stem Barks Extracts: Methanolic and Aqueous of *Pseudoedrele kotschy* (Meliaceae) on Paracetamol-Induced Hepatic Damage in Rats

significantly improved learning and memory of mice. Furthermore, the substance reversed amnesia caused by scopolamine and diazepam. In *ex vivo* tests found a reduction of acetylcholinesterase activity in the brains of animals [26].

Dhingra and Bhankher [27], demonstrated the alkaloid's ability to reduce mice immobility period, subject and not subject to

unpredictable mild stress in the forced swimming test and the tail suspension, deducing an effect of antidepressant-like substance. The compound also significantly reversed the increased brain levels of catalase, lipid peroxidation, plasma nitrite and corticosteroid induced by stress.

**Table 2.** Summary of studies of *in vitro* activities of palmatine.

Specie	Part of plant	Type of extract/fraction	Purpose of the study	Results	References
-	-	-	To investigate the effect of palmatine on isometric strength in isolated strips of rat arteries	The palmatine relaxed dose-dependently the contractile response induced by phenylephrine	[28]
-	-	-	To observe the action of palmatine on chlorine colonic secretion (Cl-) in the colonic mucosa	The substance inhibits the secretion of Cl- activated by Ca <sup>2+</sup> and AMPC	[29]
Enantia chlorantha	Bark	Methanol extract	To study the activity of the alkaloid front of Trypanosoma cruzi and Leishmania infantum	The compound exhibited significant inhibitory activity against both parasites	[30]
Coptis chinensis	-	-	To evaluate the effect of palmatine in relation to NS2B-NS3 protease of West Nile virus	The alkaloid was able to inhibit the protease activity without detectable cytotoxicity	[31]
Coptis chinensis	-	-	To study the effect of palmatine on the differentiation of osteoclasts	The palmatine showed inhibitory effect on differentiation and function of osteoclast	[32]
Coptis chinensis	Rhizome	Methanol extract (n-butanol fraction)	To investigate the effect of alkaloid isolated in adipocyte differentiation by measuring the accumulation of lipids and determine the gene expression levels of markers of adipocytes	The alkaloid, including palmatine, inhibit the accumulation of lipids in the cells and reduced the expression levels of various marker genes adipocytes	[33]

Legend: (-) not reported in the study

**Table 3.** Summary of studies of *in vivo* and *ex vivo* activities of palmatine.

Specie	Part of plant	Type of extract/fraction	Purpose of the study	Results	References
Berberis spp.	Root	Alkaloid fraction	To study the effects of various alkaloids, including the palmatine in <i>in vivo</i> models	The palmatine showed anti-inflammatory, antinociceptive and antipyretic activity for certain routes of administration	[34]
Coptis chinensis	-	-	To evaluate the cytoprotective activity of the compound in fulminant hepatic failure induced by D-galactosamine (GalN)/lipopolysaccharide (LPS)	The alkaloid relieved liver injury induced GalN/LPS by modulating cytokine response and inhibition of apoptosis	[33]

Legend: (-) not reported in the study

## Conclusion

Through the survey it was observed that studies about the pharmacological and biological activities of palmatine are being carried out since 1993. During this period, they have been proven some important activities including anti-

inflammatory, antiparasitic, antipyretic action, among others. However, there is a field for new research with the substance, which can be considered promising for use in various therapies.

## Acknowledgement

We are grateful to CNPq and FAPEPI for financial support.

## References

1. Trevisan S, Viana V. Seleção de plantas com atividade anticolinesterásica para a doença de Alzheimer. *Química Nova*. 2003; 26: 301–304.
2. Badke MR, Budó MLD, Silva FM, Ressel LB. Plantas medicinais: o saber sustentado na prática do cotidiano popular, Escola Anna Nery Revista. de Enfermagem. 2011; 15: 132–139.
3. Marques THC, Santos PS, Freitas RM. Atividade anticolinesterásica e perfil químico de uma fração cromatográfica ativa do extrato etanólico das flores *Bellis perennis* L. (Asteraceae), *Química nova*. 2013; 36: 549–553.
4. Lee S, Jung E, Ho J, Jin S, Song K, Chun B. Theriogenology Sequential treatment with resveratrol-trolox improves development of porcine embryos derived from parthenogenetic activation and somatic cell nuclear transfer. *Theriogenology*. 2015; 84: 145–154.
5. Vrba J, Havlikova M, Gerhardova D, Ulrichova J. Palmatine activates AhR and upregulates CYP1A activity in HepG2 cells but not in human hepatocytes. *Toxicol In Vitro*. 2014; 28: 693–699.
6. Costa EV, Teixeira SD, Marques FA, Duarte MCT, Delarmelina C, Pinheiro MLB, Sales Maia BH. Chemical composition and antimicrobial activity of the essential oils of the Amazon *Guatterriopsis* species. *Phytochemistry*. 2008; 69: 1895–1899.
7. Costa EV, Pinheiro B, Marques FA, Lu M, Braga RM, Helena B, Maia LNS. First report of alkaloids in the genus *Guatterriopsis*. *Annonaceae*. 2009; 37: 43–45.
8. Acirole SDG, Piccoli CF, Duque JE, Costa EV, Navarrosilva MA, Marques FA, Rebelo M. Insecticidal activity of three species of *Guatterria* (Annonaceae) against *Aedes aegypti* (Diptera?: Culicidae). *Revista Colombiana de Entomología*. 2011; 37: 262–268.
9. Britto AC, de Oliveira AC, Henriques RM, Cardoso GM, Bomfim DS, Carvalho AA, Moraes MO, Pessoa C, Pinheiro ML, Costa EV, Bezerra DP. In vitro and in vivo antitumor effects of the essential oil from the leaves of *Guatterria friesiana*. *Planta Med*. 2012; 78: 409–414.
10. Costa EV, Da Cruz PEO, Pinheiro MLB, Marques FA, Ruiz ALTG, Marchetti GM, Maia BHLNS. Aporphine and tetrahydroprotoberberine alkaloids from the leaves of *Guatterria friesiana* (Annonaceae) and their cytotoxic activities. *J Brazil Chem Soc*. 2013; 24: 788–796.
11. Dumont É, Monari A. Interaction of palmatine with DNA: an environmentally controlled phototherapy drug. *J Phys Chem B*. 2015; 119: 410–419.
12. Vrba J, Papouskova B, Pyszkova M, Zatloukalova M, Ulrichova J, Vacek J. Metabolism of palmatine by human hepatocytes and recombinant cytochromes P450. *Metabolism of palmatine by human hepatocytes and recombinant cytochromes P450*. *J Pharm Biomed Anal*. 2015; 102: 193–198.
13. Farooqui T, Farooqui AA. Aging: an important factor for the pathogenesis of neurodegenerative diseases. *Mech Ageing Dev*. 2009; 130: 203–215.
14. Anderkova L, Rektorova I. Cognitive effects of repetitive transcranial magnetic stimulation in patients with neurodegenerative diseases - Clinician's perspective. *J Neurol Sci*. 2014; 339:15–25.
15. Ray Dorsey E, George BP, Leff B, Willis AW. The coming crisis: Obtaining care for the growing burden of neurodegenerative conditions. *Neurology*. 2013; 80: 1989–1996.
16. Martelli A, Martelli FP. Alterações Cerebrais e Análise Histopatológica dos Emaranhados Neurofibrilares na Doença de Alzheimer Brain Changes and Histopathological Analysis of Neurofibrillary Tangles in Alzheimer's Disease. 2012; 45–50.
17. Sereniki A, Vital MBF. A doença de Alzheimer: aspectos fisiopatológicos e farmacológicos. *Revista de Psiquiatria*. 2008; 30.
18. Trentini CM, Gonçalves MTA. Os métodos de investigação na pesquisa junto a cuidadores de idosos com a doença de Alzheimer *Psico*. 2009; 40: 308–318.
19. Forlenza OV. Tratamento farmacológico da doença de Alzheimer. *Revista de Psiquiatria Clínica*. 2005; 32: 137–148.
20. Mak S, Luk WWK, Cui W, Hu S, Tsim KWK, Han Y. Synergistic Inhibition on Acetylcholinesterase by the Combination of Berberine and Palmatine Originally Isolated from Chinese Medicinal Herbs. *J Mol Neurosci*. 2014; 53: 511–516.
21. Jung HA, Min BS, Yokozawa T, Lee JH, Kim YS, Choi JS. Anti-Alzheimer and antioxidant activities of *Coptidis Rhizoma* alkaloids. *Biol Pharmaceut Bullet*. 2009; 32: 1433–1438.
22. Hambright HG, Bath IS, Xie J, Ghosh R, Kumar AP. Palmatine inhibits growth and invasion in prostate cancer cell: Potential role for pS6/NF- $\kappa$ B/FLIP. *Mol Carcinogen*. 1–8.
23. Li J, Wang X, Luo J, Kong L. Seasonal Variation of Alkaloid Contents and Anti-Inflammatory Activity of *Rhizoma coptidis* Based on Fingerprints Combined with Chemometrics Methods. 2015; 1131–1139.
24. Ning N, He K, Wang Y, Zou Z, Wu H, Li X, Ye X. Hypolipidemic Effect and Mechanism of Palmatine from *Coptis chinensis* in Hamsters Fed High-Fat diet. *Phytother Res*. 2015; 29: 668–673.
25. Hsieh MT, Ho YF, Peng WH, Wu CR, Chen CF. Effects of *Hemerocallis flava* on motor activity and the concentration of central monoamines and its metabolites in rats. *J Ethnopharmacol*. 1993; 52: 71–76.
26. Dhingra D, Kumar V. Memory-enhancing activity of palmatine in mice using elevated plus maze and Morris water maze. *Adv Pharmacol Sci*. 2012.

*Hepatoprotective and Antioxidant Effect of Stem Barks Extracts: Methanolic and Aqueous of Pseudocedrela kotschyi (Meliaceae) on Paracetamol-Induced Hepatic Damage in Rats*

27. Dhingra D, Bhankher A. Behavioral and biochemical evidences for antidepressant-like activity of palmatine in mice subjected to chronic unpredictable mild stress. *Pharmacol Reports*. 2014; 66: 1–9.
28. Chang YL, Usami S, Hsieh MT, Jiang MJ. Effects of palmatine on isometric force and intracellular calcium levels of arterial smooth muscle. *Life Sci*. 1999; 64: 597-606.
29. Wu DZ, Yuan JY, Shi HL, Hu ZB. Palmatine, a protoberberine alkaloid, inhibits both Ca<sup>2+</sup>- and cAMP-activated Cl<sup>-</sup> secretion in isolated rat distal colon. *Br J Pharmacol*. 2008; 153: 1203–1213.
30. Nkwengoua ET, Ngantchou I, Nyasse B, Denier C, Blonski C, Schneider B. In vitro inhibitory effects of palmatine from *Enantia chlorantha* on *Trypanosoma cruzi* and *Leishmania infantum*. *Natural Product Research*. 2009; 23:1144-1150.
31. Jia F, Zou G, Fan J, Yuan Z. Identification of palmatine as an inhibitor of West Nile virus. *Arch Virol*. 2010; 155: 1325-1329.
32. Lee JW, Mase N, Yonezawa T, Seo HJ, Jeon WB, Cha BY, Woo JT. Palmatine attenuates osteoclast differentiation and function through inhibition of receptor activator of nuclear factor- $\kappa$ B ligand expression in osteoblast cells. *Biol Pharmaceutical Bull*. 2010; 33: 1733–1739.
33. Choi JS, Kim JH, Ali MY, Min BS, Kim GD, Jung HA. Coptis chinensis alkaloids exert anti-adipogenic activity on 3T3-L1 adipocytes by downregulating C/EBP- $\alpha$  and PPAR- $\gamma$ . *Fitoterapia*. 2014; 98: 199-208.
34. K peli E, Kosar M, Yesilada E, Baser KHC, Baser C. A comparative study on the anti-inflammatory, antinociceptive and antipyretic effects of isoquinoline alkaloids from the roots of Turkish Berberis species. *Life Sciences* 2002; 72: 645–657.

**\*Correspondence to**

Chistiane Mendes Feitosa  
Universidade Federal do Pia   
Brazil