

Sustainable irrigation: The role of drip irrigation systems in modern agriculture.

Guangyong Wang*

College of Water Resources and Civil Engineering, China Agricultural University, Beijing, 100083, China

Introduction

Post-harvest preservation is crucial for maintaining the quality and shelf life of agricultural products. With the increasing demand for fresh produce, it has become increasingly important to develop new and innovative techniques for preserving the freshness of fruits and vegetables. One of the most common methods of post-harvest preservation is refrigeration, which slows down the activity of enzymes and microorganisms that cause spoilage. However, this method can also lead to chilling injury, which can affect the texture, flavor, and appearance of the produce [1].

Another preservation method is the use of controlled atmosphere storage, which involves regulating the levels of oxygen and carbon dioxide in the storage area. This method can slow down the respiration of the produce, and can also help prevent the growth of microorganisms. A relatively new preservation technique is the use of modified atmosphere packaging (MAP). This method involves packaging the produce in a special film that allows for the regulation of the gases inside the package. This can help to extend the shelf life of the produce by slowing down the respiration and preventing the growth of microorganisms [2].

Another innovative preservation technique is the use of high-pressure processing (HPP). This method uses high pressure to inactivate the microorganisms that cause spoilage, while maintaining the quality and nutritional value of the produce. This method is particularly effective for preserving the freshness of juices and other liquid products. In addition to these techniques, there are also several natural preservation methods that can be used to extend the shelf life of produce. These include the use of natural antimicrobials, such as essential oils and vinegar, and the use of natural antioxidants, such as vitamin C and vitamin E [3].

In addition to the preservation methods mentioned in the previous article, there are several other techniques that can be used to extend the shelf life of fruits and vegetables. One such method is the use of waxing. This involves coating the produce with a thin layer of wax, which can help to prevent moisture loss and reduce the penetration of oxygen. This method is commonly used for fruits such as apples, pears, and citrus fruits [4].

Another technique is the use of ethylene inhibitors. Ethylene is a natural hormone produced by fruits and vegetables that can cause ripening and eventual spoilage. By using inhibitors, such as 1-MCP (1-methylcyclopropene), it is possible to block the effects of ethylene and slow down the ripening process. This can help to extend the shelf life of fruits such as bananas, avocados, and tomatoes. Irradiation is another preservation method that can be used to extend the shelf life of fruits and vegetables. This process involves exposing the produce to ionizing radiation, which can help to inactivate bacteria, viruses, and insects. Irradiation is considered safe by the World Health Organization (WHO) and the Food and Agriculture Organization (FAO) and is approved by many countries for use in food preservation. Finally, one more preservation method is the use of natural preservatives, such as honey, salt, and sugar. These natural preservatives can help to inhibit the growth of microorganisms, and can also add flavor to the produce. They have been used for centuries to preserve food, and are still commonly used today, particularly in fermented foods like pickles, sauerkraut, and kimchi [5].

Conclusion

Post-harvest preservation is a crucial aspect of ensuring the quality and shelf life of agricultural products. With the increasing demand for fresh produce, it has become increasingly important to develop new and innovative techniques for preserving the freshness of fruits and vegetables. From refrigeration, controlled atmosphere storage, modified atmosphere packaging, high-pressure processing, to natural preservation methods, there are various options available to suit different products and requirements.

References

1. Nilsson RH, Larsson KH, Taylor AF, et al. The unite database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res.* 2019;47(1):259-64.
2. Abo KA, Fred-Jaiyesimi AA, Jaiyesimi AE. Ethnobotanical studies of medicinal plants used in the management of diabetes mellitus in South Western Nigeria. *J Ethnopharmacol.* 2008;115(1):67-71.
3. Steyskal GC. Trend curves of the rate of species description in zoology. *Sci.* 1965;149(3686):880-2.

*Correspondence to: Guangyong Wang, College of Water Resources and Civil Engineering, China Agricultural University, Beijing, 100083, China, E-mail: wangyong@cau.edu.cn

Received: 28-Jan-2023, Manuscript No. AAASCB-23-88137; Editor assigned: 29-Jan-2023, PreQC No. AAASCB-23-88137(PQ); Reviewed: 12-Feb-2023, QC No. AAASCB-23-88137; Revised: 29-Mar-2023, Manuscript No. AAASCB-23-88137(R); Published: 05-Apr-2023, DOI: 10.35841/2591-7366-7.2.168

4. Wheeler QD. Taxonomic triage and the poverty of phylogeny. *Philos Trans R Soc Lond B Biol Sci.* 2004;359(1444):571-83.
5. Dauncey EA, Irving JT, Allkin R. A review of issues of nomenclature and taxonomy of *Hypericum perforatum* L. and Kew's Medicinal Plant Names Services. *J Pharm Pharmacol.* 2019;71(1):4-14.