



The Effect of Benign Paroxysmal Positional Vertigo (BPPV) in the elderly

Weaam Al-Ameer

Bilingual SLP-AUD, Saudi Arabia

Abstract

To begin with the prevalence of vestibular and vertigo symptoms are a significant problem for elderly, around 30% of vertigo present in person older than 60 years and after the age of 85 years drawing near 50%. The benign paroxysmal positional vertigo is a major common concern for seniors, due to it is false sensation of spinning. Each human ear has a three semicircular canal which are the main role of it is to support the individual balance. Inside of each canal there are a calcium carbonate crystals (otoconia) which are embedded in gel in the utricle. If the calcium carbonate crystals (otoconia) moved by any instant and traveled from the utricle into one or more of the three semicircular canals where it should not. It causes a serious episode of vertigo, provoked by movement and changes in positions. Where the inner ear is the responsible for sending signals to the brain regarding head and body movements relative to gravity. It helps keep the individual balance to do daily tasks. The symptoms of vertigo where the feeling that you or environment around you, is moving or spinning, may vary from individual to the other. The attack of the vertigo and the sensation of it, may not be noticeable and for few seconds. For other attack it could be severe and last for a long period, where the individual will not be able to keep his balance or do daily routine. Benign paroxysmal positional vertigo may be associated with age and can occur for no known reason. The treatment of benign paroxysmal positional vertigo varies depending on the cause of it. However, it can be effectively treated with the proper maneuvers performed by a qualified healthcare specialist.

Image:



(Photo by: QAI Publishing/Universal Images Group via Getty Images)

Biography

Weaam Alameer earned her bachelor's degree in speech-language-and hearing sciences from Dar Al-Hekma university in June 2017 Jeddah, KSA. She is an audiologist who has her experience in both field's speech and audiology. As an audiologist her responsibility is to implement an effective assessment, counselling and management that will help her clients to raise awareness and acquire a good quality of life. She is also passionate about creating new ways of assessment and treatment that will help the clinicians and the patients to achieve better results at every management session.

Recent Publications:

1. You, P., Instrum, R., & Parnes, L. (2018). Benign paroxysmal positional vertigo. *Laryngoscope Investigative Otolaryngology*, 4(1), 116-123. doi: 10.1002/liv.2.230
2. Strupp, M. (2009). *Basic and clinical aspects of vertigo and dizziness*. Boston, Mass.: Blackwell.
3. Balatsouras, D., Koukoutsis, G., Fassolis, A., Moukos, A., & Aspris, A. (2018). Benign paroxysmal positional vertigo in the elderly: current insights. *Clinical Interventions In Aging*, Volume 13, 2251-2266. doi: 10.2147/cia.s144134
4. Wang, W., Lin, P., Han, X., Mao, X., Li, S., & Wen, C. et al. (2020). New views on benign paroxysmal positional vertigo. *Journal Of Bio-X Research*, 3(2), 78-82. doi: 10.1097/jbr.0000000000000069
5. Pérez-Vázquez, P., & Franco-Gutiérrez, V. (2017). Treatment of benign paroxysmal positional vertigo. A clinical review. *Journal Of Otolaryngology*, 12(4), 165-173. doi: 10.1016/j.joto.2017.08.004.

11th International Conference on Otorhinolaryngology and ENT Surgery
Webinar | February 09, 2022

Citation: Weaam Al-Ameer, *The Effect of Benign Paroxysmal Positional Vertigo (BPPV) in the elderly*, *Otorhinolaryngology* 2022, 11th International Conference on Otorhinolaryngology and ENT Surgery | February 09, 2022