

## The expressions of *Caveolin-2* mRNAs, natural antisense transcripts and proteins in mouse liver development.

Mitsuru Chiba\*

Department of Bioscience and Laboratory Medicine, Graduate School of Health Sciences, Hirosaki University, 66-1 Hon-Cho, Hirosaki, Aomori, Japan

### Abstract

*Caveolin-2 (Cav2)* has been shown to be a major component of the inner surface of caveolae and be involved in lipid metabolism, development and cancer. Natural Antisense Transcripts (NATs) are transcribed from a large number of genes in various species, including humans and mice. In the present study, we investigated the expressions and localization of *Cav2* mRNAs, NATs and proteins in the mouse liver at various developmental stages. The expressions of *Cav2* mRNAs and proteins in the liver were higher at the new-born stage (NB) than at days 14, 17, and 19 of the embryonic stage (E) and at day 3 after birth. The expressions of *Cav2* NATs were lower than that of *Cav2* mRNAs at each stage. On tissues, the expressions of *Cav2* mRNAs and NATs were uniformly low in the liver at E14. The expressions of *Cav2* mRNAs were predominantly strong in the hepatocytes at NB, whereas the expressions of *Cav2* NATs were low in the hepatocytes at NB but strongly detected in specific cells. In conclusions, *Cav2* mRNAs and NATs did not co-localized in the liver at NB. These results suggest that *Cav2* NATs may be not involved in the regulation of *Cav2* mRNA expressions at NB.

**Keywords:** *Caveolin-2*, Natural antisense transcripts, Liver development, *In situ* hybridization, New-born.

Accepted on May 23, 2018

### Introduction

Caveolae are specialized domains of the plasma membrane that are implicated in the sequestration of a variety of lipid and protein molecules [1]. It has been suggested that these important cellular organelles play a pivotal role in biochemical processes such as lipid metabolism, growth regulation, signal transduction, and apoptosis [2]. Caveolin interacts with and regulates heterotrimeric G-proteins [1,2]. Currently, there are 3 members of the caveolin multi-gene family that encode 21-24 kDa integral membrane proteins that comprise the major structural component of the caveolar membrane [3]. *Caveolin-2 (Cav2)* is abundantly expressed in fibroblasts, differentiated adipocytes, smooth and skeletal muscles, and endothelial cells [4]. The expressions of *Cav1* are similar to that of *Cav2*, whereas that of *Cav3* are limited to the muscle tissues [5]. *Cav1* regulates hepatic lipid accumulation, lipid and glucose metabolism, and hepatic proliferation [6]; however, the expressions and functions of *Cav2* in the liver are unknown.

Natural antisense transcripts (NATs) are known as one of gene expression regulators in eukaryotes [7]. In the past several years, the expression of NATs in mammals is reported [8,9]. In addition, investigation that the differential expressions of several NATs are induced during the process of liver development is made [10]. Due to the increase in the number of these studies, researchers have initiated investigations of not only the expressions of mRNAs and NATs but also the

localization and regulation of their transcripts to completely understand the molecular interactions occurring in individual loci, cells, and tissues. Changes in the level of expressions of each NAT have been described in complex diseases such as cancer and neurological [11,12]. Identification of *Cav2* NATs as an up-regulated gene during liver development is done [10]. However, little is known about the expressions of *Cav2* mRNAs and NATs during pre and postnatal development of the liver. Hence, in the present study, the possible existence of *Cav2* NATs and the localization of *Cav2* mRNAs and its NATs during the development of the liver is examined.

### Materials and Methods

#### *Liver samples*

Livers at various developmental stages, embryonic stage (E) days 14, 17, and 19, new-born stage (NB), and 3-d-old stage (3D), were obtained from the C57BL/6J mice at the RIKEN BioResource Center (Tsukuba, Ibaraki, Japan). All animal experiments were carried out according to the RIKEN guidelines for the care and use of experimental animals. Livers at E14, E17, E19, NB, and 3D were collected (each n=3).

#### *Western blotting*

Liver samples were lysed using M-PER Mammalian Protein Extraction Reagent (ThermoFisher Scientific, Inc., Waltham,

MA, USA) supplemented with Halt proteinase inhibitor cocktail (ThermoFisher Scientific, Inc.). Protein concentrations were determined using the BCA Protein Assay Kit (ThermoFisher Scientific, Inc.) and Benchmark Microplate Reader (Bio-Rad Laboratories, Inc., Hercules, CA, USA) according to the manufacturer's instructions. Protein samples (10 µg) from livers at various developmental stages were prepared using 4X sample buffer (0.25 mol/l Tris-HCl, 8% SDS, 40% glycerol, 20% 2-mercaptoethanol and 0.02% bromophenol blue; pH 6.8). Electrophoresis and immunoblotting were performed by protocol reported previously [13]. Anti-Cav2 rabbit polyclonal antibody (#SP5143P, Acris Antibodies, Herford, Germany) (1:1000 dilution) and anti-β-actin (Actb) rabbit polyclonal antibody (#ab8227; Abcam, Cambridge, UK) (1:1000 dilution) were used as primary antibodies.

### Strand-specific RT-qPCR

Total RNAs from the livers was isolated using Isogen II (Nippongene, Tokyo, Japan) according to the manufacturer's instructions. The quality and concentration of the total RNAs were assessed using the NanoDrop spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA) according to the manufacturer's instructions. All the total RNA samples had 260/280 nm absorbance ratios of 1.8-2.0 [9]. To provide an external control for the comparison of results obtained from strand-specific reverse transcription-quantitative polymerase chain reaction (strand-specific RT-qPCR), an aliquot of each RNA sample was mixed with that of the RNA fragment synthesized from pEGFP-C1 vector (EGFP: enhanced green fluorescent protein; ThermoFisher Scientific, Inc.) to attain a final amount of  $5 \times 10^{-5}$  pmol/10 µg total RNA, and the resulting mixture was subjected to the synthesis of first-strand complementary DNA (cDNA) [14].

Strand-specific RT-qPCR was performed using the total RNAs obtained from livers at different developmental stages to confirm the expression pattern of Cav2 mRNAs and NATs during liver development. First-strand cDNAs derived from mRNAs or NATs were synthesized using a Cav2 reverse primer (5'-ACA AAA CAA CTA CAT GTC TAC ATG-3') or Cav2 forward primer (5'-TTT GGA ATG CAG TGT AAA ATGTG-3'), EGFP reverse primer (5'-GAA CTC CAG CAG GAC CATGT-3'), and reverse transcriptase (Promega, Madison, WI, USA) according to the manufacturer's instructions. The mixtures were incubated at 50°C for 60 min. The resulting cDNA was incubated with RNase A at 99°C for 5 min and at 37°C for 60 min to digest the RNAs. qPCR of mRNAs and NATs was performed using the first-strand cDNAs, FastStart Universal SYBR-Green Master (Roche Diagnostics, Basel, Switzerland) and primer pairs. qPCR analyses were performed by protocol reported previously [9] using the StepOne Plus Real-Time PCR system (ThermoFisher Scientific, Inc.). To determine quantity of RNA copies, standard curve method using EGFP expression was performed. The results of qPCR are presented as means ± standard errors of the samples.

### In situ hybridization

To examine the expressions of Cav2 mRNAs and NATs on the liver, 120-nucleotide probes with a specific sequence were designed from the mRNAs of Cav2 (5'-UUU GGA AUG CAG UGU AAA AUG UGC UGU CUA GGU GUU CUC UCC AAC UUA CCC AGC AAA UGU UUU AUG ACU GGG CUU CAU CUG GCU GUG ACC AUA AGA CAU GUA GAC AUG UAG UUG UUU UGU-3'). As a negative-control RNA probe (5'-UGA CGG ACA UCG GGA AAC GCC AAA GGA GAU UAU GUA CCG AGG AAG AAU GUC GCU GGA CGG UAU GCA GGA AAA GGA GGA CGU GUG GCG AGA CAG CGA CGA AGU AUC ACC GAC AUA AUC UGC-3'), a 120-nucleotide λ-phage sequence, which had no similarity with any of the mammalian sequences registered in the DNA Data Bank of Japan ([www.ddbj.nig.ac.jp/index-j.html](http://www.ddbj.nig.ac.jp/index-j.html)), was used in all *in situ* hybridization experiments to verify that the hybridization system did not emit any non-specific hybridization signals. Digoxigenin (DIG)-labeled cRNA probes were provided by Tsukuba GeneTech Laboratories (Ibaraki, Japan).

C57BL/6J mice at different developmental stages (E14 and NB) were obtained from the RIKEN BioResource Center (Ibaraki, Japan). For *in situ* hybridization, tissues from mice at different developmental stages were first fixed *in situ* by perfusion with 4% (w/v) ice-cold paraformaldehyde solution in phosphate-buffered saline (PBS). The resulting tissues were excised and further fixed overnight in paraformaldehyde solution. The fixed tissues were embedded in paraffin and cut into 4-µm sections. These sections were placed on glass slides and subjected to hematoxylin and eosin (HE) staining and *in situ* hybridization. *In situ* hybridization was performed using the de-paraffinized sections of embryos as described above. Hybridization, washing and detection were performed by protocol reported previously [15].

### Results and Discussion

I have discovered that Cav2 NATs expressed in the liver and changed during liver development and liver regeneration using microarray analysis, previously [10,16]. However, the localization and functions of Cav2 NATs in the liver have been unknown until now. Therefore, the expressions of Cav2 mRNAs, NATs and proteins in the liver at various developmental stages were analyzed in the present study.

Livers from C57BL/6J mice at E14, E17, E19, NB, and 3D were used for the preparation of Cav2 protein expressions. Western blotting was performed as described in the Materials and Methods section. The expressions of Cav2 proteins were high in the liver at NB and low at E14 (Figure 1A). This suggests that Cav2 proteins may have important roles in the liver at NB. Liver at foetus works mainly as a hematopoietic organ. Cav2 proteins may be involved in a part of functional exchanges in the liver.

In strand-specific RT-qPCR, we showed that the amounts of Cav2 mRNAs in the liver at NB was  $130.779 \pm 75.505 \times 10^{-5}$  pmol/10 µg total RNAs and was the highest among the


examined liver development stages (Table 1). This result is consistent with the expressions of Cav2 proteins, as described in Figure 1A. There were no significant differences between the amounts of Cav2 NATs among the examined livers, whereas the expressions of Cav2 NATs were less than that of Cav2 mRNAs at each stage (Table 1). This indicates that the expressions of Cav2 NATs are low in liver during developmental process.

**Table 1.** Amounts of Cav2 mRNAs and NATs in the liver at different developmental stages.

Stage	Amount of transcripts per 10 mg total RNA	
	Cav2 mRNAs	Cav2 NATs
	( $\times 10^{-5}$ pmol)	( $\times 10^{-5}$ pmol)
	Mean $\pm$ SEM	Mean $\pm$ SEM
E14	39.813 $\pm$ 22.986	3.067 $\pm$ 1.771
E17	66.598 $\pm$ 38.450	5.176 $\pm$ 2.988
E19	73.688 $\pm$ 42.544	2.433 $\pm$ 1.404
NB	130.779 $\pm$ 75.505	3.977 $\pm$ 2.296
3D	109.211 $\pm$ 63.053	3.672 $\pm$ 2.120

The localization of Cav2 mRNAs and NATs in the liver at NB with high mRNA expressions and at E14 with low mRNA expressions was examined by *in situ* hybridization (Figure 1B). High expressions of Cav2 mRNAs were observed in the hepatocytes at NB than that at E14. Uniform localization of Cav2 mRNAs and NATs were observed in the liver at E14. Low expressions of Cav2 mRNAs were observed in the hematopoietic cells of the liver at NB; furthermore, low expressions of Cav2 NATs were observed in the hepatocytes at NB, but the expressions of Cav2 NATs were strongly detected in specific cells. High expressions of Cav2 mRNAs were observed in the hepatocytes at NB. Moreover, no co-localization of Cav2 mRNAs and NATs were observed in the liver at NB. Taken together, these results suggest that Cav2 NATs may not be directly involved in the regulation of mRNA expressions at NB. In the previous study, it has been reported that Ncam1 mRNAs and NATs co-localized in the Purkinje cells of the cerebellum [17]. Therefore, it is suggested that the functions of Cav2 NATs are different from that of Ncam1 NATs. Cav2 mRNAs and NATs may be co-localized in organs other than the liver and be involved in the regulation of gene expressions. The expressions of Cav2 mRNAs and NATs in other organs should be examined in future study.

In conclusion, the expressions of Cav2 NATs were lower than that of Cav2 mRNAs and were not co-localized with the Cav2 mRNAs. Therefore, Cav2 NATs may not regulate Cav2 mRNA expression. In future studies, the cells expressing Cav2 NATs should be identified to obtain novel information about these NATs.



**Figure 1.** Detection of Caveolin-2 (Cav2) proteins and transcripts in the liver at different developmental stages. (A) Detection of Cav2 proteins in the liver at different developmental stages by western blotting. Cav2 proteins were detected by western blotting using an anti-Cav2 antibody. The livers at various developmental stages, embryonic days (E) 14, 17 and 19, new-born (NB) and 3-d old (3D), were used.  $\beta$ -actin (Actb) was examined as a housekeeping protein; (B) Detection of Cav2 transcripts in the liver at E14 and NB. Cav2 mRNAs and natural antisense transcripts (NATs) were detected by *in situ* hybridization (ISH) with digoxigenin (DIG)-labelled complementary antisense or sense probes. DIG-labelled negative control probes were used as negative controls. Scale bar represents 50  $\mu$ m.

## References

1. Gu X, Reagan AM, McClellan ME, Elliott MH. Caveolins and caveolae in ocular physiology and pathophysiology. *Prog Retin Eye Res* 2017; 56: 84-106.
2. Gupta R, Toufaily C, Annabi B. Caveolin and cavin family members: dual roles in cancer. *Biochimie* 2014; 107: 188-202.
3. Williams TM, Lisanti MP. The Caveolin genes: from cell biology to medicine. *Ann Med* 2004; 36: 584-595.
4. Zhu LL, Cui Y, Chang YS, Fang FD. A second protein marker of caveolae: caveolin-2. *Chin Med Sci J* 2010; 25: 119-124.
5. Galbiati F, Engelman JA, Volonte D, Zhang XL, Minetti C, Li M, Hou H, Kneitz B, Edelmann W, Lisanti MP. Caveolin-3 null mice show a loss of caveolae, changes in the microdomain distribution of the dystrophin-glycoprotein complex, and t-tubule abnormalities. *J Biol Chem* 2001; 276: 21425-21433.

6. Fernandez-Rojo MA, Ramm GA. Caveolin-1 function in liver physiology and disease. *Trends Mol Med* 2016; 22: 889-904.
7. Faghihi MA, Wahlestedt C. Regulatory roles of natural antisense transcripts. *Nat Rev Mol Cell Biol* 2009; 10: 637-643.
8. Chiba M, Kimura M, Asari S. Exosomes secreted from human colorectal cancer cell lines contain mRNAs, microRNAs and natural antisense RNAs, that can transfer into the human hepatoma HepG2 and lung cancer A549 cell lines. *Oncol Rep* 2012; 28: 1551-1558.
9. Chiba M, Miura T, Kasai K, Monzen S, Kashiwakura I, Yasue H, Nakamura T. Identification of up-regulated and down-regulated cis-natural antisense transcripts in the human B lymphoblastic cell line IM-9 after X-ray irradiation. *Mol Med Rep* 2012; 5: 1151-1157.
10. Chiba M. Differential expression of natural antisense transcripts during liver development in embryonic mice. *Biomed Rep* 2014; 2: 918-922.
11. Kohno K, Chiba M, Murata S, Pak S, Nagai K, Yamamoto M, Yanagisawa K, Kobayashi A, Yasue H, Ohkohchi N. Identification of natural antisense transcripts involved in human colorectal cancer development. *Int J Oncol* 2010; 37: 1425-1432.
12. Guo JH, Cheng HP, Yu L, Zhao S. Natural antisense transcripts of Alzheimer's disease associated genes. *DNA Seq* 2006; 17: 170-173.
13. Chiba M, Watanabe N, Watanabe M, Sakamoto M, Sato A, Fujisaki M, Kubota S, Monzen S, Maruyama A, Nanashima N, Kashiwakura I, Nakamura T. Exosomes derived from SW480 colorectal cancer cells promote cell migration in HepG2 hepatocellular cancer cells via the mitogen-activated protein kinase pathway. *Int J Oncol* 2016; 48: 305-312.
14. Kiuchi S, Yamada T, Kiyokawa N, Saito T, Fujimoto J, Yasue H. Genomic structure of swine taste receptor family 1 member 3, TAS1R3, and its expression in tissues. *Cytogenet Genome Res* 2006; 115: 51-61.
15. Chiba M, Kiyosawa H, Hiraiwa N, Ohkohchi N, Yasue H. Existence of Pink1 antisense RNAs in mouse and their localization. *Cytogenet Genome Res* 2009; 126: 259-270.
16. Chiba M, Murata S, Myronovych A, Kohno K, Hiraiwa N, Nishibori M, Yasue H, Ohkohchi N. Elevation and characteristics of Rab30 and S100a8/S100a9 expression in an early phase of liver regeneration in the mouse. *Int J Mol Med* 2011; 27:567-574.
17. Chiba M. Detection of natural antisense non-coding RNAs transcribed from Ncam1 in mice tissues at several developmental stages. *Biomed Rep* 2015; 3: 663-667.

**\*Correspondence to**

Mitsuru Chiba  
 Department of Bioscience and Laboratory Medicine  
 Graduate School of Health Sciences  
 Hirosaki University  
 66-1 Hon-Cho  
 Hirosaki  
 Aomori  
 Japan