

Understanding the risk factors for first febrile seizures in young children.

Preethi Elli*

Department of Brain and Neuroscience, Stanford University, United States

Introduction

Febrile seizures are a common type of seizure that occurs in young children, typically between the ages of 6 months and 5 years. They are triggered by a fever, and often occur during the first day of the illness. While febrile seizures are usually harmless and do not cause long-term harm to the child, they can be frightening for parents to witness. There are several risk factors that can increase a child's likelihood of experiencing a febrile seizure for the first time. In this article, we will explore these risk factors and discuss what parents can do to reduce their child's risk of having a febrile seizure [1].

Age of children between the ages of 6 months and 5 years are at the highest risk of having a febrile seizure. The risk decreases significantly after the age of 5. Family history of febrile seizures is more likely to experience them themselves. In fact, the risk is three to four times higher if a sibling or parent has had a febrile seizure. The higher the fever, the higher the risk of having a febrile seizure. However, it's important to note that not all children with high fevers will have a seizure. Certain types of illnesses, such as ear infections, upper respiratory infections, and gastroenteritis, can increase a child's risk of having a febrile seizure. Boys are slightly more likely than girls to experience a febrile seizure. Children with developmental delays or other neurological conditions may be at an increased risk of febrile seizures. While vaccinations themselves do not cause febrile seizures, some children may experience a fever as a side effect of certain vaccines, which can trigger a seizure [2].

While there is no way to completely prevent febrile seizures, there are steps parents can take to reduce their child's risk. For example, keeping the child's fever under control with acetaminophen or ibuprofen can help. It's also important to keep the child hydrated and comfortable during illness. In addition, parents should be vigilant in monitoring their child's fever and seek medical attention if it becomes too high or lasts for an extended period of time. Children with a family history of febrile seizures may benefit from close monitoring during illnesses. Febrile seizures are a common occurrence in young children, but they can be scary for parents to witness. Understanding the risk factors for the first febrile seizure can help parents take steps to reduce their child's risk. By keeping a close eye on their child's fever and seeking medical attention when necessary, parents can help ensure their child's safety and well-being. Febrile seizures are classified into two types: simple and complex. Simple febrile seizures last less than 15

minutes and do not occur more than once within 24 hours. They do not involve a specific part of the brain and do not cause any long-term harm to the child. On the other hand, complex febrile seizures last longer than 15 minutes, occur more than once in a 24-hour period, and involve a specific part of the brain. These seizures can cause long-term harm to the child and require further medical evaluation [3].

It's important to note that while febrile seizures are generally harmless, they can be an indication of a serious underlying medical condition. Children who experience a febrile seizure for the first time should be evaluated by a healthcare provider to determine the cause of the fever and rule out any other underlying conditions. While there is no definitive way to prevent febrile seizures, some studies suggest that certain lifestyle factors may play a role in reducing a child's risk. For example, a healthy diet and regular exercise can help boost a child's immune system and reduce the likelihood of infections that may trigger fevers and seizures. It's also important for parents to follow proper medication dosing instructions when giving their child fever-reducing medications, as overdosing can increase the risk of a febrile seizure. In addition, parents should ensure that their child receives all recommended vaccinations to reduce the risk of infections that can trigger fevers and seizures. In rare cases, children with frequent or prolonged febrile seizures may benefit from medication to prevent future seizures. However, this decision should be made in consultation with a healthcare provider [4].

Febrile seizures are a common occurrence in young children, but they can be frightening for parents to witness. Understanding the risk factors for the first febrile seizure and taking steps to reduce a child's risk can help ease parents' concerns. By seeking medical attention when necessary, maintaining a healthy lifestyle, and following proper medication dosing instructions, parents can help ensure their child's safety and well-being [5].

Conclusion

Febrile seizures occur commonly in young children that can be triggered by a fever and may cause great concern for parents. While they can be frightening, it's important to remember that febrile seizures are typically harmless and do not cause long-term harm to the child. Understanding the risk factors for the first febrile seizure can help parents take steps to reduce their child's risk and promote overall health and well-being. With proper care and monitoring, parents can help ensure their child's safety and comfort during illness. It's important to seek

*Correspondence to: Preethi Elli, Department of Brain and Neuroscience, Stanford University, United States, E-mail: preethieli@stanford.edu

Received: 12-Apr-2023, Manuscript No. AAJBN-23-97945; Editor assigned: 15-Apr-2023, PreQC No. AAJBN-23-97945(PQ); Reviewed: 29-Apr-2023, QC No. AAJBN-23-97945;

Revised: 03-May-2023, Manuscript No. AAJBN-23-97945(R); Published: 10-May-2023, DOI: 10.35841/aaibn-6.3.142

medical attention when necessary and follow recommended guidelines to prevent future febrile seizures.

References

1. Steering Steering Committee on Quality I Management SoFSAAoP . Febrile seizures: clinical practice guideline for the long-term management of the child with simple febrile seizures. *Pediatrics*. (2008) 121(6):1281-6.
2. Kanemoto K, Kawasaki J, Miyamoto T, et al. Interleukin (IL)1beta, IL-1alpha, and IL-1 receptor antagonist gene polymorphisms in patients with temporal lobe epilepsy. *Ann Neurol*. 2000;47(5):571-4.
3. Nakamura Y, Shi X, Numata T, Mori Y, et al. Novel HCN2 mutation contributes to febrile seizures by shifting the channel's kinetics in a temperature-dependent manner. *PLoS ONE*. 2013;8(12):e80376.
4. Santoro B, Lee JY, Englot DJ, et al. Increased seizure severity and seizure-related death in mice lacking HCN1 channels. *Epilepsia*. 2010;51(8):1624-7.
5. Hesdorffer DC, Shinnar S, Lewis DV, Moshe SL, Nordli DR, Jr, Pellock JM, et al.. Design and phenomenology of the FEBSTAT study. *Epilepsia*. 2012;53(9):1471-80.